


UC-NRLF



C 3 118 173





A GENERAL CATALOGUE
OF
NEBULÆ AND CLUSTERS OF STARS,

ARRANGED IN ORDER OF RIGHT ASCENSION AND REDUCED
TO THE COMMON EPOCH 1860.0

(WITH PRECESSIONS COMPUTED FOR THE EPOCH 1880.0).

BY

SIR JOHN F. W. HERSCHEL, BART., K.H., F.R.S.,

M.A., D.C.L., F.R.S.E., HON. M.R.I.A., F.R.A.S., F.G.S., M.C.U.P.S.,

Member of the Institute of France,

and

Correspondent or Honorary Member of the Imperial, Royal, and National Academies of Sciences of Amsterdam,
Berlin, Breslau, Brussels, Copenhagen, Göttingen, Haarlem, Massachusetts (U.S.), Modena, Moscow,

Naples, Pesth, Petersburg, Stockholm, Turin, Vienna, and Washington (U.S.);

The Italian and Helvetic Societies;

The Academies, Institutes, &c. of Albany (U.S.), Bologna, Catania, Dijon, Lausanne,

Nantes, Padua, Palermo, Rome, Venice, Utrecht, and Wilna;

The Philomathic Society of Paris; the Asiatic Society of Bengal; South African Lit. and Phil. Society; Literary and

Historical Society of Quebec; Historical Society of New York; Batavian Society of Experimental

Philosophy of Rotterdam; Royal Med.-Chir. Soc., Instit. of Civil Engineers,

and Soc. of Arts, Manufactures, and Commerce, London;

Leeds and Manchester Literary and Philosophical Societies; Geographical Society of Berlin;

Astronomical and Meteorological Society of British Guiana; &c. &c. &c.

From the PHILOSOPHICAL TRANSACTIONS.—PART I. 1864.



LONDON:

PRINTED BY TAYLOR AND FRANCIS, RED LION COURT, FLEET STREET.

1864.

15851
H6

A GENERAL CATALOGUE

NEBULAE AND CLUSTERS OF STARS

ARRANGED IN ORDER OF RIGHT ASCENSION AND REDUCED
TO THE COMMON EPOCH 1850.0

(WITH THE PROPORTIONS COMPUTED FOR THE EPOCH 1850.0)



BY JOHN F. W. HIERSCHEL, PART. R.H. F.R.S.

27915

NEW ASTROPHYSICAL OBSERVATORY, PART I. 1864

LONDON:

PRINTED BY TAYLOR AND FRANCIS, 42, COURT, FLEET STREET.

1864



PHILOSOPHICAL TRANSACTIONS.

I. Catalogue of Nebulæ and Clusters of Stars.

By Sir JOHN FREDERICK WILLIAM HERSCHEL, Bart., F.R.S.

Received October 16,—Read November 19, 1863.

Introduction.

THE study of the Nebulæ has, within the last quarter of a century, attracted much more of the attention of observers than heretofore—as well on account of the singularity of the phenomena presented by many of these objects, as in consequence of the increased optical power of the telescopes which the skill and industry of modern inventors and artists have placed within their reach. The brighter nebulæ cannot be viewed to any advantage, and the fainter cannot be seen at all, except by the aid of telescopes of large aperture; and, thanks to the exertions of Lord ROSSE, Mr. LASSELL, Messrs. NASMYTH and DE LA RUE in England, and Messrs. STEINHEIL, FOUCAULT, and PORRO in Germany and France, as regards reflecting telescopes, and to those of FRAUNHOFER, MERZ, CAUCHOIX, CLARKE, COOK, SECRETAN, ROSS, and DALLMEYER as regards refractors; instruments of abundantly sufficient optical capacity not only to repeat and verify the earlier observations, but to disclose new and more interesting features in many cases, have now come into the hands of many observers, both professional astronomers and amateurs, and may be had by any one who is willing to incur a cost which may be considered moderate when it is remembered that instruments of similar dimensions and goodness could not be obtained fifty years ago at any price. In consequence we find a continually increasing attention directed to this department of astronomy. Not to insist on the observations of the Earl of ROSSE and Mr. LASSELL with their transcendent reflectors, we find a systematic examination and review of them undertaken by M. D'ARREST in the year 1855, by the aid of a refractor of 6-feet focal length and $4\frac{1}{2}$ inches aperture in the Leipzig Observatory, whose results, consisting in the carefully determined places, by repeated observations, of about 230 nebulæ, were published in 1856, in a work entitled “Resultate aus Beobachtungen der Nebelflecken und Sternhaufen” (Erste Reihe, Leipzig). This review has since been carried on by the same excellent astronomer, with the great refractor by MERZ of 11 inches in aperture and 16-feet focus, erected in the year 1861 at the Royal Observa-

tory of Copenhagen. Again, from the Observatory of the Collegio Romano, under the direction of Signor SECCHI, have emanated many valuable observations, and from that at Harvard College, Cambridge, U.S., under the late and present Professors BOND, some of the most striking pictorial representations of particular nebulae which we possess. Neither ought a short but very valuable memoir by the late E. MASON, printed in the 7th volume of the Memoirs of the American Academy of Arts and Sciences, to be passed in silence; containing as it does a very elaborate and minute examination, and some excellent delineations of several highly interesting nebulae, particularly those in the great nebulous region of Cygnus. To M. AUWERS also we owe many accurate and valuable observations, besides a Catalogue comprising the whole series of Sir WILLIAM HERSCHEL's nebulae arranged in order of right ascension and reduced to a common epoch, of which more hereafter. Should the efforts which are now making to procure for the University of Melbourne in Australia a reflector of the first magnitude prove, as is to be hoped, successful, it is understood that one of the principal uses to which it will be devoted will be the examination and exact delineation of the numerous and wonderful objects of this class which the southern hemisphere presents.

These circumstances, but more especially the last-mentioned, render it extremely desirable to have presented in one work, without the necessity of turning over many volumes, a general catalogue of all the nebulae and clusters of stars actually known, both northern and southern, arranged in order of right ascension and reduced to a common and sufficiently advanced epoch which may serve as a general index to them, and enable an observer at once to turn his instrument on any one of them, as well as to put it in his power immediately to ascertain whether any object of this nature which he may encounter in his observations is new, or should be set down as one previously observed. For want of such a general catalogue, in fact, a great many nebulae have been, from time to time, in the 'Astronomische Nachrichten' and elsewhere, introduced to the world as new discoveries, which have since been identified with nebulae already described and well known. Many a supposed comet, too, would have been recognized at once as a nebula, had such a general catalogue been at hand, and much valuable time been thus saved to their observers in looking out for them again.

Besides these there are other considerations which have weighed with me in undertaking the task of compiling such a general catalogue. Having, in the course of my own observations, received the greatest possible assistance from the possession of a Manuscript Catalogue of all the nebulae and clusters discovered by my Father, brought to the common epoch 1800.0, and arranged in zones of 1° in breadth in polar distance, by his sister the late Miss CAROLINA L. HERSCHEL, it seemed to me nothing less than a debt of gratitude, not merely to acknowledge that assistance, but to avail myself still further of it to complete the list of his nebulae by supplying from that catalogue the places of all those nebulae among them which had escaped my own observation (a very numerous list), and by inserting from it all those places of nebulae observed by myself which were deficient in either element (of R.A. or P.D.), or in which I had reason to apprehend

greater errors than those which probably affected her results. This I have accordingly done. But to do it effectually, and at the same time to effect a thoroughly correct identification of the objects in my catalogues with those of the older series, involved, as a necessary preliminary step, the reduction to 1830·0 of the whole of her catalogue, an operation in which I received the assistance of my sons; the computations being executed for each nebula in duplicate and checked by myself, and which, taken leisurely, zone by zone, as time and circumstance permitted, proved less onerous and wearisome than might have been expected. The Catalogue thus reduced to the same epoch as my own, afforded the means of detecting and rectifying a great many errors of nomenclature in the latter. And it was in the course of this part of the inquiry, in which many cases of considerable intricacy and difficulty occurred (as will be evident on a perusal of the notes appended to this Catalogue), and in which it became necessary to recur both to the original sweeps and to a series of registered extracts from them (the nature of which will be more distinctly stated hereafter), that I learned fully to appreciate the skill, diligence, and accuracy which that indefatigable lady brought to bear on a task which only the most boundless devotion could have induced her to undertake or enabled her to accomplish.

Arrived at this stage—that is to say, the mean results of all the observations in my own Catalogues taken, and all the deficient or imperfectly observed nebulae in my Father's list supplied, as above stated, and the whole arranged, not in zones, but in general order of right ascension,—it then became necessary, in order to produce a work available for future observation, to bring the whole up to a still more advanced epoch. The work required for this purpose, calling no longer for any discussion, or collation of the original observations or registers, but being one of simple arithmetical computation from a definite formula—the Royal Society, at my application, very liberally undertook to supply, from the funds at their annual disposal, the amount necessary to procure its execution by an experienced computer (Mr. KERSCHNER, one of the occasional computists for the Royal Observatory of Greenwich). This work the Astronomer Royal most obligingly offered to superintend, affording at the same time his advice as to the general principle on which the computation should be conducted. The plan suggested by him and adopted in effect was this. Each object in the Catalogue was first roughly brought up to the year 1880 by the application of approximate precessions in R.A. and P.D. The places so obtained were then employed to compute the exact precessions in both by the usual formulæ, with coefficients for the year 1880·0, viz.

$$\text{Precession in R.A.} = 3^{\text{s}}\cdot072 + 1^{\text{s}}\cdot337 \cdot \sin \text{R.A.} \cotan \text{P.D.}$$

$$\text{Precession in P.D.} = -20''\cdot06 \cos \text{R.A.}$$

And the precessions, so calculated, were then used to bring up the places from 1830 to 1860, the epoch of the Catalogue; so that, the places being given for 1860 and the precessions for twenty years in advance, the application of those precessions to those places shall give dependable places for any year up to the year 1930, at which time the small error in excess or defect of the true precession consequent on using the fifty years'

antecedent place of the object will be exactly compensated by the further change of place in the same direction in the *subsequent* fifty. Two cases of excessive proximity to the poles, northern and southern, viz. those of the nebulae Nos. 2043 and 1652 of the present Catalogue, are excepted, the precessions changing so rapidly, and with so much deviation from uniformity, that a rigorous computation, at least in R.A., will always be necessary. In the case of No. 2043, the effect of precession in the thirty years from 1830 to 1860 has been to change the R.A. from $2^{\text{h}} 32^{\text{m}}$ to $10^{\text{h}} 8^{\text{m}}$.

This computation was completed, and a fair copy of the resulting places, arranged *de novo* in their order of R.A. for 1860, forwarded to me on the 6th of February last (1863). The nomenclature of the objects having in the interim been settled satisfactorily by myself, and a description of each nebula, from a careful comparison of all the descriptions given, prepared, it remained only to fill in the columns left blank for these and the other necessary particulars, and to complete the Catalogue by the insertion in their proper places of the places and descriptions of all such other nebulae, non-observed by either my Father or myself, similarly reduced, of which I could gather any accounts. These will be found enumerated further on in the "Explanation and arrangement of the Catalogue."

On the 23rd of February last, while engaged in this work, I received, by the kindness of the Astronomer Royal, a copy of the important work of M. AUWERS before alluded to, entitled "WILLIAM HERSCHEL'S Verzeichniss von Nebelflecken und Sternhaufen, bearbeitet von ARTHUR AUWERS. Königsberg, 1862," of whose existence this was my first notice. It contains a complete and most elaborate reduction to 1830, from the observed differences in R.A. and P.D. with known stars, recorded in the Philosophical Transactions, of all the nebulae and clusters in my Father's three Catalogues; together with a separate catalogue of all those collected by MESSIER from his own observations, or those of MECHAIN and others (101 in number), similarly reduced; another of LACAILLE's southern nebulae, and one of 50 "new nebulae," comprising nearly all those observed by other astronomers (Lord ROSSE excepted) in this hemisphere—all brought up to the same epoch.

It may be readily supposed that I lost no time in comparing my own previous work with this of M. AUWERS; the places of which having been obtained by the aid of far better and more dependable catalogues of stars, to give the true positions of the zero-points or determining stars in the differential observations, as well as of more exact precessions, and doubtless, a much more systematic process of treatment, would be entitled, *observation for observation*, to be considered as representing the original sweeps more faithfully than could be expected from my own preparatory catalogue. On the other hand, however, the Zone Catalogue from which that was derived possessed the advantage of having been deduced, not from a single difference of R.A. and P.D. between each nebula and a single determining star, but from *all* the observations of each nebula; often in many different sweeps, and in the same sweep often from more than one star; thus eliminating, no doubt, a great deal of casual error. In that catalogue, too, as in my own catalogues of 1833 and that of the southern nebulae, the individual results of each observation, or, to speak more exactly, of each differential comparison, is separately

recorded, so that any suspiciously large deviation from the mean of all may be at once noticed and traced to its origin in the sweeping books. My reduction was of course based on the means of all these (rejecting such as were obviously and grossly faulty), and might therefore, *pro tanto*, be regarded as of superior authority. This consideration, joined to that before adduced, decided me to retain those places in the present Catalogue which had been derived from this source, except in a few instances (specified in the notes) when it proved, by careful examination of the causes of discordance, that actual mistakes had been committed. And I must not omit to add that the comparison so instituted with M. AUWERS's results has led me to the detection of several grave errors in my own work which would certainly have otherwise escaped notice (and in some cases have caused the loss of future observations by missetting the telescope), and whose rectification has added materially to its value. On the other hand, as no human work is perfect, I have been led to notice some errors in M. AUWERS's work itself, which are set down in a list of *errata* and *corrigenda* at the end of this Catalogue; and besides, a good many cases in which, owing to mistakes in the printed catalogue in the volumes of the Philosophical Transactions (many of which stood corrected in MS. in the margin of the copy of those Transactions in my possession, and many more have been silently detected and rectified by Miss C. H. in her subsequent computations), his calculations have been founded on erroneous data, and have therefore led him to assign erroneous places to the objects so affected. Thus on every account the result has been what may be considered a complete expurgation of both our catalogues.

It remains for me to say a few words on the way in which the reduction to 1860 and the calculation of the precessions have been performed by Mr. KERSCHNER, the computist employed by the Astronomer Royal for that purpose. The whole work has been executed on printed forms, which being preserved may at any time be referred to. Since error in computation, however practised the computer, and however checked, is always possible, and occasional error of copying, especially when the order of the entries has to be rearranged, is absolutely unavoidable, I considered it incumbent on me to recalculate, *seriatim* from my original MS. Catalogue for 1830, and taking for granted the precessions set down in the fair copy, for 1880, the places both in R.A. and P.D. of every object included in the Catalogue; keeping an eye meanwhile to the precessions themselves, and their signs, to seize the least indication of error in that quarter. It would have been too laborious to recompute these. As for the precessions in P.D., their regular progression of itself ensures their correctness, as far at least as the integer seconds and the first decimal place. A pretty considerable number of errors (most of them of little moment) was thus detected and corrected—not more, however, than might reasonably be expected in the work of the most expert computist in so extensive a work, consisting of between nine and ten thousand computed entries (taking both elements), and traceable moreover in many instances to obvious misreading, and in some to actual misentry on my part, of figures in the original MS., which but for this further examination would also have escaped notice altogether.

The correction of these and the other errors already spoken of necessitated, in a great

many instances, a change in the order of R.A., and a consequent erasure and interlineation in the MS. The introduction, too, of the other nebulae (those of M. AUWERS's catalogue of "novæ," those communicated to me for insertion by M. D'ARREST, and those noticed by Lord ROSSE in his memoir of 1861, amounting altogether to 433 objects) necessitated many more interlineations, often occurring very inconveniently, two or three together, in a way to disfigure the MS. considerably. Unfortunately, too, in the MS. itself the column headed "No. in the Catalogue," which I had intended to have been left blank till all the rest of the work was completed, had been filled in by the transcriber with a series of numbers in regular progression, from 1 to 4629, the actual number of lines of which it then consisted. This made it necessary to renumber the whole *ab initio* in red ink, striking out the former numbers, and thus producing a still more unsightly appearance. Under these circumstances, I debated whether or not to recopy the whole. But, to say nothing of the sacrifice of time (since I could have entrusted it to no other hand), I believe it impossible to copy so voluminous a mass of figures and abbreviated writing without numerous errors. And being satisfied, from the repeated and careful revision it has undergone, of its present correctness, and equally so that with ordinary care on the part of the compositor (should the Council of the Royal Society decide on printing it) no *mistake* can arise from any of the alterations and interlineations it contains, I have decided in favour of presenting it as it stands, with the exception of two sheets which it was absolutely necessary to recopy owing to the extreme closeness of the interlineations, the smallness of the writing, and the transpositions needed. These have each been twice carefully read with the original.

In presenting to the Royal Society this Catalogue, it will be accompanied by the following series of records and documents which it may become desirable hereafter to refer to in elucidation of any point which may arise respecting the history or reduction of such of the objects as occur in my Father's classes and numbers printed in the Philosophical Transactions, viz.—

1st. A series of "*register sheets*," in which are entered up *all* the observations of *each* nebula or cluster copied verbatim from the sweeps, the nebulae, &c. being arranged in the order of their dates of discovery. These are the "register sheets" referred to in the notes on this Catalogue, and cited by their *general* (*i. e.* current) *number*, as H, 1; H, 2; ... H, 2508.

2nd. A similar set of register sheets of all the observations of each of MESSIER's nebulae, arranged according to MESSIER's numbers.

3rd. A general index of the 2508 nebulae in classes and numbers, to find the "general number" of each to facilitate reference to the register sheets. (This index was drawn up by myself.)

4th. An index list of the same nebulae, &c. arranged according to the "general number," to find the class and number of each.

5th. A more complete ditto ditto, containing also the rough approximate R.A. and P.D. of each object for 1800, and the determining stars as in the Philosophical Transactions.

6th. A catalogue in zones of P.D. of all the said nebulae and clusters arranged in each

zone in order of R.A., and reduced to the year 1800 by Miss C. L. HERSCHEL, exhibiting the reduced result of each separate observation of each nebula; together with the determining star or stars in each case, and the differences of R.A. and P.D. from such star, with references to the current number of the sweep in which the observation is contained.

7th. The original sweeps with the 20-feet reflector at Slough in which the nebulæ were observed, contained in three small quarto and four folio volumes of MS.

All these manuscripts, with exception of the index No. 3, are in the original hand-writings of my Father and his Sister, in most cases easily distinguishable, in some others not so readily. The Zone Catalogue No. 6 is entirely the autograph of the latter.

Explanation and arrangement of the Catalogue.

The Catalogue is arranged in twelve columns, of which the first contains the general or current number in order from 1 up to 5063, the total number of objects comprised, including six supplementary ones, whose insertion in their proper order in R.A. would have involved altering all the numbering both of the catalogue and the annotations, &c., and would have proved a source of confusion and unavoidable error. Nevertheless, to prevent their being overlooked by any observer who may consult the catalogue for the purpose of a general review of the nebulæ, or for the verification of a new one, their numbers are interpolated into the general series so as to catch the eye, and a reference made to the supplementary catalogue in each case in the column of descriptions.

Column 2 contains the numbers of those nebulæ of which observations are given in my two former catalogues, and those of the two nubeculæ; the numbers from 1 to 2307 inclusive being from that in Philosophical Transactions 1833, and from 2308 to 4021 from my Cape observations. Where a number in this column is enclosed in hooks thus [], it is taken from the Catalogue of Objects in the Nubecula minor in pp. 153 to 155 of that work. Where in parentheses thus (), from those in the Nubecula major, pp. 156 to 163.

Column 3 contains the classes and numbers of nebulæ as given by my Father in his three Catalogues in the Philosophical Transactions for 1786, 1789, and 1802. One only is omitted, viz. V. 35. It is an immense diffused nebosity, extending from $5^h 27^m$ to $5^h 42^m$ in R.A., and from $98^\circ 6'$ to $87^\circ 43'$ in P.D. A special list of these great diffusions of nebula is given by M. AUWERS in p. 42 of the work above cited.

Column 4 contains references to other authorities, and gives either the name of the first discoverer of the nebula, or a reference to the particular list or catalogue of nebulæ which has been taken as the authority for the place set down. The principal of these are—1st. The list of “new nebulæ” (Verzeichniss neuer Nebelflecke), in pp. 73 to 76 of the work of M. AUWERS already cited. These are referred to in the following form:—Auw. N. 1, Auw. N. 2, &c. 2ndly. Under the form D’Arr. 1, 2, &c., are given a series of objects contained in a MS. list of 125 nebulæ, kindly communicated to me by their discoverer, M. D’ARREST, Director of the Royal Observatory of Copenhagen, and reduced by him to the epoch (1860.0) of this Catalogue, with their precessions for 1880. 3rdly. A great number of nebulæ cited under the form “R. novæ,” whose places have been approxi-



mately obtained from the diagrams accompanied by micrometrical measures of position and distance, or from more loose and general indications contained in Lord Rosse's paper in the Philosophical Transactions for 1861, the comparisons being in all cases made with those nebulae in my Catalogue of 1833 whose numbers stand annexed to them in column 2, with an italic letter appended, thus:—

322, *a* R. nova; 319, *a* R. 3 novæ.

In cases of which latter kind it is intended to express merely that nebulae to the number indicated, not otherwise identifiable, will be found on due search in the immediate neighbourhood of the place approximately set down. Lastly. The names of Professor G. P. BOND, Mr. S. COOLIDGE, and Mr. J. T. SAFFORD in this column of the supplementary list of nebulae refer to the places of nebulae and clusters in a list of objects of that description discovered at the Observatory of Harvard College, obligingly communicated to me by Professor BOND, Director of that establishment, too late for their introduction into the body of the Catalogue.

Besides these references, in which the places set down have been adopted from the catalogues above mentioned, column 4 also contains synonyms or identifications of objects observed by myself with those contained in MESSIER's lists communicated to the French Academy, or to the *Connaissance des Temps* for 1783 and 1784. These are cited by the number they bear in MESSIER's own list, thus, M. 1, M. 2, &c. They have, with very few exceptions, been observed and described by myself or my Father, and their places here set down are given as results from our observations. In the few excepted cases they are taken from M. AUWERS's catalogue already spoken of. The nebulae also whose identity has been (sometimes satisfactorily, but for the most part very doubtfully) made out with objects in Mr. DUNLOP's Catalogue of Southern Nebulae, are indicated by the letter Δ , thus, Δ . 169, &c. In a few cases, chiefly those of nebulous stars, planetary nebulae, or very star-like objects, which have been set down as stars in catalogues of authority; these are also referred to by name and number in column 4.

Many of Mr. DUNLOP's nebulae are contained in LACAILLE's catalogue, as also some of MESSIER's, but of that catalogue two objects only, not so identifiable, viz. Nos. 38 and 40 of M. AUWERS's catalogue of LACAILLE's nebulae, have been considered as definitely enough described (*nébuleuses sans étoiles*) by that astronomer to be included in the present Catalogue.

Column 5 contains the Right Ascension in time for 1860.0 of each object in the Catalogue. When this is given to decimals of seconds, it is to be understood as having been brought up from the mean of the observations given in my former Catalogues, or from the mean of those (where not observed by myself) in Miss C. HERSCHEL's Zone Catalogue above mentioned*. When the R.A. is given only to the nearest minute or degree, it will of course be understood that the place is too loosely determined to render

* In some cases a careful subsequent revision of the catalogued observations *seriatim* has necessitated altering these R.A.'s by a few decimals of a second (seldom more) *after the process of reduction to 1860*. In all such cases the alteration has been applied as a *correction* to Mr. KERSCHNER's figures, so as not to disturb the amounts of precession allowed—a procedure perfectly legitimate and productive of no error. The same remark applies to col. 8.

further precision of statement other than illusory. This is the case with the greater part of those set down as “R. novæ.”

Column 6 contains the precession, in seconds and decimals, in R.A. for 1880.0.

Column 7 contains the number of observations in R.A. which have been actually used in concluding the R.A. for 1830, from which that for 1860 has been computed. In all cases (unless where the contrary is especially indicated in a note, or otherwise as by the letters B.A.C. or A.S.C., Au., &c. inserted in place of a number in the column itself—which indicate that the R.A. is that of a star in one of those catalogues, or rests upon that other authority), the observations used for all objects included in my former catalogues are brought up from the data there registered, to the exclusion of all others; and in such cases (the vast majority) no parenthesis or other distinctive mark is applied. When, however, no satisfactory R.A. is there recorded, or when the R.A. is there expressly stated to have been set down from the “working list,” the R.A. adopted is that brought up from the Zone Catalogue of C.H., and in such cases the number of observations used is enclosed in parentheses (). Dots attached (:) indicate some uncertainty in the R.A.; (::) a very considerable doubt, extending, perhaps, to a whole minute; ? and ?? express still wider limits of uncertainty. In those nebulae of my Father’s catalogues which have no number corresponding in column 2 (indicating the absence of any observations of my own), the places set down both in R.A. and Declination are those brought up from the Zone Catalogue of Miss C. H., and the numbers of observations on which they rely are set down in the appropriate column without any parenthesis or distinctive marks, the absence of any number in column 2 being a sufficient indication. In the case of M. D’ARREST’S nebulae, the numbers in column 6 enclosed thus [] indicate the number of his observations of the nebula employed by himself to give the place.

Columns 8, 9, and 10 contain, in like manner, the North Polar Distance for 1860, the precessions for 1880, and the numbers of observations used for P.D. in the case of each object; and the same remarks apply to these as to columns 5, 6, and 7.

In column 11 is given a short description of the nebula or cluster in abbreviated words, made out from an assemblage and comparison of *all* the descriptions of each object given in my Father’s *and* my own observations. As regards the former, recourse was had, not to the printed account in the Philosophical Transactions (which gives only a single description), but to a series of manuscript sheets in the nature of a REGISTER (and *as such cited* in the notes which follow this Catalogue), into which have been transcribed, *verbatim*, from the original sweeps, all the descriptive parts of each and every observation of each cluster or nebula in the order of their dates, and the data for computing their places, derived from the sweeps by applying the index and clock corrections pertaining to each. In this Register the nebulae are entered, each with its class and number, and each on a separate sheet; the whole series being arranged, however, not in the order of their classes and numbers, but in the order of the dates of their discovery, from No. 1, corresponding to October 28, 1783, to No. 2508, corresponding to September 30, 1802. Of these, the first 2500 only are included in the catalogues com-

municated to the Royal Society; the other 8 are printed in the form of an Appendix to my Cape Catalogue, in p. 128 of the "Results of Observations," &c. A similar and separate Register in sheets has been kept for my Father's observations of MESSIER's nebulae, and these have in like manner been collated with my own observations of the same objects in framing the ultimate, or, as it may be termed, the average description of each.

In making out these descriptions, it was found to a certain degree practicable, in the particulars of brightness, size, and extension, to make a kind of arithmetical approximation to a mean conclusion, by arranging the degrees of brightness, &c. in a progressive upward scale from 1 to 10, and taking a mean of these numbers in each case, as indicating the designating words to be finally adopted. Thus, taking the extreme degree of faintness when a nebula was declared to be "excessively faint," or "barely visible," or "hardly more than suspected" for 1, and "extremely" or "excessively bright" for 10, the intermediate degrees, such as *very faint*, *faint*, *considerably faint*, *pretty faint*, *pretty bright*, *considerably bright*, *Bright*, *very bright*, were denoted by the intermediate numbers 2, 3, 4, 5, 6, 7, 8, 9; and similarly for the scale of sizes, exchanging the words Small and Large for Faint and Bright. In the case of extension, the scale 1 to 10 was supposed arranged in the order, *Round*, *very little extended*, *elliptic* or *oval*, *considerably extended*, *pretty much extended*, *much extended*, *very much extended*, *extremely extended*, or *a long ray*. It is obvious that the qualifying words, such as "pretty" and "considerably," admit of a good deal of latitude of interpretation, and that, in reference to brightness or faintness, greatness or smallness, their meaning is rather relative than absolute; and especially, that as between *bright* or *faint*, and "considerably bright" or "considerably faint," for instance, there is so little real distinction of an absolute kind, that it is impossible to say which is to be accepted as indicating the superior degree. In the case of extension there is the same indistinctness as to precedence between the qualifying phrases "considerably" and "pretty much." Nicety, however, in this respect would be misplaced, when it is considered that when several descriptions of the same nebula, observed at different times, come to be compared, they can hardly ever be reconciled except by allowing to each qualification a latitude of meaning extending over several degrees of our arbitrary scale. In many instances, indeed, the discordance, or rather contradiction is so great, as to authorize a strong suspicion of variability in the object itself. In a few cases where, from the low altitude of the object in England, coupled with corresponding discordances of description, it was evident that it must have been seen to much greater advantage from the Cape station (as, for example, in that of h. 3375 = H. III. 754), additional weight has been attributed to the Cape observations.

In the descriptions, I have found it absolutely necessary to abstain from any specification of the estimated sizes of nebulae or clusters in angular measure. In comparing estimations of this kind I find the discordance so great, and (to speak only of my own practice) so little evidence of adherence to any definite standard of estimation, that nothing but confusion would have arisen from introducing such estimates. Nevertheless, as in the use of such a catalogue as the present some guide is necessary for the

observer, to advertise him of what sort of object he may expect to see, the following scale may be taken as conveying a general idea of the magnitudes intended by the conventional words used. Thus, a round nebula of 3" or 4" in diameter would be called *extremely small*;

one of 10" or 12", *very small*;

20" or 30", *small*, or *considerably small*;

50" or 60", *pretty small*, or *pretty large*;

3' or 4', *considerably large*, or *large*;

8' or 10', *very large*;

20' and upwards, *extremely large*.

In estimating clusters of stars (that is to say, of well separated and scattered stars) a wider acceptation must be understood, so that, for instance, a cluster of only 1' in extent would be considered *extremely* or *very small*; one of 15' or 20' *large*, and one of 30' or 40' *very large*. This amplification of scale, however, must not be held applicable to those resolved or resolvable clusters of a "globular" character marked in the descriptions as $\oplus$, which must be understood as belonging to "nebulae" and not to "clusters," so far as the conventional terms used in the descriptions are concerned. I should observe also, that when in making out the average appropriate phrase in size I have found any extravagant discordance between the estimate in words and that in figures, as, for instance, where a nebula has been described in words as *very large*, and the diameter then set down as 2', a compromise has usually been made, and the word modified, as, for instance, to *large* or *considerably large*.

The abbreviations employed in the column of descriptions and elsewhere, in the notes, &c., are as follows:—

ab.	about.
alm.	almost.
am.	among.
app.	appended.
att.	attached.
Auw.	Auwers.
A.S.C.	Astronomical Society's Catalogue.
b.	brighter.
bet.	between.
biN.	binuclear.
bn.	brightest towards the north side.
bs.	brightest towards the south side.
bp.	brightest towards the preceding side.
bf.	brightest towards the following side.
B.	Bright.
Br.	Brisbane (Sir T.'s) Catalogue of Stars.
Bo.	Bodo.
B.A.C.	British Association Catalogue.
c.	considerably.
co.	coarse, coarsely.

ch.	chevelure.
com.	cometic.
cont.	in contact.
C.	Compressed.
Cl.	cluster.
C.G.H.	"Results of observations, &c. at the Cape of Good Hope."
C.H.	Miss Carolina Herschel. When it occurs in column 4 it indicates that the object was discovered by her.
d.	diameter.
dist.	distance.
	distant.
dif.	diffused.
diffc.	difficult.
D.	double.
D'Arr.	D'Arrest.
Δ .	Dunlop.
def.	defined.
e.	extremely.

eo.	excessively.
er.	easily resolvable.
exc.	excentric.
E.	extended.
f.	following.
F.	faint.
g.	gradually.
gr.	group.
H.	Sir William Herschel.
h.	Sir John Herschel.
h.o.n.	list of omitted nebulae in C.G.H.
i.	irregular.
inv.	involved.
	involving.
iF.	irregular figure.
l.	little (adv.).
	long (adj.).
L.	Large.
Lac.	Lacaille.
Lal.	Lalande.
Lass.	Lassell.
m.	much.
mm.	mixed magnitudes.
mn.	milky nebulosity.
mon.	monograph.
M.	Middle, or in the middle.
M.	(in col. 3) Messier.
Mess.	Messier.
n.	north.
neb.	nebula.
np.	north preceeding.
nf.	north following.
nr.	near.
N.	nucleus, or to a nucleus.
o.	omitted.
ON.	omitted nebula.
p.	preceeding.
p.	pretty (before F, B, L, S, &c.).
pg.	pretty gradually.
pm.	pretty much.
ps.	pretty suddenly.
P.	poor.
Pi.	Piazzi.
P.T.	Philosophical Transactions.
quad.	quadrilateral.
quar.	quartile.
r.	resolvable, barely (mottled as if with stars).

rr.	partially resolved—some stars visible.
rrr.	well resolved—clearly seen to consist of stars.
R.	round.
RR.	exactly round.
R.nova.	New nebula discovered by Lord Rosse.
R. MS.	Manuscript notes furnished by His Lordship.
Ri.	Rich.
R.	The Earl of Rosse.
s.	suddenly.
s.	south.
sp.	south preceeding.
sf.	south following.
sc.	scattered.
st.	stars.
sev.	several
susp.	suspected.
sh.	shaped.
stell.	stellar.
S.	small.
sm.	smaller.
sw.	sweep.
Σ.	Struve.
tri-N.	tri-nuclear.
trap.	trapezium.
v.	very.
vv.	an intensive of v.
var.	variable.
W.H.	Sir W. Herschel.

Besides these abbreviations of words, the following arbitrary signs are used.

* a star; *10 a star of the 10th magnitude.

** a double star; *** a triple star.

! a remarkable object; !! very much so; !!! a magnificent or otherwise exceedingly interesting object.

? doubtful; ?? very doubtful, either as to accuracy of place or reality of existence, according to the column in which it occurs.

: , :: , see explanations already given.

△ a triangle. Forms a triangle with.

⊕ a globular cluster of stars.

○ a planetary nebula.

⊙ an annular nebula.

st. 9 Stars from the ninth (or other) magnitude downwards.

st. 9 13 Stars from the ninth down to the 13th magnitude.

As examples of the interpretation and expansion of these abbreviations some examples are subjoined.

Ex. 1. pB; vL; vg, vsmbMN 15"; pmE 162°·3; "pretty Bright; very Large; at first very gradually, then very suddenly much brighter in the middle to a nucleus 15" in diameter; pretty much extended—the position of the longer dimension micrometrically measured 162°·3 (*i. e.* reckoned from the north round to 162°·3 in the direction nsp)."

The angles of position in all cases are to be understood as so reckoned. When decimals of degrees are annexed (or if integer, written decimally thus 151°·0), they have been micrometrically measured. If thus, E 0° or E 45°, E 90°, they mean only in or near the meridian, or parallel or oblique to the meridian from nf to sp, &c., as the case may be. If with a $\pm$ annexed, the position is from a more or less careful estimation.

Ex. 2. R; psbM ill def O; pB*10·125°·4, 70"; "Round, pretty suddenly brighter in the middle to an ill-defined planetary disc; has a pretty Bright star of the 10th magnitude, whose position measured *from* the centre of the nebula is 125°·4, and whose distance also from the centre is 70" by estimation."

The relative situations of neighbouring stars or nebulae are *invariably* to be understood as thus reckoned, *i. e.* taking the centre of the nebula or other object described as a starting-point or origin of angle or distance. Thus S*s will mean that a small star is south of the nebula, *np nr that a star is near the nebula in a north preceding direction from it; *₄^f, 3'n, that a double star follows the centre of the nebula 4 seconds of time, and is 3' to the north of it.

Ex. 3. Cl; pRi; pmC; L; st6, 10 ... 15. "A cluster; pretty rich; pretty much compressed; Large; consisting of stars one of which is of the 6th, and the rest from the 10th to the 15th magnitudes."

Attached or vicinary stars or small nebulae are always placed at the ends of the descriptions. Thus $\oplus$ sf means that the nebula described "has a globular cluster following and to the southward of it." When, however, the description of a cluster ends abruptly thus, *_{*}, it is to be understood that "the place taken is that of a conspicuous double star."

The 12th column of the Catalogue contains the number of times that each nebula has been observed by both my Father and myself, whether its place were taken or not, comprising all the cases in which the object has been seen, and whether described or not. Since attention has been drawn to the real or supposed variability of nebulae, and since it can hardly be doubted that comets have occasionally been observed as nebulae, this enumeration is not without its importance. In this column the abbreviation "mon" occasionally occurs. In such cases the nebulae have been so often and diligently observed for the purpose of exact delineation or "monographing," that a special enumeration of the observations would be impossible or useless.

Finally, at the end of the line allotted to each nebula occur occasionally one or both of the marks * and †. The former refers to the notes appended to the Catalogue, the latter to the list of figured nebulae in which the publications wherein are contained figures of the nebulae are referred to by plate and figure—those at least which seem entitled, in the present state of astronomical instrument-making and pictorial representation, to be pointed out to the observer as conveying any idea of their appearance.

Notes on the Catalogue.

No.

- 12 h. 5. D'Arrest says, "h. II. positio certe erronea," but gives no indication of the correction required in R.A. or P.D.
- 29 h. 13; II. 241=II. 243. In P.T. the determining * is omitted, and in the statement of the places of these nebulae, as well as of II. 239, 240, 242, and III. 199, there is much confusion, for the correction of which see the list of errata subjoined. Auwers has threaded the intricacies of this maze with singular felicity, but has been misled in the case of II. 243 into assigning to it a totally erroneous place ($22^{\text{h}} 48^{\text{m}}$ R.A., $73^{\circ} 37'$ P.D. 1830), and, in consequence, has not perceived its identity with II. 241.
- 78 II. 3. Auwers makes the P.D. of this neb. (1830)= $99^{\circ} 32'$, from P.T., which places it $2^{\circ} \pm n$ of 17 Ceti. C.H. makes it $1^{\circ} 51' n$ of the same star, or for 1830, $99^{\circ} 42'$. In fact H. has two observations of it, neither of them more than eye-drafts with neighbouring stars, and the P.D. is concluded graphically by C.H. from these diagrams.
- 88 III. 876. The P.D. of Auwers ($81^{\circ} 16'$) is 1° wrong. The place given in P.T. is $1^{\circ} 43' n$ of 51 Piscium; so also in Register (H. 2296).
- 119 Auw. N. 4=D'Arr. 6. The place given is that brought up from D'Arrest's observations, the R.A. being set down only roughly in Auw.
- 132 h. 57=V. 20. Once looked for by Lord Rosse and not seen. Having been observed both by H. and h., there can be no doubt of its existence.
- 138 h. 61=h. 2345=V. 1. In h.'s sweep 733 the position reading is set down as $324^{\circ} 5$. This is in contradiction with a diagram made at the time, and is an obvious mistake for $234^{\circ} 5$, which = $180^{\circ} + 54^{\circ} 5$, agreeing well with the diagram and with 2 obs. of W.H., in both of which it is described as "*nf* to *sp*." There is also an erratum in the C.G.H. Catal., *for* $143^{\circ} 8$ *read* $144^{\circ} 5$, since $324 \cdot 5 - 180 = 144 \cdot 5$.
- 145 h. 64=II. 621=II. 703. Auwers remarks that A Ceti, the determining star of W.H., does not exist; but C.H. has perceived this, and by using another determining star (13 Ceti, sw. 756, W.H.), has fixed the place of the nebula II. 703 for 1800 at R.A. $0^{\text{h}} 37^{\text{m}} 47^{\text{s}}$, P.D. $93^{\circ} 53'$ (= $93^{\circ} 43'$, 1830), thereby identifying it with II. 621. Auwers, using a conjectural star, sets down the P.D. erroneously as $92^{\circ} 52'$ (1830).
- 165 h. 2356. This is the main body of the nubecula minor.
- 169 h. 2359. A complex object with several nuclei. There is an erratum in the R.A. set down in C.G.H. as resulting from sw. 488, *for* $46^{\text{m}} 12^{\text{s}} \cdot 1$ *read* $47^{\text{m}} 12^{\text{s}} \cdot 1$.
- 177 79, *a*, *b*. In Lord Rosse's diagram, α =h. 79, β =h. 78, γ =nova, accidentally omitted in the body of the Catalogue, but inserted as No. 5058 at the end. The whole Catalogue having been finally numbered before the omission was detected, it could not be inserted in its place. δ is a star; ϵ =h. 79, *a*.

- No.
 178 } h. 4007, 4008, 4012. In the Catalogue of C.G.H. these nebulae are placed
 179 } erroneously in the 23^h of R.A., owing to a mistake of a whole hour in
 196 } reducing.
 202 }
 203 } These constitute the group laid down by Lord Rosse as seen in and about the
 205 } places of h. 84, 85, 86, viz. his α , β , γ , γ' , δ , ϵ , ζ , θ . Of these, α is No. 202=h. 84;
 206 } β =No. 203=h. 85; γ =No. 206=h. 86; γ' =No. 205=h. 86, a ; δ =No. 209=h. 86, b ;
 207 } ϵ =No. 208=D'Arrest No. 10; ζ =No. 207=D'Arr. 9, and θ =h. 86, c . In the MS.
 208 } notes furnished me by Lord R. it is stated that α =h. 84, β =h. 85, and θ =h. 86.
 209 } The latter identification, however, is incorrect.
 210 }
 214 h. 88=I. 54. This is *not* the I. 54 of the P.T., which proved to be one of
 Messier's nebulae, but another subsequently inserted by W.H., so as not to
 break the order of the numbers, as appears from a MS. correction in P.T., and
 from Register (H. 570).
 275 } These constitute Lord Rosse's group seen in or near the place of h. 103, and
 276 } marked in his diagram as A, β , δ , ϵ , and another unlettered (which call γ).
 277 } These I identify as follows:—A=No. 276=h. 103; β =No. 277=h. 103, b ;
 280 } γ =No. 275=h. 103, a ; δ =No. 280=h. 103, c ; and ϵ =No. 290=h. 103, d .
 290 }
 297 In reference to M. Auwers's remark on the nebulae 170, 171, as also 167, 168
 311 (H. class III.), after very careful examination of all the data, I can arrive at no
 317 } other conclusion than that embodied in the present Catalogue under these Nos.:
 319 } h. 118 is certainly not III. 171, neither is h. 120. Both places and descriptions
 325 } disagree.
 313 } h. 119 was taken for III. 556, but no R.A. was obtained, that set down being the
 314 } R.A. brought up from C.H. The descriptions differ so materially, especially in
 the particular of extension, that they are most probably distinct nebulae.
 330 h. 124=VII. 48. Auwers remarks in his 'Verbesserungen zu h,' that this cluster,
 h. 124, is not nova, but VII. 48. This is correct. Re-examining sweep 216, I
 find an error of 1° committed in reducing the P.D.
 358 This is not in M. D'Arrest's final list, communicated to me in MS.; but being set
 down by M. Auwers as No. 15 in his 'Verzeichniss neuer Nebelflecke,' I felt
 bound to retain it.
 418 h. 160=h. 2442=I. 62. This nebula, though set down by W.H. as of the 1st
 class (*i. e.* as a bright nebula), could not be seen by D'Arrest with the Leipzig
 Fraunhofer of 6-feet focus and 4½ inches aperture. It is marked in this Cata-
 logue, however, by a mean of 4 observations, only as "F."
 428 55 Andromedæ. Although this star has been eight times examined by Lord Rosse
 without perceiving any nebulous atmosphere, yet as my observation is corrobora-

No.

rative of Piazzi's designation of it as "Nebulosa," it is retained for occasional future examination.

442) h. 169, II. 221. The places agree almost exactly, but the descriptions are irre-
 444) conciliable. One makes the nebula round, the other much extended. They
 are therefore almost certainly distinct nebulae, and there is therefore probably
 some error in the R.A. of II. 221. The neighbourhood is rich in nebulae (see
 the next note, however).

442) }
 444 } In Lord Rosse's diagram of the group about h. 169, assuming α to be h. 169
 445 } =No. 444, the others will be β =No. 445=169, α ; γ =No. 446=169, b ;
 446 } δ =No. 447=169, c ; and ϵ =II. 221.
 447 }

462 h. 179=50 Cassiopeiæ. Retained in the Catalogue for future occasional obser-
 vation. Nothing can be more difficult than to verify or disprove the nebulosity
 of a considerable star under ordinary atmospheric circumstances.

472 h. 184=III. 583. Though Lord Rosse on one occasion did not find this nebula,
 its existence cannot be doubted, having been found by h. nearly in the place
 assigned by C.H.

487 h. 193=I. 152. M. D'Arrest found this nebula too faint for observation with the
 Leipzig refractor, though placed by W.H. in Class I., and standing in this Cata-
 logue (from a mean of 3 observations) as a "bright" nebula.

501 h. 204=III. 604. C.H. and Auwers make the R.A. 1^m less. Both H. and h. rely
 on single observations. Sweep 188 h. examined and reduction found correct.

510 h. 206=III. 457. Not found by Lord Rosse; once looked for. See notes on
 Nos. 472 and 132.

516 h. 210=II. 246. Singularly enough, h. and H. are at issue about the two adja-
 cent stars. h. makes the stars south of the nebula; H., on the contrary, places
 the nebula south of the stars, and says expressly that both this nebula and
 III. 201, observed just previously, were similarly situated with regard to their
 attendant stars. Now in h.'s obs. of III. 201 (No. 513) the attendant star is
 stated to be sf the nebula, and in that of II. 246 the larger of the two stars is
 south and only a very few degrees preceding. I believe the error to lie on the
 side of the older observations, as I have a diagram of the small star nearer to
 II. 246, sf, which shows that I made no mistake of n and s.

536 I. 153. Auwers makes the R.A. for 1830 $1^h 28^m 45^s$, whereas C.H. makes it
 $2^h 15^m 13^s$. The cause of the discordance lies in an erratum in P.T. (see list
 of errata). In C.H.'s reductions the error is corrected, and I find the correc-
 tion verified on reference both to the Register (H. 1488) and the original sweep
 (sw. 596). The nebula *follows* (not precedes) the determining star.

549 h. 226=I. 154. Auwers makes the R.A. of this for 1830, $2^h 23^m 8^s$; C.H.

No.

$2^h 20^m 57^s.8$, by the observations in different sweeps differing only 18^s in R.A. The latter is the more correct; so that M. Auwers's remarks on this nebula are not confirmed. The cause of the disagreement lies in a misprint in P.T. (See List of Errata.)

557 } In Lord Rosse's description of this group, α =No.557=h.231; β =No.563=h.234;
 558 } γ =No.558=231, α ; δ =No.559=231, δ . The other nebula, "about $12'$ south
 559 } following," is probably No.563=h.234. No.561=h.233 seems to have escaped
 561 } notice.
 563 }

571 h. 240=II. 238=III. 198. C.H. has overlooked or omitted an obs. of W.H. of III. 198 in sw. 574, which, referred to, confirms Mr. Marth's surmise that the nebulae are identical.

573 II. 6. This was probably really a comet, as indicated by its description, having been subsequently looked for and not found.

574 h. 244=I. 102. M. D'Arrest found this nebula, when observed with the Leipzig refractor of $4\frac{1}{2}$ inches aperture, inferior to a 1st class nebula. In this Catalogue, from a mean of 5 observations, it ranks as "considerably bright."

591 h. 258=I. 1. M. D'Arrest found this nebula, when examined with the Leipzig refractor, not entitled to rank above the 2nd class. With this our present Catalogue agrees, it being set down from a mean of 8 observations as "pretty faint."

614 This nebula of Bessel was also looked for and not found by D'Arrest, who therefore supposes it to have been a comet.

636 h. 280=II. 502. II. 502 is described by H. as eS; F; stellar. Either then the identity is doubtful, or some change must be suspected. The place, however, agrees well.

639 h. 281=IV. 43. Once looked for by Lord Rosse, but not found. (See notes on 134, 472, 510.)

646 h. 284=III. 578. The same remark. Twice looked for unsuccessfully by Lord Rosse. On one occasion clouds were passing.

654 } In Lord Rosse's diagram of this pair and the neighbouring stars γ and δ , the figure
 655 } is in contradiction with the measures. The position of $\alpha\gamma$, instead of 2° , should, I presume, have been stated thus, $\gamma\alpha=178^\circ$, or, which comes to the same thing, $\alpha\gamma=-2^\circ$. This has been assumed in deducing the place of No. 655=289, α from No. 654=h. 289.

656 h. 291=III. 591. H. makes this nebula to be the nf of two, but both those of h. the sf.

674 h. 293=II. 603. H.'s description is pB; stellar; a pc* with eS, vF chevelure. The place, however, agrees well with that of h. 293.

684 III. 195. Auwers makes the R.A. (1830)= $3^h 11^m 50^s$ and C.H. $3^h 10^m 13^s$; but
 MDCCCLXIV.

No.

- a misprint in P.T. (see List of Errata) accounts for the difference of the minute at least.
- 708) III. 959; I. 60. The catalogued places contradict the described position of and
709) np; but this is owing to the error in R.A. of I. 60, which D'Arrest makes less by 40^s , which would place I. 60 at $3^h 19^m 35^s$ (1860).
- 710 Au. N. 17. The discovery of this nebula is attributed by Au. to Schönfeld in 1858, but it seems to be identical with that described by Tuttle (Astronom. Notices, xix. p. 224). Auwers's place is preferred, Tuttle's being only approximate.
- 768 Au. N. 18. The celebrated variable nebula of Tempel, discovered Oct. 19, 1859.
- 774 II. 594. Auwers considers this as identical with II. 548, with 1° mistaken in P.D.
- 778 h. 309=I. 155. Auwers makes the R.A. of I. 155 for 1830= $3^h 53^m 33^s$, destroying the identity of these two nebulae. But his place is deduced from an erroneous entry in P.T. (see List of Errata). C.H., by 2 observations in sweeps 608, 638 agreeing to 3^s in R.A. and $2'$ in P.D., gives a place which, brought up to 1830, gives R.A. $3^h 37^m 58^s$; P.D. $94^\circ 29' 7''$.
810. h. 311=IV. 69. M. D'Arrest found the nebulous atmosphere around the central star of this nebula very conspicuous with the Leipzig $4\frac{1}{2}$ -inch refractor.
- 826 h. 2618=IV. 26. D'Arrest's R.A. is preferred, that of h. 2618 being clearly shown to be erroneous.
- 836 II. 464. The P.D. is given by W.H. as the same with that of 44 Eridani. C.H., using an erroneous place of this star, makes the P.D. $5'$ too small. This is here corrected, and the result agrees with Auwers.
- 839 Auw. N. 20. This is the remarkable variable nebula discovered by Mr. Hind on Oct. 11, 1852. M. D'Arrest testifies to its complete disappearance on the 3rd and 4th of Oct. 1861, "Hujus nebulae ne umbram quidem detegere valeo."—"Caelo serenissimo regionem summâ curâ perlustravi adjuvante Dr. Schjellerup. *Nebula reverâ* deest." (In 1855 and 1856 it was found by M. D'Arrest within $2'$ of Mr. Hind's original place.) On Dec. 29, 1861, it was seen by M. Otto Struve with the great Pulkowa refractor, but so excessively faint as to be barely within the power of that instrument. On March 22, 1862, with the same telescope, it was again seen, but considerably brighter, so as to bear a faint illumination of the wires.
- 851 h. 314=III. 587. Not seen by Lord Rosse, once looked for, clouds passing. See notes on Nos. 639, 646, &c.
- 880 h. 322. The bright star preceding is ν Eridani.
- 908 h. 333=II. 547. Not seen by Lord Rosse, once looked for. See notes 132, 472, &c.
- 926 h. 335. Erroneously identified in my Catalogue of 1833 with III. 453 (No. 981). See the note on that nebula.

- No.
- 953 h. 341=D'Arrest 48. Observed by him as "nova," but since recognized as unquestionably =h. 341.
- 970 VIII. 43. Auwers makes the P.D. of this cluster for 1830 = $66^{\circ} 25'$, which is incorrect. The determining star is 109, *n*, Tauri, the cluster being $1^{\circ} 29'$ north of the star. This would give $66^{\circ} 39'$ for the P.D. for 1800, agreeing with C.H., and $66^{\circ} 36'$ for 1830.
- 975 h. 343. A very large diffused nebosity, distributed in zigzags. This has been looked for seven times by Lord Rosse and not found. Its existence is therefore very doubtful.
- 979 h. 2709. The place graphically determined by measurement of a diagram, as compared with h. 2710.
- 981 III. 453. This was erroneously identified with h. 335 in my Catalogue of 1833. By an unlucky coincidence, its place per working list, roughly brought up from C.H., agreed so well with the latter nebula as taken in sw. 322 (h.), that it was unhesitatingly assumed to be the same. It appears, however, that in C.H.'s reduction an error of 10^m in R.A. has been committed, the star of comparison being 10 Orionis, and the nebula *following* the star by $5^m 7^s$ (as ascertained by reference both to the register sheet (H.1160) and the original sweep (sw. 462, H.)). M. Auwers, misled by my erroneous identification, has assumed that the nebula must have *preceded* the star, which would (nearly) account for the difference, and in consequence, his R.A. of this nebula is 10^m too small. C.H.'s error probably arose from misapplying in like manner the sign of the Δ . R.A.
- 998 III. 268. Auwers's R.A. ($4^h 57^m 23^s$, 1830) is adopted in preference to $5^h 0^m 28^s$, that brought up from C.H. to the same epoch. In the sweep 367 (H.) three stars of comparison are given, 58 Eridani, α Leporis, and 19 Leporis. The Δ . R.A. of α and 19 comes out correct, but that of 58 from each is wrong by $3^m 5^s$, so that the star must have been mistaken. C.H. has used 58 and α , and has rightly brought out the place of the nebula by the former (the wrong star), and wrongly by the right one; and by an odd coincidence the two results agree well, though both wrong.
- 1030 h. 349=VII. 4. Described by D'Arrest as "Ein Ausserordentlich reicher Hauf," an extraordinarily rich cluster.
- 1133 h. 356. Looked for four times by Lord Rosse, in two of which the sky was fancied to have a milky appearance.
- 1138) h. 2841. Double nebula. In my Cape Catalogue, sweep 538, for "first" and
1139) "second" read "larger" and "smaller." The smaller is sp. The position 260° is right. It is very remarkable that in sweeps 508, 522, 658, and 761 the smaller of the two was not noticed. Is it variable?
- 1167 III. 747. Auwers makes the P.D. $8' 20''$ greater. It is difficult to identify the determining star used by C.H.

No.

- 1165 } h. 2866, 2867, 2868, 2869. $16^s.2$ added to all the R.A.'s of these nebulae in the
 1168 } Cape Catalogue to compensate an error detected in sw. 538. The correction is
 1171 } deduced from a comparison of the diagram fig. 20, Pl. VI. C.G.H. with the
 1174 } place of No. 1171.
- 1179 h. 360. $3^s.3$ added to h.'s P.D. to bring it to the place in B.A.C.
- 1180 V. 30. The place of V. 30 corrected by $+3^s.2$ in R.A. and $+25' 45''.4$ in P.D. to bring it to the place of ϵ 42 Orionis in the B.A.C.
- 1183 h. 361=V. 31. h.'s place corrected by $+0^s.4$ in R.A. and $-0' 27''.2$ in P.D. to bring it to that of ι 44 Orionis in B.A.C.
- 1185 III. 1. ?? There are two observations by H. of III. 1, but they differ enormously. One agrees with M. 43. The place of M. 43 is corrected to agree with its place in the Catalogue of Stars, &c. in the great nebula in Orion, C.G.H. p. 28.
- 1191 Chacornac's recently discovered nebula. Place from Moigno's "les Mondes," No. 9, p. 241.
- 1196 III. 269. Auwers gives as the R.A. of this nebula for 1830 $6^h 27^m 57^s$, which is mistaken by 1^h . The Philosophical Transactions says that it precedes 19 Leporis by $32^m 23^s$, and that this is no misprint appears from C.H.'s reductions.
- 1226 IV. 24. Annular according to Lord Rosse.
- 1287 III. 270. Auwers places this nebula in R.A. $6^h 40^m 20^s$, or an hour too late. Its place is very distinctly settled by two determining stars, α Leporis and 19 Leporis, the former of which it followed by $15^m 4^s$, and preceded the latter by $20^m 0^s$.
- 1425 h. 393=IV. 3. Lord Rosse's account of this nebula is extremely remarkable. "This h. 393," he says, "is an enormous nebulosity which I have traced f and n of it to a great distance—*some degrees*. It narrows at times to a band across the finding eyepiece about $6'$ or $8'$."
- 1440 h. 401=V. 27=VIII. 5. Retained as a cluster, though but a poor one. Nine times examined by Lord Rosse for nebulosity, but none seen.
- 1452 III. 271. Auwers places this nebula in R.A. $8^h 3^m 35^s$, P.D. $76^\circ 21'$ (1830). There has been some mistake. III. 271 is stated to follow 8 (ν 3) Canis, $8^m 0^s$, and to be $4'$ n of that star, which gives a place agreeing with C.H. and with the present Catalogue.
- 1454 h. 441=M. 41. This nebula was also observed by Flamsteed.
- 1455 } In Lord Rosse's diagram of this group, α is No. 1457=h. 410; β =No. 1455=410, α ;
 1456 } γ =No. 1456=410, b ; δ =No. 1458=h. 409; and ϵ =No. 1460=410, c . But
 1457 } some suspicion seems to have arisen that the principal nebulae observed were
 1458 } not really h. 409, 410, but h. 406, 407. In that case the identification will
 1460 } stand as follows:—
- α =No. 1448=h. 406.
 β =h. 406— $5^s.2$ in R.A., and $-1' 25''$ in P.D.
 γ =No. 1449=h. 407.

No.

 $\delta = \text{h. } 406 + 1^{\text{h}} 6$ in R.A., and $-5' 6''$ in P.D. $\epsilon = \text{h. } 406 + 14^{\text{h}} 7$ in R.A., and $-5' 2''$ in P.D.

1480 h. 423. This nebula is entered by C.H. as VIII. 1. B, with a remark "not in print."

1508 h. 439=VI. 6. The R.A. is nearly 2^{m} in excess of C.H. and of Auwers. Examined sweep (h.) 393 in which it was observed. Found all clear and correctly reduced.

1527 } Compared with Lord Rosse's two diagrams of the nebulae composing this group.
 1528 } None of them are "novæ." $\alpha = \text{h. } 449$; $\beta = \text{h. } 448$; $\gamma = \text{h. } 447$; $\delta = \beta$; $\epsilon = \gamma$;
 1530 } $\zeta = \text{h. } 446$.
 1531 }

1533 VIII. 44. Auwers's P.D. is 84° , instead of 82° , owing to an erratum in P.T. (See List of Errata.)

1578 h. 468=III. 479. No nebulosity seen by Lord Rosse in 5 observations. In H.'s single observation the nebula is "suspected," and in those of h. it is not positively ascertained. The object seems therefore to be merely a small resolved cluster of vFst.

1594. M. 47. Auwers assigns a R.A. greater by 4^{m} . The cluster has not since been observed. It is probably a very loose and poor one.1611 h. 480=VI. 37. h.'s P.D. corrected by $-10'$ as the presumed error of reading in the single observation obtained. Harding in 1827 (it appears) observed its P.D.= $100^\circ 10'$ (for 1830), and W.H.'s place for that epoch is $100^\circ 12'$, that of h. being $100^\circ 19' 4''$.

1615 } In Lord Rosse's diagram, $\alpha = \text{No. } 1617 = \text{h. } 483$; $\beta = \text{No. } 1616 = \text{D'Arr. } 51$;
 1616 } $\gamma = \text{No. } 1615 = 483$, α . D'Arrest's place for β is preferred to that which results
 1617 } from comparison with the diagram. h. 284 could not have been in the field, being almost a degree distant.

1633 h. 493=II. 719. h.'s R.A. in P.T. diminished by 1^{m} for an error of 1^{m} detected in the reduction of the observation. This brings it nearer to Auwers.1652 h. 3176. *Polarissima Australis*. This nebula is so near the south pole that its precession in R.A. varies from year to year with great rapidity, so that its R.A. cannot be computed correctly by the ordinary approximate method.

1666 } The four nebulae h. 508, 510; 510, a ; 510, b evidently include among them that
 1667 } third nebula referred to by Lord Rosse as the accompanying "nova" "forming
 1668 } a triangle with h. 507, 508—of the last degree of faintness." h. 507, however, is 30° distant in P.D., so that in the observation of Feb. 9, 1850, the P.D. of h. 507 must doubtless have been read as 36° instead of 66° , giving rise to a mistaken identity with one of the *two* really new nebulae at that time in view.

1696 III. 50. I find a memorandum to the effect that this nebula is lost, and was probably a comet; but I cannot recover my authority for the statement. It is described by H. as "of the last degree of faintness," and it is therefore no way

No.

surprising that it should not have been again perceived without some time and trouble bestowed, and in clear weather.

1707 h. 527=II. 48. M. Auwers, owing to an erratum in P.T. (see List of Errata), makes the R.A. of II. 48 two minutes too great, and is thus led to doubt its identity with h. 527. There still remains the rather considerable disagreement of 5' in P.D. D'Arrest found neither of these nebulae; but there can be no doubt of the existence of one at least, in or near the place here given. This is *not* the nebula seen by Lord Rosse "nearly in contact with h. 526." This latter (described already by h. as "bi-nuclear") was seen by R. as distinctly double.

1712 h. 531=M. 67. Discovered by Oriani.

1720 h. 535=II. 823. W.H. describes this nebula as "Round;" h. as "much extended," while Lord Rosse saw it as bi-nuclear, or a double nebula joined by faint nebulosity. Is it separating into two, like Biela's comet?

1735} h. 542 and II. 557. The descriptions are irreconcilable, and they must be two
1736} distinct nebulae. The R.A. of h. 542 was *not* observed, and its P.D. is set down as "hardly more than conjectural," having been looked for by working list as II. 557 and set down as such.

1742 h. 545=II. 834. Misprinted II. 844 by Auwers in the Catalogue, but the number is correct in his general list of the nebulae by numbers and classes.

1743 h. 546. Not seen by Lord Rosse in one observation. Examined sweep 21 (h.) and found all right.

1756 III. 291=D'Arr. 60. These are assuredly one and the same nebula. Auwers's declination of III. 291 ($+27^{\circ} 7'$) should be $+26^{\circ} 7'$.

1773 h. 565=III. 61. The P.D. according to H. is 70° .

1788 II. 708. Owing to an erratum in the determining star in Phil. Trans. (see List of Errata), Auwers has given the place of this nebula for 1830 R.A. $9^{\text{h}} 12^{\text{m}} 39^{\text{s}}$; P.D. $39^{\circ} 17'$, instead of $9^{\text{h}} 6^{\text{m}} 29^{\text{s}}$; $47^{\circ} 20'$.

1791}
1794} h. 577; h. 578. Not seen by Lord Rosse in one observation. (See next note.)

1792 D'Arrest 62. This nebula must surely be variable, as it is inconceivable else that it should not have been seen by h., when h. 578, to which it is almost close, was observed and its place taken. D'Arrest says, "Fugerat Herschelium nec non me anno 1862." Neither of the three (Nos. 1791, 1792, 1794) were seen by Lord Rosse. Sweep 59 (h.) and the reductions re-examined. Found all clearly written and all correct.

1804} h. 581, 582; 581, *a, b, c, d*, 582, *a, b, c, d, e, f, g*; D'Arr. 63. Of this very complex
to } group of 15 nebulae or "knots" (as they are called by Lord Rosse), six have
1815 } been determined from his diagram, and six more by the aid of notes subse-
1817 } quently furnished me from the records of the observatory at Birr Castle, con-
1818 } taining differences of R.A. and P.D. from one or other of the former. These
1821 }

No.

are indicated by the letters MS. attached in the column of descriptions. The others I identify as follows:—

α	(in Lord R.'s diagram)	is No. 1813=582, <i>c</i> .
β	" "	1812=582, <i>b</i> .
γ	" "	1811=h. 582.
δ	" "	1806=h. 581.
ϵ	" "	1815=582, <i>e</i> .
ζ	" "	1821=582, <i>g</i> .



One of those for which no data are given must have been D'Arr. 63, and the two remaining ones are included under the entries Nos. 1817, 1818 as 582, *f*.

1832 h. 590. Not seen by Lord Rosse; once looked for. Re-examined the sweep and reductions. Found all correct.

1868 h. 3171. In the omitted observations of nebulae in the last page of the C.G.H. observations, *for* h. 3170 *read* h. 3171; and this observation, combined with the two in the body of the work, gives the mean result for 1830 employed to deduce the place in the present Catalogue.

1911 h. 3185=III. 289. In consequence of a misprint in P.T. (see List of Errata), the P.D. of Auwers is 5' too small. Corrected by this, his place agrees well with my observation.

1953 M. 81 ?? A nebula observed by W.H. as described, but differing most materially in place from M. 81. It would certainly be very extraordinary should *three* nebulae so extremely remarkable as M. 81 and 82 and this be found to lie so near together.

1959 } h. 3198, 3202 are distinct nebulae, and were observed consecutively in one and
1962 } the same sweep—sw. 561 (h.).

1960 } h. 3199 and 3201 are also distinct nebulae, and were observed consecutively in
1961 } sweep 562 (h.).

1974 III. 293. M. Auwers makes the place of this nebula $9^h 24^m 4^s$; $66^\circ 30'$ (1830), instead of $9^h 48^m 48^s$; $60^\circ 13'$. The cause of the error is an erratum (see List) in P.T., where the determining star is set down as 23 Leonis instead of 23 Leonis Minoris, another of the instances of confusion arising from the use of this silly and barbarous nomenclature.

2014 h. 669=III. 65. Not seen by Lord Rosse in one observation. It was found by h. in its place *per* working list.

2019 h. 672. Not seen by Lord Rosse in one observation. Examined the sweep and reductions, and found all correct.

2043 h. 250. This nebula is so very close to the North Pole, that its place cannot be calculated by a precession proportional to the time in the usual approximate mode, the R.A. changing from year to year with extreme rapidity.

- No.
 2054 }
 2055 } In Lord Rosse's diagram, α =No. 2058=h. 692=II. 44; β =No. 2061=h. 693
 2057 } =II. 45; γ =No. 2055=692, b ; δ =No. 2054=D'Arr. 61; ϵ =No. 2057=692, c ,
 2058 } not lettered in the diagram.
 2061 }
- 2088 } II. 28, 29. Both D'Arrest and Secchi agree in placing this double nebula more
 2089 } to the south than W.H. by $15' \pm$, and D'Arrest supposes the P.D. to have been
 misread to that extent. As so great a proper motion is most improbable, and
 the identity is indisputable, I have adopted this supposition and made the neces-
 sary correction.
- 2094 h. 706. Not seen by Lord Rosse in 6 observations. Re-examined the record of
 the original obs. Sweep 115 (h.), No. 68, and the reductions. The entries are
 all clear and perfectly legible. Reduction in P.D. correct; reduction in R.A.
 erroneous by $-0^m 26^s.6$. This, however, could not have caused its non-observ-
 ation by R. This then was a comet, or is a lost nebula. The error of reduction
 is corrected in the present Catalogue.
- 2111 III. 316. C.H.'s reduction of this nebula being affected with a considerable error,
 Auwers's R.A. is adopted, after verification.
- 2144 h. 3276. Place approximate, by equatoreal zone review.
- 2189 h. 745=V. 52. Not seen by Lord Rosse when once looked for (see note on
 No. 132, &c.).
- 2192 h. 3294. The minute in R.A. doubtful.
- 2197 h. 3295. The great nebula about η Argus. According to a letter from Mr. Eyre
 B. Powell of Madras, a most extraordinary change has taken place in this nebula
 since my figure of it was delineated. He states that the southern end of the
 curious oval vacuity close to the great star, which was *decidedly closed* when I
 depicted it, is now *decidedly open*. Should this be established, it will be the
 most extraordinary fact that has yet appeared in the history of a nebula.
- 2201 h. 754=II. 99. M. D'Arrest found this nebula in the Leipzig refractor, bright
 enough to be ranked in the first class. And it is marked as "very bright" in
 this Catalogue by a mean of 5 observations. It must have been ill seen in the
 earlier observation when classed as II.
- 2231 } IV. 6=II. 131 and h. 777=III. 88. I adopt, on due consideration, the opinion
 2234 } of Auwers, that III. 88 and II. 131 are not the same. Their having been
 successively observed in the same sweep is decisive. Also, that IV. 6 is not
 III. 88, but in reality identical with II. 131. The descriptions are made out
 in conformity with this.
- 2233 } I. 118 and h. 779. The degree of P.D. is probably mistaken in I. 118. Marth,
 2236 } according to Auw., suggests that the determining star 46 Ursæ (which though

No.

not so called in B.A.C., is doubtless No. 3741 of that catalogue) was mistaken, and should have been called 46 Leonis minoris. Consulting the original sweep (sw. 487, H.), I find this surmise *not* corroborated; for the nebula, when reduced by the star next preceding it (37 Leonis minoris), gives the same Polar distance, and, within a few seconds, the same R.A. But there is some faint indication of the figure 6 in the reading of the Polar distance piece $56^{\circ} 55'$ having been written over a 7, which would have thrown the nebula somewhat below the southern limit of the sweep, and might have caused a suspicion of error at the time. I found no nebula in the catalogued place in my sweep No. 337 (h.), so that the probability of an erroneous degree is strengthened. At the same time, it is not impossible that this nebula may be identical with No. 2236=h. 779, the mistake in the degree lying the other way.

2238 h. 780=I. 172. h., in Ph.Tr., suggests that this nebula may have moved. There is, however, no ground for this supposition, as its place agrees quite remarkably with that brought up from C.H. But query if the double star have not moved, since one of the observations places it "in the middle," and a subsequent one makes the southern extremity of the nebula touch the large star of the double star.

2276 h. 806=II. 101. Found to rank as a first-class nebula by M. D'Arrest with the $4\frac{1}{2}$ -in. Leipzig refractor. In this Catalogue it stands described as "very Bright," by a mean of 4 observations. See remark in note 2201.

2310 h. 823=III. 111. There is a strange amount of discordance between the observed and reduced places of this nebula. Auwers makes the P.D. for 1830= $84^{\circ} 29'$. C.H. has reduced the single observation of W.H. by two stars 84, τ Leonis and 349 Bode Leonis, and her results differ by $10'$; τ , which gives the greater, being stated to be "too far distant in P.D." The several results stand thus:—

P.D. 1830, by Auwers	$84^{\circ} 29'$
„ by τ Leonis (C.H.)	$84^{\circ} 20'$
„ by h. obs.	$84^{\circ} 15'$
„ by 349 B. Leonis (C.H.)	$84^{\circ} 9\frac{1}{2}'$

My observed P.D. is nearly a mean between those of C.H.

2315 h. 828=II. 42. Not seen by Lord Rosse when once looked for (see notes on No. 132, &c.).

2319 h. 829=III. 351. The observations of this nebula, which are numerous, disagree so very remarkably in the particular of brightness, that a considerable suspicion of variability exists.

2373 h. 854=M. 65. There is a misprint, 45° for 75° np to sf, in the position of extension in my Catalogue of 1833. The diagram in the original sweep also corroborates this, as does also the figure (fig. 53) accompanying that Catalogue.

No.

- W.H. twice says mE in merid. (180°)—h.'s position 75° np to sf= 165° ; a mean of those of Winnecke and Auwers= 172° .
- 2377 h. 857, h. 875; M. 66. No doubt these are the same. fig. 54 P.T. 1833 corroborates their identity. The accompanying stars and their positions agree entirely. The R.A. of h. 875, however, requires to be corrected by -3^m , allowing the seconds and the P.D. observed in that observation their weight.
- 2382 II. 30. Auwers deduces his R.A. for 1830 ($11^h 12^m 21^s$) from the statement in P.T. "following 68, δ Leonis, $6^m 30^s$." C.H. from the same data concludes R.A. $11^h 11^m 31^s$ (also for 1830). The latter is (within 2^s) the correct result.
- 2388 h. 867=h. 861? These are very probably the same. But as, after all, the difference of the observed R.A.'s is sufficient to have allowed one to escape while observing the other, so that they *may* be different, and as moreover one is described as "Round," and the other as "extended," both are retained.
- 2405 h. 882=I. 20. This nebula would seem to have decreased in brightness. The bright * is 1341. A.S.C.
- 2411 h. 886=I. 131. Ranked by M. D'Arrest in the second class with the $4\frac{1}{2}$ -inch Leipzig telescope. In this Catalogue it stands as "pretty Bright" from a mean of three observations.
- 2417 III. 112. Auwers has reduced this nebula by the star given in P.T. ϕ , 74 Leonis. But I find a MS. note that this star was not dependable, and that Mayer's No. 510 is the proper determining star. The nebula was subsequently looked for and found, not in the place given by ϕ , but $8'$ from the P.D. concluded from Mayer 510. A mean of these two determinations is therefore used in this Catalogue.
- 2440 h. 907=III. 353. Auwers doubts the identity of these nebulae. But this is in consequence of a misprint in P.T. (see List of Errata), 53^m for 43^m . The error is found also in the Register Sheet (H. 937), but C.H. has avoided it and used 43^m in her reduction so as to give a R.A. agreeing within 35^s with that of h. 907.
- 2461 h. 918=II. 784. Lord Rosse, in his observation of this nebula, mentions "another brush-like, $20'$ np." This was no doubt II. 783=No. 2454.
- 2501 h. 945=I. 94. W.H. makes this nebula by one observation extended, n to s, by another nf to sp, while h. has two observations agreeing in making it extended in the parallel. Surely it does not rotate?
- 2540 h. 967. 1^m added to the R.A. It is evidently the first of the group of 4.
- 2577 III. 113. This nebula is reduced also in Auwers's catalogue by ϕ Leonis, the star set down in P.T. But C.H. remarks that ϕ was above the sweep, and otherwise observed under unfavourable circumstances, and Mayer's 510 zod. star. s. $0^\circ 31'$ is preferred, which gives a result differing by $+24'$ in P.D. and -48^s in R.A. The place adopted in the present Catalogue is in conformity with this remark. (See note on No. 2417.)

No.

- 2591 h. 1000=III. 616. The star 6m, 5' n only noticed by W.H. The other 7m, f in the parallel only by h. Are there really two stars? and are they both variable?
- 2597 h. 1002=I. 203. Auwers, in consequence of an erratum in P.T. (see List of Errata), makes the R.A. of this nebula 7^m too small. The error is corrected in the Register (H. 1889) and in C.H.'s reduction.
- 2604 h. 1009=I. 202. The same misprint in P.T. mentioned in the last note on h. 2597 has also vitiated M. Auwers's R.A. of this nebula. It is corrected in the Register Sheet (H. 1886) and in C.H.
- 2608 h. 1013=III. 381. I adopt Mr. Marth's identification of these nebulae. The place of III. 381 in the catalogue of C.H., from which my working lists were made out, is vitiated by some great mistake. The P.D. is supposed to be derived from 1 Comæ, the neb. being 1° 12' south of the star. This, however, would give 68° 9' 29" for 1830 instead of 65° 45' 0", that brought up from C.H.
- 2650 h. 1039. This cannot be identical with h. 1036, and its brightness precludes its being accepted as III. 354. But there is extreme uncertainty as to its P.D. The degree may even be wrong.
- 2652 h. 1041=II. 733. According to W.H. the position of extension is "near the meridian." If *meridian* be not a mistake for *parallel* it has changed. h. has a measure 62° 3, and an estimation 65° in another observation.
- 2653 h. 1042. This cannot be III. 3, as C.H. has reduced two obs. of this latter well agreeing, and giving a R.A. 2^m exceeding that of h. 1042, which also rests on 2 obs. of h.
- 2668 h. 1050=I. 253. The difference of descriptions is extraordinary, so that they seem hardly to pertain to the same object; but the places agree.
- 2683
- 2684
- 2685
- 2686 h. 1062, 3, 4, 5, 7, 8, 1070, 1, 3, 5, III. 391, 2, 3, 4, 5, 6. The places set down
- 2689 for the nebulae of this extensive group are made out by a most careful consideration of all the observations and records in the sweeping books which seem
- 2690 irreconcilable with a group of six nebulae only. The group, however, needs a
- 2693 thorough re-examination.
- 2694
- 2697
- 2699
- 2701
- 2702
- 2730 II. 14. Owing to an erratum in P.T. (see List of Errata) Auwers gives quite an erroneous place for this nebula (11^h 39^m 27^s R.A., 81° 9' P.D. 1830).
- 2747 h. 1103=III. 814. Auwers suspects some error of the press, since his P.D. for 1830 comes out 36° 58', while that of h. 1103 is 35° 56'. There is, however,

No.

no error, either of printing, registry, or reduction in any part of the older work. The determining star is rightly set down as 5 Canum, whose P.D. for 1800 (the epoch of C.H.'s catalogue) is $37^{\circ} 19' 42''$, and III. 814 is declared to be $1^{\circ} 32'$ north of it, so that $35^{\circ} 48'$, the P.D. of C.H., is correct, and reduced to 1830 ($=35^{\circ} 58'$) agrees with my place within $2'$. Neither is there any error of the press or of reduction, or any apparent mistake of a clerical nature in all the process of h. 1103, and the nebula observed is set down *in* the sweeping book (of course from the working list) *as* III. 814. I consider their identity therefore as fully established.

2771 } h. 1211=II. 372. II. says, the most northerly of the pair II. 372, III. 360 the
2773 } largest: h., "by diagram," makes the following nebula, III. 360=No. 2773,
the larger of the two.

2814 II. 109. The reductions of the sweep 187 (H.) in which this occurs are somewhat precarious, and in C.H.'s revision of the sweep the Δ . P.D. from 6 Comæ is set down at $1^{\circ} 50'$, that in the P.T. at $1^{\circ} 54'$ (these changes are never made without good reason), and this accounts for $4'$ out of the $5'$ difference between her P.D. and that of M. Auwers.

2846 III. 535. In a sweep two years subsequent to the obs. of this nebula by H. it was looked for again but not found. ? if a comet.

2849 D'Arr. 89. M. D'Arrest makes mention in a letter which he has done me the honour to address to me, of a nebula having the same R.A. as this, but a P.D. (1860)= $83^{\circ} 46' 42''$. He does not include it in his final list. It should, however, be looked for.

2852 } h. 1183, 7, 9, 1190, 4; II. 568, 9, 570, 1, 2, 3. There cannot be a doubt that
2856 } II. 568, 569, 570, 571, are in 82° P.D., and II. 572, 3, in 83° . It is equally
2857 } certain that h. 1183, 1189, 1190, 1194 are in 83° . They were observed in two
2862 } distinct sweeps (sw. 111 and 238); I observed also II. 572 in sw. 238, and III.
2865 } 573 in sw. 250. There must be a set of nebulae, at least 8 in number, hereabouts.
2869 } N.B. W.H. makes II. 568, 569, 570, 571, $34'$ n. of 11 Virginis. If n. be a
mistake for s, these agree with h. 1187, 1189, 1190, 1194.

2855 h. 1186=I. 90=II. 322. Marth's conjecture is right (see Auwers's note on I. 90) as regards II. 322, but not so his conclusion that II. 322=II. 377.

2878 h. 1202=I. 139=M. 61. Discovered by Oriani. N.B. The first discoverers of the nebulae in Messier's list, when not Messier himself, are mentioned by M. Auwers in his catalogue of those nebulae (pp. 66-71), except in the cases of Oriani's nebulae, M. 14?, 18?, 35?, 61, 67.

2884 1202, *a*. Under h. 1196 and 1202, two nebulae, unidentifiable, are described as companions, but there must be some great error in Lord Rosse's account of them, as the place of one is referred to a scarlet star " $10'$ south of a scarlet star R.A. $12^h 25'$." Now h. 1202 is in R.A. $12^h 14^m$. To afford a fair chance

No.

- of reobserving them, the companion 10' nf h. 1202 is entered here as 1202, α , and that south of the scarlet star, under No. 3060 as 1196, α .
- 2892 D'Arr. 90. "Reperta a me Mart. 4, 1862. Eandem reperit Schönfeldus, April 1, 1862. Vide Comptes Rendus, &c."
- 2951 II. 87. This *may* be h. 1240, but 7' in P.D. is a large error.
- 2961 h. 1253=M. 86. The nebula of Lord Rosse 14' sp this is no doubt II. 168.
- 2976 h. 1261=III. 492. III. 492 was looked for April 11, 1787, by W.H. in the place assigned to it, but was not seen. Auwers, however, makes it identical with h. 1261. Yet the descriptions are radically different, and after all there may be another nebula, the real III. 492, in the neighbourhood.
- 2992 } R. novæ. 1274, α ; 1275, α . Of the eleven "knots" seen by Lord Rosse in this
2995 } place these two are the only really "novæ." The other 9 were h. 1237, 1244, 1250 (1 & 2), 1253, 1259, 1274, 1275, and Auw. N. 30, numbered in this Catalogue 2931, 2949, 2955, 2956, 2961, 2965, 2974, 2991, 2994. h. 1203, numbered by Lord Rosse as one of the group, seems too far remote in R.A. to have been seen on that occasion.
- 2999 h. 1279=II. 156. H. says "F;" h. "vB." The latter preferred, since F might arise from fog or haze.
- 3003 h. 1282. II. 56 and II. 90. Both II. 56 and II. 90 were seen in one sweep, March 1, 1784, at 1^m interval of time (by the same star, 25 Comæ), II. 56 being 1' more north, and II. 90 3' more south than the star. This is a case of positive disappearance, for in sweep 334 (h.) the neighbourhood was carefully examined and only one nebula found.
- 3008 I. 23. By ϵ Virginis, sw. 174; n. $1^{\circ} 31'$; $\therefore$ P.D. (1830) $77^{\circ} 18' 29''$. By 34 Virginis in sw. 199, s. $0^{\circ} 19'$, whence P.D.= $77^{\circ} 25' 33''$, mean $77^{\circ} 22'$. Auwers makes it $77^{\circ} 16'$. This nebula is placed in the 2nd class by M. D'Arrest as seen with the Leipzig refractor. In this Catalogue it is set down from a mean of two observations, as "pretty bright."
- 3011 h. 1289=II. 212=II. 750. The two nebulae so designated were not observed by H. in one sweep, and are, no doubt, identical.
- 3013 h. 1290=II. 122=II. 174. These two nebulae of the 2nd class were also not observed by H. in the same sweep, and are presumed to be identical, as the places agree.
- 3021 h. 1294=M. 49. Discovered by Oriani in 1771.
- 3026 h. 1295=II. 117=II. 629. The same remark applies as in the notes on Nos. 3011, 3013.
- 3029 II. 116. Not seen by D'Arrest.
- 3043 h. 1307=I. 83. Not found by Lord Rosse when once looked for. There can be no doubt, however, of its existence in or near this place.
- 3060 1196, α =R. nova. See note on No. 2884.

No.

3075 h. 1329=I. 31=I. 38. H. describes I. 31 as "between two bright stars." The places differ $15'$ in P.D.; h. describes I. 38 (the place well agreeing with that of H.) in one observation as having a large star f, and in two others as having a star 9m, p; that is, in effect, as lying between two bright stars. N.B. The star used for I. 31 is $31\ d\ 1$ Virginis, and for I. 38, $32\ d\ 2$ Virginis. The declination of $31\ d\ 1$ is $30'$ wrong in A.S.C. (No. 1469). In B.A.C. it is right. The P.D.'s of the two nebulae of H. differ, as already remarked, by $15'$. The R.A.'s fail to agree. They must be identical with a mistake of $15'$ in I. 31. D'Arrest says he is sure there are not two nebulae here.

3078 III. 26. Place as per C.H., $12^h\ 25^m\ 32^s$, $68^\circ\ 32'$ for 1830; as per Auwers, $12^h\ 25^m\ 40^s$, $68^\circ\ 47'$ (see List of Errata). The correction of the place in P.T. is not, properly speaking, an erratum, but the substitution of a good observation for a bad one. In the obs. sw. 177 (H.), where 20 Comae was used as the determining star, the place is given only by description. In a sweep long subsequent (sw. 944) it was compared with 26 Comae in the regular form of observation, and this is of course to be preferred. Auwers's place is deduced from the earlier, and that of C.H. from the later observation, rejecting the other.

3079 h. 1322=8 Canum. This very remarkable object occurs among the list of those observed by Lord Rosse in his paper in P.T. 1861, but without a word of remark or description; and it does *not* occur among his list of nebulosities looked for but not perceived. Surely it might be inferred from this that the nebulosity surrounding the star *was* seen, or its absence would have been noticed, as in the instance of 55 Andromedae. Yet Mr. Lassell saw no nebulosity about 8 Canum.

3097 h. 1348=M. 89. Lord Rosse has h. 1343 and 1348, and in his account of them says, "two others, about $20'$ s. of 1348;" one of these must have been h. 1343, and the other h. 1349.

3103 h. 1353=I. 119. This nebula was barely perceptible, with straining the attention, by M. D'Arrest with the $4\frac{1}{2}$ -inch Leipzig refractor. It is described in this Catalogue as "considerably bright" by two observations.

3108 h. 1358, 1359, 1363=IV. 8, 9. The obs. of 1363 in my Catalogue of 1833, in which the R.A. is uncertain, undoubtedly refers to the same very remarkable double nebula, IV. 8, 9. D'Arrest is sure that there is no other double nebula in this neighbourhood.

3111 M. 90. The place is from two observations by W.H., as also the description.

3127 h. 1374=I. 273. The descriptions of H. differ so much that it is not impossible there may be another bright nebula near this place.

3138 h. 1379=II. 577. Two diagrams by h. in sweeps 141, 143, agreeing, represent this nebula as making a considerably acute-angled, nearly isosceles triangle with

No.

two following stars. H. says, "Between two Bright stars, making a triangle with them." No one now, looking at those diagrams, would call the situation of the nebula between the stars. A suspicion of proper motion arises in such a case.

3148 h. 1384=II. 148. In my Catalogue of 1833 this nebula is identified with II. 20, and in the Register Sheets (H. 320), under the head of II. 148, there is a memorandum, "Probably the same as II. 20 (H. 47)." But on examining all the observations of both nebulae, I arrive at the conclusion that they are different, II. 20 being nearly 2^m later in R.A.

3170 h. 1401. Query if not =II. 38, with one degree mistaken in P.D.

3174 See note on 3148, above.

3177 h. 1406, 1407=II. 794 (1 & 2), III. 778; h. 1428, 1435=II. 795, 796. Auwers
3179 } remarks, and justly, on the great apparent discordance of the observations of h.
3206 } and his places of II. 794, 5, 6, and those of W.H. The fact is that the places
3216 } of these in the P.T. all rest on comparisons with ϵ Ursæ in sweeps 921 and
3224 } 1001 (H.); and the observation of that star has been erroneous or mistaken in
sw. 921 by about 11' in P.D., as appears from an obs. of 73 Ursæ in the same
sweep. The nebulae affected by this error are those here enumerated, and it
requires very careful consideration to disentangle all the observations of each
nebula by both stars, and to decide on their identities. My final conclusions
are,—1st, that in these sweeps two distinct nebulae, II. 794, 1 and II. 794, 2,
were observed, and confounded together under one number (=H. 2079 register).
These are my h. 1406, 1407. 2ndly, that h. 1407 and III. 778, II. 795, 796
are correctly determined in sw. 1001 (H.). 3rdly, that in sw. 921 (H.) the
nebula set down as II. 794 was not the same as that called II. 794 in the
reduction of sw. 1001; *i. e.* that it was in fact h. 1406, and that in this obser-
vation there is also an error of 6' in P.D., or that, if not, there must be still
another nebula in P.D. 33° 54' (1860). Finally, that the place of III. 778
given in Phil. Tr., which is affected by the same *general* cause of error, requires
a correction of +9' in P.D.

3180 h. 1405=III. 44. This is the companion of M. 60, and is placed by M. D'Arrest
in the first class, even with the 4½-inch Leipzig refractor. Perhaps the very
superior light of M. 60 may have led both H. and h. to under-estimate that of
its, anyhow, much fainter companion.

3189 } h. 1414, 1415=I. 176, 177. These two, according to Lord Rosse, are connected
3190 } by faint nebulosity.

3206 III. 778. See note on 3174.

3214 h. 1426=II. 181. Auwers points out a discordance of 19' in P.D. between my
observation and that of II. 181. This is owing mainly, however, to a misprint
in Phil. Trans. (See List of Errata.)

No.

3216 }
 3224 } II. 795, 796. See note on 3174.

3228 I. 8=III. 6. The later of these nebulae is expressly stated in the register (H. 38) to be of the 1st class, though set down (it does not appear why) in the 3rd.

3254 h. 1452=I. 41. The case of this nebula is a very odd one. H. has two observations of it. One on April 5, 1784, where it is described as a "L; B; r neb; sbM; iR Fig; Class I." Another on March 3, 1789, calls it "pB; cL; i Fig; er. Many of the st. visible." So that it may be called a cluster. Both the places of H. and that of h. agree so well, that the object in all must have been the same. Here seems evidence of change.

3256 h. 1453=II. 73. Contradictory descriptions, and possibly two nebulae differing 1^m in R.A.

3311 h. 1480=I. 141. Query if not changed. h.'s observations are positive as to the clearness of the sky. But query as to the state of the speculum.

3319 h. 1485=II. 384. Not seen by Lord Rosse in two observations (hazy).

3337 } h. 1497=I. 68; II. 299; h. 1511=I. 69; h. 1536=II. 301; h. 1574=III. 382.

3338 } Auwers finds 5' Δ .P.D. between H. I. 68 and h. 1497. His place is from P.T.

3358 } 53 Virginis n. 1° 4', whereas C.H. in her reductions uses n. 1° 11', and my

3420 } observations of this and the other nebulae in this list justify the departure. I

3483 } subjoin her note on this nebula (in zone 103° C.H.):—

"I. 68, I. 69, III. 282 are each 7' more north than they are given in the "printed Catalogue. The disagreement is the result of the recalculation, and "is probably owing to my attempting more accuracy in valuing the 'numbers " 'to a degree,' &c. &c." (*i. e.* in the index reductions of the Polar distance readings which were parts of an arbitrary scale). And in the next zone (104° C.H.) occurs,

"II. 299 and II. 301 require the same memorandum." In point of fact, comparing my own observations with those reduced by M. Auwers, the differences, as stated by him, run thus:

I. 68	. . .	Δ .P.D. H-h=	+5'
I. 69			+7'
III. 282			+7'
II. 299			—
II. 301			+6'

so that in each case, where I have observed the object, the alteration is justified. This is only one out of the innumerable instances of painstaking and laborious scrutiny bestowed by her upon these reductions which have occurred to me in the collation of her zone catalogue with the original observations and with my own results.

3356 h. 1509=I. 143. Auwers places this nebula 1° 13' too much to the south in consequence of an erratum in P.T. (see List of Errata).

No.

- 3358 See note on 3337.
- 3363 V. 3. Auwers makes the R.A. of this neb. for 1830 $13^h 2^m 31^s$, which is 10^m too great. The P.T., which in this instance is correct, makes it follow 75 Leonis $1^h 44^m$.
- 3393 h. 1527. This is not impossibly III. 937, but as both R.A.'s and P.D.'s differ very much, they *may* be different, and are therefore separately stated.
- 3415 h. 1535. Not seen by Lord Rosse in one observation; clouds passing h. has two observations, both agreeing well.
- 3420 See note on No. 3337.
- 3421 II. 185. Auwers, misled by an error in P.T. (see List of Errata), makes the R.A. of this neb. too small by 10^m .
- 3426 Auw. N. 31. Not visible in the Königsberg Heliometer.
- 3483 See note on No. 3337.
- 3506 II. 22. P.D. extremely doubtful.
- 3512 II. 826. Place re-reduced by the star used by H. and A.S.C.
- 3527 h. 1597=II. 314. Auwers makes Δ .R.A. II.—h.= $+107^s$, and remarks that there is perhaps some error in P.T. This is the case (see List of Errata), and with the correction there indicated the agreement is satisfactory.
- 3550 D'Arr. 94. D'Arrest says "not found again, Feb. 19, 1863. Sky perfectly clear. Perhaps a comet."
- 3588 h. 1633=III. 926. H. says it is sp a considerable star. h. has "a $\ast 9^m$ with a very dilute nebulous atmosphere." Has the star or the nebula moved?
- 3650 III. 946. Auwers makes the declination $+89^\circ 17'$, a misprint for $+80^\circ 17'$.
- 3662 h. 1674=L. 255. Evidently ill seen by h. The description of H. preferred.
- 3664 } h. 1676, 1679=III. 422, 423. Auwers makes the P.D. $12'$ too great by reason
3668 } of an erratum in P.T. (see List of Errata).
- 3728 h. 1720=III. 666. Auwers finding Δ .R.A. II.—h.= $+52^s$, supposes a mistake of 1^m . Examined sweep 146 (h.), and found all clearly written and right reduced.
- 3750 } h. 1734, 1735=II. 309, 310. H. says the second is the larger, h. the smaller of
3751 } the two.
- 3760 }
- 3762 }
- 3763 } h. 1744=M. 101, and its attendants in more or less intimate nebulous connexion.
3764 } Of those in Lord Rosse's woodcut, P.T. 1861, p. 729, N, the principal nucleus,
3766 } is No. 3770=h. 1774; n_1 =No. 3774=1744, i; n_2 No. 3773=1744, h. The
3767 } others are not lettered, and are made out from the joint evidence of this dia-
3770 } gram and the measures of position and distance of the stars compared with the
3771 } copper plate, fig. 35.—1744, α is not improbably=III. 787.
3773 }
- 3774 }

No.

- 3820 h. 1763=III. 804=III. 835. The identity of these nebulae rests on a memorandum in MS. in my copy of Ph. Tr., supported by the reductions of all the obs. by C.H. in 3 sweeps, each with two determining stars. Auwers makes them differ by 14' in P.D.
- 3836 III. 551. Place concluded from h. 1772=III. 552 from H.'s description, viz. that it precedes that nebula by 3' or 4' (3' 30")=14^s of time.
- 3844 h. 1777=III. 347. Auwers makes Δ .P.D.=−59', but observes that there must be some misprint. Examining all, I find that such is the case (see List of Errata), which recognized, shows that 1° has been mistaken, and the identity is therefore proved.
- 3846 h. 1779=I. 144. Auwers makes the P.D. (1830)=86° 30', and H.—h.=1° 14'. The cause of the discordance is a misprint in P.T. (see List of Errata), in consequence of which the nebula is 1° 13' north of its printed place.
- 3858 } h. 1789, 1788, 1791=III. 416, 417. Lord Rosse says that of these three only
3859 } two were found. The obs. in sw. 28 re-examined—1789 and 1791 were both
3860 } observed. Moreover, in sw. 337, III. 417=h. 1791 and h. 1788 were both
observed, and 1791 is expressly stated to have been the sf of two seen in moonlight. Now the np of these could not be h. 1789, which is eF and not north, but south preceding, whereas h. 1788 by its place in sw. 338 is np. All three, therefore, really *existed* at the date of these observations. It was h. 1789 (eF) which escaped Lord Rosse's notice, though looked for with greater instrumental power. Perhaps it may have changed.
- 3863 III. 135. Auwers's P.D. for 1830 is 63° 0'. C.H. reduced to 1830 gives 62° 50' 20". Auwers has used (P.T.) 1° 5' n. of δ , 12 Bootis; C.H. 1° 16' n. of the same *. C.H. is to be preferred on every account to P.T. Her Δ .P.D.'s are grounded on a most complete and searching re-examination and recomputation (*according to the then existing star catalogues*) of all the data (in the earlier sweeps most obscure—*foliis sibyllinis obscuriora*) for determining the degrees and minutes of P.D. from the index numbers. In almost every case I find her corrections (or rather interpretations) to be justified; and I have no doubt that in this particular instance such will prove the case, though *here* I confess myself, after consulting the original sweep, unable to perceive the reason for the deviation.
- 3888 III. 319. Auwers, following P.T., which places the nebula 2° 26' north of β Ursæ min., makes the P.D. 1830=12° 46', and so it stands in the Register sheet (H. 864). But it should be 2° 26' south. So C.H. has used it, and so it proves to be on reference to the original sweep, sw. 391 (H.), giving for the P.D. 17° 36' 12".
- 3920 h. 1832=II. 695. Not seen by Lord Rosse in one observation. See note on No. 132.
- 3922 h. 3573= Δ . 342. In Auwers's list of Lacaille's nebulae, he sets down for the

No.

declination of this $-55^{\circ} 58'.8$. For $58'.8$ read $48'.8$, if it be the same object, but of that some doubt remains.

3967 VI. 8. Auwers, using χ Virginis, the determining star in P.T., places this cluster in R.A. $14^h 53^m 37^s$ (1830), $99^{\circ} 55'$ P.D. This, however, is declared by a subsequent MS. note to be a mistake for Mayer's 577 zod. star, whence the place in this Catalogue is accordingly derived. But this star, too, must have been mistaken, and on consulting the original sweep (sw. 209, H.) I find no star in the sweep whose identity can be satisfactorily ascertained. All that can be certainly affirmed is that, within a degree one way or the other in P.D., and from 5 to 10 minutes of time in R.A. of the place set down, there exists a fine cluster of the 6th class which should be looked for. Fortunately it is the only nebula observed in the sweep, a very short one.

3977 h. 1866=I. 184. Some suspicion of variability, inasmuch as one description calls it R, another E, and another mE, besides other indications in respect of brightness.

3998 III. 373. C.H., by three distinct observations in three different sweeps (400, 730, 917, H.) from the same determining star 11 Libræ (s. $0^{\circ} 13'$, s. $0^{\circ} 14'$, and s. $0^{\circ} 15'$), deduces a P.D., which reduced to 1830= $91^{\circ} 49' 39''$. Auwers, using the same star, s. $0^{\circ} 12'$ as *per* P.T., places it in P.D. $91^{\circ} 17'$, which, however, is probably a misprint for $91^{\circ} 47'$. Two of H.'s observations place the small star *south*, and one *north* of the nebula.

3999 h. 1881=II. 576. The binuclear character verified by R, who says that it is a close double nebula.

4016 h. 1892=III. 131. Query if not variable in brightness. H. in two observations calls it F and cB; h., in two others, vF and eF.

4025 } II. 756=h. 1898?. In the two observations by H. of II. 756 it is described as

4029 } cF; pL; iF; r;
pB; s; E;

and no mention is made of a double star near it, so that though the places agree within the *possible* limits of discordance, they are most probably two distinct nebulae.

4043 } 1901, *a*. Two of six seen by Lord Rosse. The others must have been h. 1901,
4044 } h. 1902, II. 541 and III. 511.

4048 } III. 886, 887. Auwers has made an error of $-12'$ in the declination, or $+12'$ in
4049 } the P.D. of this double nebula as determined from P.T. ($20'$ n. of 7 Serpentis).
The P.D. here set down is that correctly reduced, C.H. having on her part committed an error of $+2'$ in P.D.

4051 h. 1905=II. 751. In Auwers's declination, *for* $+20^{\circ} 44'$ read $+20^{\circ} 14'$, an evident misprint.

4065 II. 818. Owing to an erroneous designation of the determining star in P.T. (see

No.

List of Errata), Auwers has given the place of this nebula (1830) as R.A. $14^h 41^m 3^s$; Decl. $+60^\circ 5'$.

- 4124 } h. 1934, &c. In Lord Rosse's diagram of the group h. 1934, A, the most con-
 4127 } spicuous, would naturally be selected as identical with that nebula, but in that
 4128 } case II. 766 would not be included in the group. On the other hand, if B be
 4131 } taken for h. 1934, the identifications will stand as follows:—A=No. 4131
 4133 } =II. 766; B=No. 4128=h. 1934; C=No. 4127=1934, *b*; D=No. 4124
 =1934, *a*. This, however, supposes an error of 45^s of R.A. in H.'s place of
 II. 766, which is not probable, while on the other hand it is difficult to account
 otherwise for its not having been noticed at all. All things considered, I have
 thought it best to enter A as a new nebula, No. 4133=1934, *c*, leaving 766
 untouched.
- 4167 h. 1948=III. 74. Not seen by Lord Rosse, once looked for (see note on No. 132).
- 4173 h. 3624=M. 80. This is Pogson's globular cluster, with a variable star in the
 centre, for whose most singular history see the Monthly Notices of the R. Ast.
 Soc. xxi. pp. 32, 33, by Mr. Pogson. Mr. P. in that statement says that Sir J.
 Herschel (among others mentioned) had described it as either "cometary" or
 "nebulous." This is incorrect. In both my observations of this object it
 stands described as a globular cluster, *all completely resolved into stars*. (See
 C.G.H. h. 3624.)
- 4234 h. 1970= Σ . 5. D'Arrest calls this planetary nebula *blue*. The place used is a
 mean of his observations, that of h. (Catal. of 1833) being only Struve's roughly
 brought up. M. D'Arrest makes the diameter = $14''.6$.
- 4247 III. 727. The comparison of the place here set down with that of Auwers is
 curious for the great number of perfectly accidental errors which have heaped
 themselves together. The place (C.H.) is rightly reduced by her from σ Her-
 culis, $f 16^m 11^s$; $n 0' 14''$, which is that given in P.T., and which, reduced to 1830,
 gives for the R.A. $16^h 44^m 46^s.8$ and for the P.D. $47^\circ 58' 16''$, differing $+8^s.8$
 and $+11''$ from the exact result. In M. Auwers's catalogue it is entered thus:
 III. 127; R.A. $16^h 14^m 47^s$; Decl. $+43^\circ 1'$ (corresponding to P.D. $46^\circ 59'$).
 That is to say, there is a misprint *in each of the three particulars*. This is not
 to be taken as a specimen of M. Auwers's work, which is an admirable example
 of painstaking devotion, and far beyond any eulogy in my power to offer. But
 it is a striking instance of the way in which, in the great run of chances,
 unlucky coincidences will happen.
- 4259 h. 1974. Doubtful whether a nebula or a very faint double or triple star.
- 4294 M. 92 (= also Lalande No. 31544). Not observed by h., but 8 times by H.
 Place from Wollaston's catalogue, which is almost identical with Auwers
 (Δ .R.A.= $0^s.1$, Δ .P.D.= $0' 3''$).
- 4302 h. 1981=h. 3686=IV. 11. The annular form only perceived in the southern

No.

observations. Both H. and h., in their northern observations, describe it as of equable light throughout. It appears from Lord Rosse's observations that the annular form is much more common among these "planetary" nebulae than H. or h. had any idea of.

4364 h. 3723=II. 200. On a ground astonishingly rich.

4368 V. 13. P.D. by Auwers = $113^{\circ} 36'$ (1830), owing to an error in P.T. (see List of Errata).

4372 h. 3726= Δ . 473. There is a singular statement respecting this cluster by Cacciatores in No. 113 of the *Astronomische Nachrichten*. He observed it as a nebula, he says, on the 19th of March, 1826 (of course, therefore, Dunlop has the priority in point of date). But *where* he saw it Lacaille, he says, noted his star 1483 (*Cælum Australe*). Also, Piazzini in 1794 and 1801 in the same place saw only a star. Cacciatores in 1809 and 1810 observed the same star, but saw no nebula, only a star 9m following it (P. xvii. 341, 346). In looking for the comet of 1826, however, "fui colpito," he says, "da questa bella nebulosa." Unfortunately for this curious history, the place of Piazzini's star referred to (and which he identifies with 1483 C.A.) differs by no less than $18'$ in P.D. from that of the nebula in question, which was therefore out of the field of view, both of his own and of Piazzini's telescope, when observing the star.

4390 h. 2000. Σ . 6. Omitted by Auwers from his catalogue of new nebulae, which contains many far less remarkable. Diameter, according to D'Arrest, = $7''\cdot05$. Bessel's place = h. + $0^{\circ}\cdot8$, - $0' 22''$.

4397 h. 2004=M. 24. H.'s two observations hardly consist with this description, and their deviation in R.A. of nearly 4^m from Messier's place makes it very doubtful whether he really saw this object.

4411 M. 69. Piazzini, in a note on xviii. 122 of his catalogue, says that both M. 69 and M. 70 are 1° more to the south. But he is wrong.

4415 Auwers, N. 40. This is the nebula discovered by Tuttle on Sept. 1, 1859, and it would appear to be variable, for M. D'Arrest says (in a letter of May 8, 1863), "La nébuleuse de M. Tuttle (*Astron. Nachr.* No. 1337. p. 272) était, le 24 Sept. 1862, si brillante et si remarquable dans le chercheur (*grandis et præclara, ovalis, 2' longa, 80'' lata*), que je suis persuadé qu'elle n'a pas été telle du temps de Messier et de votre père, et de vos propres observations. Voici la position que j'ai obtenue. 1861.0 R.A. $275^{\circ} 55'\cdot6$, N.P.D.= $15^{\circ} 30'\cdot1$." The place given in the present Catalogue is that of M. Auwers, and differs somewhat, though not considerably, from this determination.

4428 M. 70. See the note on No. 4411.

4462 III. 742. This agrees too well with M. D'Arrest's place of his No. 113 not to be the same. His description is F; S; R; $*10p 12'\cdot6$, s $2' 30''$.

4473 Auwers, N. 44. This is the nebula discovered by Mr. Hind on March 30, 1845.

No.

- It was observed in May 1852 as a nebula of the first class; subsequently as "pretty faint and diluted." M. Auwers found it "surprisingly faint," and of the 2nd class at the highest.
- 4487 h. 2037=III. 743. This was seen as a planetary nebula in the twilight by M. D'Arrest with the $4\frac{1}{2}$ -inch refractor, and can therefore hardly be ranked so low as Class III.
- 4536 h. 2062=III. 144. Not seen by Lord Rosse; once looked for. (See note on No. 132, &c.)
- 4570 h. 2073. Not seen by Lord Rosse; twice looked for. h. has three observations agreeing well. The object is an equivocal one.
- 4585 } h. 2081=I. 103. According to an observation of Olbers, cited by Auwers, this
4586 } is identical with No. 4585=I. 103, the place of the latter nebula, as assigned by H., being 20' wrong in P.D. This had escaped my notice until the nebulae in this Catalogue had been finally numbered and much other work accumulated on them; and it was considered better to let No. 4585 stand, though erroneous, than to hazard confusion by striking it out and altering all the subsequent numbering.
- 4618 h. 2093. In conformity with Mr. Mason's remarks on my observations of this nebula, and with his elaborate and excellent monograph of the great nebulous system of which it forms a part, I have diminished the P.D. in my Catalogue of 1833 by 1° . It is evident that the index reading must have been mistaken, 1° for 0° . Sweep 8 examined; the writing is clear and the reduction correct, but the conclusion from Mr. Mason's observations is irresistible.
- 4628 h. 2098=IV. 1. According to Lassell this is annular, an elliptic ring with a star in the centre.
- 4654 h. 2113. Not seen by Lord Rosse; twice looked for. Examined sw. 86 (h.), in which it was observed. All found apparently correct, the observation clearly written and right reduced: and it is added, "the double star" (h. 934 in my "3rd series of observations, &c. &c.," Mem. Ast. Soc. vol. iii.) "is a good guide." A diagram accompanying the observations, by indicating lines points out the relative situation of the double star and nebula.
- 4710 h. 2133. Not seen by Lord Rosse in four observations.
- 4714 h. 3897. Not found by Mr. Lassell within 30' all round the place.
- 4723 h. 2137=III. 920. Not seen by Lord Rosse in one observation.
- 4756 h. 2148. Not seen by Lord Rosse in three observations. In one a cloud passing.
- 4775 h. 2156=III. 932. H. says, "just sf a S* to which it seems almost to be attached, but is free from it." h. says, "has a * 13m at a distance from the edge = 1 diameter by diagram." Sw. 274 (h.). This sweep re-examined. The diagram makes the star north of the nebula. The description says, "Diagram certainly right."
- 4816 2172, a. In this group Lord Rosse has given only measures of relative position,

No.

and none of distance; so that it is impossible to assign specific places to the individuals of which it consists. He speaks of five *near* to h. 2172. The diagram exhibits only four. One may possibly be III. 166.

- 4848 2184, *a*. In Lord Rosse's diagram of the group to which this belongs, α is h. 2183 = No. 4845; β = D'Arr. 117 = No. 4844; γ = h. 2184 = III. 217 = No. 4846; δ = D'Arr. 118 = No. 4847. That marked as 2184, *a* is not lettered in the diagram, and is "nova."
- 4892 h. 2205 = I. 55. Placed in the second class only by M. D'Arrest with the 4½-inch Leipzig refractor. In this Catalogue it is set down as only "pretty Bright," from a mean of seven observations.
- 4894 h. 3971 = h. 3972. These are assuredly identical; but the minute of R.A. being doubtful, that of the earlier 3971 is preferred. The mean of the seconds and the Polar distances is taken, blending the two, and also the descriptions.
- 4922 h. 2223 = III. 222. Three times called by h. "pretty Bright," and three times by h. and H., eF; vF; eF. Is this a case of variability?
- 4933 h. 2228 = h. 3982 = I. 104. Placed in the second class by M. D'Arrest. With this the present Catalogue agrees; making it "pretty Faint" by a mean of three observations.
- 4941 D'Arr. Not included by M. D'Arrest in his final list; but there are four observations of it recorded in his "Resultate," all agreeing well.
- 4964 h. 2241 = IV. 18. According to Mr. Lassell this superb "planetary nebula" is *bi-annular*, consisting of a nucleus and *two oval rings*.
- 4966 h. 2242 = III. 226. Called by h. in four observations, pB; pB; pB; pB, and in two by H. eF; vF.
- 4980 h. 2250 = III. 213. Not seen by Lord Rosse in 4 observations. In my observations of sweep 103, a very short sweep, using the quadrant instead of the index arc, and with no good zero star, both R.A. and P.D. may be a good deal wrong. My place, however, agrees pretty well with that of H. (Δ .R.A. = 5°, Δ .P.D. = 4'), and the existence of a nebula as described, *hereabouts*, is certain, but it should be looked for within somewhat wider limits.
- 4998 h. 2261 = I. 110. H. has two observations in which this nebula is called cB; h. has one where it is called eF; adding "sky quite clear."
- 5003 } h. 2263 = II. 208. These can hardly be the same. The R.A.'s differ by nearly
5004 } 2^m and the P.D.'s by 6'. The descriptions also disagree. 255°, the position of the star 14m in h. 2263, is not np but sp, and the estimates of their magnitudes differ materially.
- 5015 h. 2271 = III. 854. A very problematic object, and in which there is great difficulty in making out its nature. Stars and nebula oddly mixed.
- 5020 } h. 2274 = II. 230; 2274, *a*; h. 2275 = II. 231. In Lord Rosse's diagram of this
5021 } group, α = h. 2274; β = h. 2275; γ = nova = 2274, *a*. h. sweep 91 makes II.
5022 } 230 the np of two, and II. 231 "to have II. 230, 45° sp." This is contradicted

No.

by the diagram. There is some confusion among the observations as to whether the two nebulae II. 230, 231 really lie np or sp from each other, and it might be suspected that the P.D.'s had been read crossways, the R.A.'s being rightly set down; but Lord Rosse's diagram and measures decide the point in favour of the relative situation being here correctly given.

5051 h. 2302. Not seen by Lord Rosse in two observations. Examined the original observation, all clear and apparently correct. The nebula certainly exists in or very near the place here set down.

5061 2849, *a*. A nebula mentioned by M. D'Arrest, but not included in his MS. list of well-determined nebulae. Should, however, be looked for.

References to Figures of Nebulae in various works.

In the following list of figured nebulae, the first column contains the current number of the nebula or cluster in the present Catalogue; the second the number attached to it in my Catalogues in P.T. 1833 and C.G.II.; or if not found in either of these, the class and number in my Father's Catalogues or other sufficient designation. The third contains an abbreviated reference to the publication in which the figure will be found, viz.—

P.T. 33. The volume of the Philosophical Transactions of the Royal Society for A.D. 1833.

P.T. 44. Ditto, Ditto, for 1844

P.T. 50. Ditto, Ditto, for 1850

P.T. 61. Ditto, Ditto, for 1861

} Lord Rosse's papers.

C.G.II. Results of astronomical observations at the Cape of Good Hope by J.F.W.H.

R. di. The woodcut diagrams in Lord Rosse's paper, Philosophical Transactions, 1861; such only being referred to as express some distinct peculiarity not elsewhere figured.

B.A.A. Professor Bond's Memoirs in vol. iii. N.S. of the Transactions of the American Academy of Arts and Sciences.

M.A.A. Mr. Mason's Memoirs in vol. vii. of the Transactions of the American Academy.

D'Arr. M. D'Arrest's Inaugural dissertation and description of the Copenhagen Equatoreal, 1861.

Lam. Dr. Lamont's "Oeffentliche Vorlesung über die Nebelflecken." München 1837.

Lass. Mr. Lassell's Memoirs in vol. xxiii. of the Transactions of the Royal Astronomical Society.

Column 4 contains the number of the Plate in the volume referred to where the figure will be found, and column 5 the number of the figure in that Plate.

The figures annexed to Mr. Dunlop's catalogue are not included, as for the main part they offer no resemblance to the objects figured (when identifiable), and would serve only

to mislead. The same remark applies to most of the older figures of nebulae scattered through the volumes of the *Histoire de l'Académie Française*, and other collections. Of the older figures of the nebula in Orion, however, for curiosity's sake, a list is sub-joined. The figures accompanying my Father's memoir in *Philosophical Transactions*, 1811, are also omitted. They do not profess to be resemblances, and are given rather as types of certain classes of objects into which he there considers the nebulae to be distributable. At least they are made from very rude diagrams.

References to published figures of Nebulae.

No. in Catalogue.	h. &c.	Work cited.	No. of plate.	No. of fig.		No. in Catalogue.	h. &c.	Work cited.	No. of plate.	No. of fig.
27	2315	C.G.H.	iv.	8		1157	357	P.T. 33	viii.	81
31	15	P.T. 61	xxv.	1				P.T. 44	xix.	81
52	2322	C.G.H.	iii.	1				R. di.		
67	2327	C.G.H.	vi.	19				D'Arr.	ii.	4
105	44	B.A.A.	opp. p. 86			1163	2864	Lass.	ii.	1
106	45	B.A.A.	Ditto.			1164	2865	C.G.H.	iv.	7
116	50	B.A.A.	Ditto.			1165	2866	C.G.H.	vi.	20
117	51	B.A.A.	Ditto.			1168	2867	C.G.H.	vi.	20
138	61	P.T. 33	vi.	52		1171	Δ. 136	C.G.H.	vi.	20
169	2359	C.G.H.	v.	10		1171	2868	C.G.H.	vi.	20
187	2370	C.G.H.	iv.	6		1174	2872	C.G.H.	iv.	7
298	112	P.T. 33	v.	38		1175	2869	C.G.H.	vi.	20
303	116	R. di.				1176	2875	C.G.H.	iv.	7
352	131	P.T. 50	xxxvi.	5		1177	2876	C.G.H.	iv.	7
		P.T. 61	xxvi.	10		1179	360	C.G.H.	viii.	1
372	142	R. di.						B.A.A.	opp. p. 96	
400	151	P.T. 33	vi.	58				Lass.	i.	1
412	156	P.T. 61	xxv.	2				*	see note	
527	218	P.T. 33	ii.	28		1180	V. 30	C.G.H.	ii.	3
		R. di.				1183	361	C.G.H.	ii.	3
544	223	D'Arr.	ii.	7				P.T. 50	xxxviii.	6
560	232	P.T. 61	xxv.	3				Lass.	ii.	3
567	2487	C.G.H.	vi.	14		1185	M. 43	C.G.H.	viii.	1
572	241	P.T. 61	xxv.	4				B.A.A.	opp. p. 96	
575	242	P.T. 33	vi.	56				Lass.	i.	1
		P.T. 61	xxv.	5				*	see note	
600	262	P.T. 61	xxv.	6		1225	365	D'Arr.	ii.	2
705	2534	C.G.H.	vi.	7				Lass.	ii.	2
731	2552	C.G.H.	iv.	1		1226	iv. 24	D'Arr.	ii.	10
810	311	P.T. 33	ii.	31		1233	2910	C.G.H.	iii.	5
		P.T. 61	xxv.	17		1235	2913	C.G.H.	iii.	5
822	2620	C.G.H.	v.	11		1238	2916	C.G.H.	iii.	5
823	2621	C.G.H.	v.	11		1243	2918	C.G.H.	iii.	5
826	2618	D'Arr.	ii.	9		1248	2923	C.G.H.	iv.	9
		Lass.	ii.	4		1249	2925	C.G.H.	iv.	9
853	315	P.T. 61	xxv.	8		1258	2935	C.G.H.	iv.	9
888	327	P.T. 61	xxv.	9		1259	2933	C.G.H.	iv.	9
979	2709	C.G.H.	iii.	3		1260	2936	C.G.H.	iv.	9
980	2710	C.G.H.	iii.	3		1265	2938	C.G.H.	iv.	9
981	2711	C.G.H.	iii.	3		1266	2939	C.G.H.	iv.	9
987	2716	C.G.H.	iii.	3		1267	368	P.T. 33	iv.	36
1057	2775	C.G.H.	vi.	1				R. di.		
1082	2802	C.G.H.	iii.	6		1269	2941	C.G.H.	ii.	4
1084	2803	C.G.H.	iii.	6		1276	2948	C.G.H.	iii.	4
1085	2804	C.G.H.	iii.	6		1277	2949	C.G.H.	iii.	4
1086	2805	C.G.H.	iii.	6		1278	2950	C.G.H.	iii.	4
1089	2808	C.G.H.	iii.	6		1279	2951	C.G.H.	iii.	4
1090	2810	C.G.H.	iii.	6		1281	2952	C.G.H.	iii.	4
1135	2840	C.G.H.	iii.	2		1282	2953	C.G.H.	iii.	4
1137	355	P.T. 33	v.	49		1283	2954	C.G.H.	iii.	4
1140	2842	C.G.H.	iii.	2		1419	390	R. di.		
1141	2843	C.G.H.	iii.	2		1425	393	P.T. 61	xxvii.	11
1142	2844	C.G.H.	iii.	2		1437	399	P.T. 33	vi.	64
1143	2845	C.G.H.	iii.	2				P.T. 50	xxxvii.	10
1156	2859	C.G.H.	iv.	7				Lass.	ii.	8

TABLE (continued).

No. in Catalogue.	h. &c.	Work cited.	No. of plate.	No. of fig.		No. in Catalogue.	h. &c.	Work cited.	No. of plate.	No. of fig.
1467	415	P.T. 33	viii.	91		2841	1175	P.T. 33	vi.	55
1477	421	P.T. 61	xxvii.	12		2870	1196	P.T. 61	xxvii.	21
1511	3075	C.G.H.	iv.	4		2878	1202	P.T. 33	vii.	69
1519	444	P.T. 33	vii.	72		2884	1196, <i>a</i>	P.T. 61	xxvii.	21
		P.T. 50	xxxvii.	6		2910	1225	P.T. 33	vi.	57
1520	445	Lass.	ii.	9		2950	1245	P.T. 61	xxvii.	22
1521						2958	1252	P.T. 33	vii.	68
1532	450	P.T. 50	xxxviii.	15		2962	1252	P.T. 33	vii.	68
		Lass.	ii.	6		2972	1258	R. di.		
1565	{ 464 }	P.T. 50	xxxviii.	12		3041	1306	P.T. 61	xxvii.	23
	{ 3093 }	Lass.	ii.	5		3042	1308	P.T. 61	xxvii.	23
1567	3095	Lass.	ii.	7		3085	1337	P.T. 33	iv.	37
1677	3131	C.G.H.	vi.	12				P.T. 61	xxviii.	24
1721	536	P.T. 33	vi.	61		3101	1352	P.T. 33	viii.	83
		Lam.	i.	8		3106	1357	P.T. 50	xxxvii.	9
1728	537	P.T. 33	vi.	65		3108	{ 1358 }	P.T. 33	vii.	78
1745	3145	C.G.H.	v.	12			{ 1363 }			
1801	3154	C.G.H.	v.	8		3109	{ 1359 }	P.T. 33	vii.	78
1861	604	P.T. 33	vii.	70			{ 1365 }			
1863						3113	1362	P.T. 33	vi.	66
		P.T. 50	xxxvi.	3		3132	1376	P.T. 33	vi.	50
1911	639	P.T. 61	xxvii.	13		3151	1385	P.T. 61	xxviii.	25
2003	3221	C.G.H.	v.	9		3152				
2017	3228	C.G.H.	vi.	9		3165	1397	P.T. 33	vii.	76
		Lass.	ii.	10				P.T. 50	xxxvii.	9
2058	692	P.T. 61	xxvii.	14		3180	1405	P.T. 33	vii.	74
2063	3241	C.G.H.	vi.	2		3182	1408	P.T. 33	vii.	74
2067	3239	C.G.H.	iv.	3		3189	1414	P.T. 33	vii.	75
2102	3248	C.G.H.	vi.	5		3190	1415	P.T. 61	xxviii.	26
		Lass.	ii.	11		3240	1441	P.T. 61	xxviii.	27
2158	731	P.T. 33	v.	40		3249	1451	R. di.		
2197	3295	C.G.H.	ix.	1	Argds.	3258	1456	P.T. 33	v.	41
2216	765	P.T. 61	xxvii.	15		3275	3435	C.G.H.	i.	2
2217	766	P.T. 61	xxvii.	15		3278	1466	P.T. 33	viii.	84
2333	3324	C.G.H.	iv.	10		3321	1486	P.T. 33	ii.	27
2336	3325	C.G.H.	iv.	10		3340	1499	P.T. 33	vi.	62
2337	3326	C.G.H.	iv.	10		3356	1509	P.T. 33	vi.	67
2338	3327	C.G.H.	iv.	10		3511	1589	P.T. 61	xxviii.	28
2340	3329	C.G.H.	iv.	10		3525	3501	C.G.H.	iv.	2
2342	3330	C.G.H.	iv.	10		3531	3504	C.G.H.	v.	7
2343	888	P.T. 33	ii.	32		3570	3514	C.G.H.	vi.	1
		P.T. 50	xxxvii.	11		3572	1622	P.T. 33	ii.	25
2373	854	P.T. 33	vi.	53				P.T. 50	xxxv.	1
		P.T. 50	xxxvii.	7		3606	3523	R. di.		
		Lam.	i.	6		3614	1649	C.G.H.	iv.	5
2377	{ 857 }	P.T. 33	vi.	54		3615	1650	P.T. 33	v.	39
	{ 875 }	P.T. 61	xxvi.	16		3651	3541	P.T. 61	xxviii.	29
2378	859	P.T. 33	vi.	51		3706	3548	C.G.H.	vi.	15
2379	858	R. di.				3717	1713	C.G.H.	vi.	10
2445	910	R. di.				3750	1734	P.T. 61	xxviii.	30
2486	{ 934 }	P.T. 33	vii.	79		3766	III. 787	R. di.		
	{ 3355 }					3770	1744	P.T. 61	xxix.	35
2488	{ 936 }	P.T. 33	vii.	79		3778	III. 788	P.T. 61	xxix.	35
	{ 3356 }					3779	III. 789	P.T. 61	xxix.	35
2559	982	R. di.				4051	1905	P.T. 33	vii.	77
2597	1002	R. di.				4052				
2606	1011	P.T. 61	xxvi.	17		4058	1909	P.T. 61	xxviii.	31
2652	1041	P.T. 50	xxxvii.	7		4066	3594	P.T. 50	xxxvii.	8
2670	1052	P.T. 61	xxvi.	16		4083	1916	C.G.H.	vi.	8
2671	1053	P.T. 61	xxvi.	16		4087	1917	P.T. 33	viii.	87
2680	1061	P.T. 61	xxvii.	19		4118	1929	R. di.		
2733	1092	R. di.				4125	3610	P.T. 33	viii.	89
2756	1111	P.T. 61	xxvii.	20		4160	1946	C.G.H.	vi.	7
2760	1113	P.T. 61	xxvii.	20		4224	3641	P.T. 61	xxviii.	32
2804	1146	P.T. 33	vii.	71		4229	3644	C.G.H.	v.	4
2806	1148	P.T. 33	vi.	59		4230	1968	C.G.H.	v.	6
2807	1149	P.T. 50	xxxvii.	8				P.T. 33	viii.	86
2838	1173	P.T. 50	xxxv.	2				P.T. 61	xxviii.	33
						4234	1970	Lam.	i.	1

TABLE (continued).

No. in Cata- logue.	h. &c.	Work cited.	No. of plate.	No. of fig.		No. in Cata- logue.	h. &c.	Work cited.	No. of plate.	No. of fig.
4261	3661	C.G.H.	vi.	13		4572	2075	P.T. 33	v.	47
4284	3675	C.G.H.	vi.	6				P.T. 61	xxviii.	34
4290	3680	C.G.H.	vi.	3				Lam.	i.	5
	3680, 2	C.G.H.	v.	3	Milky Way.	4594	2084	P.T. 61	xxx.	36
4302	{ 1891 }	C.G.H.	vi.	4		4600	2088	P.T. 33	iii.	33
	{ 3686 }	C.G.H.	vi.	4		4616	2092	P.T. 33	iii.	34
4305	3688	C.G.H.	vi.	18				M.A.A.	vii.	1
	3702, 2	C.G.H.	v.	1	Milky Way.	4618	2093	P.T. 33	viii.	82
4335	3707	C.G.H.	v.	5				M.A.A.	vii.	1
4342	3713, 2	C.G.H.	v.	2		4627	2099	P.T. 61	xxx.	37
4343	1989	P.T. 33	v.	42		4628	2098	P.T. 33	v.	44
4355	{ 1991 }	P.T. 33	viii.	80				P.T. 50	xxxviii.	14
	{ 3718 }	P.T. 33	viii.	80				D'Arr.	ii.	1
		C.G.H.	ii.	2				Lam.	i.	4
		M.A.A.	iv.	1		4678	2125	P.T. 33	viii.	88
4361	3722	C.G.H.	i.	1				P.T. 44	xviii.	88
4375	3727	C.G.H.	vi.	16		4687	{ 2128 }	P.T. 33	viii.	90
4395	2002	P.T. 33	ii.	30			{ 3878 }			
4403	2008	P.T. 33	iv.	35		4729	3908	C.G.H.	iv.	11
		C.G.H.	ii.	1		4730	3909	C.G.H.	iv.	11
		Lam.	i.	10		4731	3910	C.G.H.	iv.	11
		M.A.A.	vi.	1		4733	3911	C.G.H.	iv.	11
4437	2019	Lam.	i.	9		4734	2139	P.T. 61	xxx.	38
4447	2023	P.T. 33	ii.	29		4815	2172	P.T. 61	xxx.	39
		P.T. 44	xix.	29	†	4876	2197	P.T. 33	vii.	73
		D'Arr.	ii.	5		4877	2198	P.T. 33	vii.	73
4487	2037	Lam.	i.	7		4892	2205	P.T. 33	vi.	63
4510	2047	P.T. 33	v.	46				P.T. 50	xxxvi.	4
		D'Arr.	ii.	3				D'Arr.	ii.	6
		Lam.	i.	2		4950	2236	P.T. 33	vi.	60
4514	2050	P.T. 33	v.	43		4964	2241	P.T. 33	v.	45
4532	2060	P.T. 33	ii.	26				P.T. 50	xxxviii.	13
		P.T. 44	xix.	26				P.T. 61	xxx.	40
		P.T. 50	xxxviii.	17				Lam.	i.	3
		P.T. 61	xxxi.	43		4971	2245	P.T. 33	viii.	85
		D'Arr.	ii.	8				P.T. 61	xxx.	41
4565	2072	P.T. 33	v.	48		5046	2297	P.T. 61	xxx.	42

* No. 1179=h. 360. Other figures of the great nebula in Orion will be found in Huyghens's *Systema Saturnium*, 1659; ditto, copied by Le Gentil in *Mém. Acad. Sci. Par.* 1759, pl. 21. fig. 1; Le Gentil's own figure in *do. do.* fig. 2; by Picard, *do. do.* fig. 5; another by Le Gentil, *do. do.* fig. 6. See also:—

Mairan, "Sur la Lumière Zodiacale," copied in Lalande's 'Astronomy.' These older representations, however, are mere curiosities, and present no points of exact resemblance.

Messier, *Hist. de l'Acad. Sci. Par.* 1771, p. 435...461. Plate 8 is a careful and (for the time) elaborate figure.

J. F. W. Herschel, *Mem. Astron. Soc.* ii. 1826.

De Vico, *Memoria intorno ad alcune osservazioni fatte nel Collegio Romano nel corrente anno 1838*, nebulosa d'Oriono osservata al Telescopio di Cauchoix. 1839.

Bond. A very fine engraving—not yet published.

† No. 4447. P.T. 44. xix. fig. 29. There is an erratum in this figure. For Decl. 32° 49' n read 22° 49' n.

The following nebulae have been indicated by Lord Rosse as being either "of spiral structure (S), having in them dark spaces (D), as knotted (K), or as in the form of rays (*i. e.* much elongated forms) with splits or clefts (R).

No. in Catalogue.	h. &c.		No. in Catalogue.	h. &c.		No. in Catalogue.	h. &c.		No. in Catalogue.	h. &c.	
202	84	K	2158	731	D	2717	1085	S	3249	1451	S
372	142	S	2194	749	S	2733	1092	S	3258	1456	S
594	257	K	2248	788	D	2749	1107	D	3474	1570	S
600	262	S	2373	854	S	2807	1149	R	3572	1622	S
604	264	D	2377	857	D	2870	1196	S	3750	1734	S
888	327	S	2379	858	S	2878	1202	S	3843	1776	S
895	329	K	2413	887	D	2890	1211	S	4045	1901	K
1267	368	D	2445	910	S	2910	1225	D	4058	1909	D
1458	409	K	2499	943	S	2991	1274	K	4087	1917	R
1527	446	K	2559	982	S	3049	1312	S	4572	2075	S
1676	514	D	2597	1002	S	3050			4815	2172	S
1806	581	K	2652	1041	R	3106	1357	R	4964	2241	D
2058	692	D	2670	1052	S	3121	1368	S	4971	2245	S
2066	695	S	2680	1061	S						

List of Errata and Corrigenda in Sir William Herschel's Catalogue of 2500 Nebulae in the Philosophical Transactions.

Class.	No.	No. in Catalogue.	Error and Correction.
I.	6	3702	for f. 3 ^m 56 ^s read f. 33 ^m 56 ^s
	54	214	for f. 12 ^m 44 ^s ; s. 2° 50' read f. 18 ^m 36 ^s ; s. 1° 26'
	87	2274	for f. 9 ^m 30 ^s read f. 10 ^m 30 ^s
	137	1837	for 42 Lyncis read 41 Lyncis
	143	3356	for s. 2° 7' read s. 0° 54'
	144	3846	for n. 0° 24' read n. 2° 7'
	153	536	for p. 23 ^m 16 ^s read f. 23 ^m 16 ^s
	154	549	for f. 1 ^m 23 ^s read p. 1 ^m 23 ^s
	155	778	for f. 7 ^m 49 ^s read p. 7 ^m 49 ^s
	202	2604	for f. 0 ^m 47 ^s read f. 7 ^m 47 ^s
	203	2597	for f. 7 ^m 42 ^s read f. 14 ^m 42 ^s
II.	1	4738	for p. 15 ^m ::, s. 1° :: read p. 11 ^m 45 ^s , n. 0° 17'
	11	2824	for f. 1 ^m 24 ^s , n. 0° 24' read f. 1 ^m 13 ^s , n. 0° 30'
	14	2730	for 3 (γ) Virginis f. 2 ^m 20 ^s , n. 1° 22' read 59 e Virginis p. 69 ^m 0 ^s , n. 0° 11'
	48	1707	for f. 56 ^m 45 ^s read f. 54 ^m 45 ^s
	181	3214	for s. 0° 48' read s. 1° 15'
	185	3421	for p. 11 ^m 0 ^s read p. 1 ^m 0 ^s
	239	634	for 27 (x) Persei p. 8 ^m 20 ^s , n. 0° 2' read 30 Persei p. 14 ^m 41 ^s , n. 0° 51'
	240	5046	for read 39 Pisc. p. 2 ^m 24 ^s , n. 1° 0'
	241	29	for read 39 Pisc. p. 14 ^m 24 ^s , s. 0° 11'
	242	4973	for 48 (μ) Pegasi read 87 (u) Pegasi
	264	1335	for 47 (δ) Cancri read 25 (δ) Canis
	265	1384	for 1 χ Can. read ξ 1 Can.
	286	654	for p. read 13 (ζ) Eridani p.
	314	3528	for f. 17 ^m 57 ^s read f. 15 ^m 57 ^s
	372	2771	for p. 74 ^m 24 ^s read p. 14 ^m 24 ^s
	658	1718	for 44 Lyncis read 43 Lyncis
	708	1788	for 37 Lyncis read 36 Lyncis
	794	{ 3177 3179 }	for s. 0° 49' read s. 1° 0' (see note on this No. in Catal.)
	795	3216	for s. 1° 13' read s. 1° 24' (see note on this No. in Catal.)
	796	3224	for s. 1° 25' read s. 1° 36' (see note on this No. in Catal.)
	818	4056	for 12 Draconis read 12 Draconis Hevelii
	853	14	for p. 25 ^m 38 ^s read p. 25 ^m 48 ^s
II.	6	3228	for 59 (e) Virginis p. 28 ^m 11 ^s read d Virginis f. 2 ^m 42 ^s , n. 0° 57'. The obs. belongs to I. 8
	26	3078	for 20 Comæ f. 4 ^m 30 ^s , s. 0° 37' read 26 Comæ p. 5 ^m 5 ^s , s. 0° 32'
	112	2417	for 74 φ Leonis f. 10 ^m 6 ^s , s. 1° 52' read Mayer 510. z. p. 61 ^m 48 ^s , s. 1° 10'
	113	2577	for φ Leonis f. 34 ^m 18 ^s , s. 1° 3' read Mayer 510. z. p. 37 ^m 36 ^s , s. 0° 31'
	178	631	for 17 (γ) Persei f. 9 ^m 6 ^s read 17 (r) Persei f. 10 ^m 0 ^s
	192	419	for 72 Ceti read 62 Ceti

TABLE (continued).

Class.	No.	No. in Catalogue.	Error and Correction.
III.	195	684	for f. 42 ^m 42 ^s read f. 41 ^m 6 ^s
	199	628	for 27 (κ) Persei p. 8 ^m 27 ^s , n. 0° 2' read 30 Persei p. 14 ^m 44 ^s , n. 0° 55'
	256	1641	for s. 0° 48' read s. 0° 58'
	289	1911	for s. 0° 25' read s. 0° 31'
	293	1974	for 23 Leonis read 23 Leonis minoris
	319	3888	for n. 2° 26' read s. 2° 26'
	347	3844	for s. 1° 17' read s. 0° 17'
	353	2440	for f. 53 ^m 4 ^s read f. 43 ^m 4 ^s
	369	3618	for — 25 ^m 41 ^s read f. 25 ^m 41 ^s
	422	3664	for n. 0° 44' read n. 0° 36'
	423	3668	
	511	4042	
	607	1645	for f. 3 ^m 5 ^s read f. 3 ^m 11 ^s
	627	1820	for p. 12 ^m 33 ^s read p. 12 ^m 23 ^s
	739	4149	for 43 Lyncis read 42 Lyncis
	751	1897	for p. 32 ^m 30 ^s read p. 32 ^m 47 ^s
	778	3206	for 39 Lyncis read 38 Lyncis
			for s. 1° 4' read s. 1° 15' (see note on this No. in Catal.)
IV.	29	2255	for f. 3 ^m 36 ^s read f. 3 ^m 46 ^s
	31	4802	for — 0° 37' read s. 0° 37'
V.	13	4368	for n. 0° 39' read s. 0° 38' ::
VI.	8	3967	for 26 χ Virginis f. 23 ^m 44 ^s , s. 0° 6' read Mayer 577. z. f. 1 ^m 48 ^s , n. 1° 26'
VII.	6	1509	for 50 Geminorum read 51 Geminorum
VIII.	11	1534	for 50 Geminorum read 51 Geminorum
	28	1229	for (1λ) Orionis read (1st χ) Orionis
	44	1533	for 5 (η) Can. min. read 4 (γ) Can. min.

The following nebulae are declared in MS. notes to be identical.

II. 6=I. 1; II. 119=II. 94; II. 148=II. 20; II. 176=II. 70; II. 243=II. 241; II. 703=II. 621;

III. 6=I. 8; III. 198=II. 238; III. 835=III. 804.

Errata and Corrigenda in M. Auwers's Catalogue.

Page.		For	Read	Page.		For	Read
19		II. 844	II. 834	37	III. 946 in Decl.	89°	80°
20	III. 291 in Decl.	27°	26°	39	III. 347 in Decl.	24°	25°
24	II. 30 in R.A.	11 ^h 12 ^m 21 ^s	11 ^h 11 ^m 33 ^s	40	II. 751 in Decl.	20° 44'	20° 14'
25	IV. 59 under Δα	—5	—31	40		I. 282	I. 182
26	III. 385 in R.A.	16 ^h	11 ^h	42	III. 127.....	III. 127	III. 727
26	III. 388 in R.A.	10 ^h	11 ^h	42	Do. in R.A.	14 ^m	44 ^m
26	II. 342 in R.A.	10 ^h	11 ^h	42	Do. in Decl.	43°	42°
28	III. 814 in Decl.	53°	54°	72	No. 27 Decl.	58° 8	48° 8
32		III. 858	III. 850	77	List of Errata, II. 341...	16 ^m & 11 ^m	16 ⁿ & 11 ^a
33	III. 778 in R.A.	13 ^h 37 ^m	12 ^h 37 ^m	77	Ditto, III. 680...	26 ^m & 16 ^m	26 ⁿ & 16 ⁿ

M. Auwers has given a list of errata and corrigenda required in my two previous Catalogues. They are very numerous, but relate almost exclusively to errors of identification with my Father's classes and numbers. They had been, with hardly an exception, detected and rectified during the process of preparing and arranging the present Catalogue, which being therefore expurgated of them, it is unnecessary to annex a list of them here.

One very important erratum, however, must be noticed, not having been set down by M. Auwers. In p. 494, explanation of plates, Phil. Trans. 1833, figs. 13...18, for pmbM; vbM; vmbM read psbM; sbM; vsbM.

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1	h.	II.	D'Arrest, 1	h m s	s	[4]	63 4 0	-20.05	[4]	F; S; R; bet*11 and *14 ...	0
2	4014			0 0 5	+3.07	3	120 41 11.5	20.05	3	eF; cL; mE; vglbM	3
3	4015			0 1 13.8	3.065	2	124 38 54.5	20.05	2	F; cL; vLE; glbM	2
4	1	III. 868		0 1 28.3	3.062	2	86 7 38.5	20.05	2	eF; pL; vglbM	3
5	2	III. 866		0 1 34.2	3.073	2	57 20 43.5	20.05	2	eF; pL; vglbM	3
6	2, a		R. nova	0 1 35.7	3.084	2	57 20 43.5	20.05	2	vF; vS; Sst+neb	3
7	3	II. 591		0 1		...	57 20		...		0
8	4			0 1 37.0	3.076	3	74 57 52.5	20.05	3	vF; pS; R; glbM	4
9		III. 147		0 1 52.1	3.083	1	63 3 27.5	20.05	1	pB; S; R; bM.....	1
10	2308	III. 461		0 2 33.4	3.081	1	64 49 57.5	20.05	1	3Sst+neb	1
11	2309			0 2 47.2	3.061	1	115 44 58.5	20.05	1	vF; cL; mE; gbM	2
12	5	IV. 15		0 2 57.1	3.053	2	147 48 14.5	20.05	2	vF; S; R	2
13	2310			0 3 14.6	3.085	(1)	63 4 58.5	20.05	1	vF; vS; stell	2*
14	6	II. 853		0 3 25.2	3.033	1	147 46 24.5	20.05	1	eF; p of 2.....	1
15	2311			0 3 31.8	3.089	2	57 25 28.5	20.05	2	pB; pL; E 0°+	3
16			Auw. N. 1	0 3 39.0	3.020	2	147 46 21.5	20.05	2	eeF; S; R; f of 2.....	2
17	2312			0 3 41.7	3.078	...	71 59 0.7	20.05	...	F (Schmidt 1861, Oct. 10) ...	0
18	7	III. 861		0 4 22.0	3.023	1	147 43 45.5	20.05	1	eF; S; R	1
19		III. 456		0 5 7.5	3.093	2	59 44 3.5	20.05	3	vF; pS; R	4
20	8	IV. 58		0 5 11.3	3.076	1	84 21 57.5	20.05	1	vF; pS; iF	1
21	9			0 5 31.4	3.188	3	18 15 26.5	20.05	3	vF; vS; R; vsmB*10; *12 241°4; 25°.	4
22	10			0 5 45.3	3.095	1	59 51 46.5	20.05	1	eF; *12, 45°, 325°	1
23	2313			0 5 59.4	3.096	1	59 29 20.5	20.05	1	eF; vS	1
24			Auw. N. 2	0 6 47.8	3.052	1	113 57 14.8	20.04	1	eF; L; vglbM; L*cont, f ...	1
25	11	III. 183		0 6 59.2	3.074	...	84 47 28.8	20.04	...	A nebula (Markree Cat. 1852, Oct. 22).	0
26	2314			0 7 47.8	3.089	1	72 14 19.8	20.04	1	vF; S; E	2
27	2315		Δ. 507	0 7 53.1	2.978	1	151 6 13.8	20.04	1	eF; S; R; bM.....	1
28	12			0 8 0.4	3.028	3	129 59 33.8	20.04	3	vB; vL; vmiE; tri-N	3†
29	13	II. 241 = II. 243		0 8 8.9	3.083	1	78 19 58.8	20.04	1::	eL; eF; diff.....	1
30	14	III. 248		0 8 18.0	3.088	1	73 26 48.8	20.04	1	F; S; R; sbM	4*
31	15	V. 16		0 9 14.6	3.065	1	97 5 46.8	20.04	1	vF; S; iR; psvlbM	3
32	15, a		R. 6 novæ	0 9 14.6	3.112	1	60 42 9.1	20.03	1	eF; L; 3 or 4st+neb	2†
33	16			0 11 6.2		...	60 42		...	Nos. 32...37 incl.....	0
34	17			0 11		...	60 42		...	F; S; R; psbM	1
35	18	II. 257		0 13 54.8	3.106	1	68 24 45.7	20.01	1	F; S; R; psbM	1
36	19			0 14 5.3	3.107	1	68 19 15.7	20.01	1	E; bi-N; 3Bst near	1
37	20			0 14±		...	68 19±		...	{ Several F, S (3 novæ at least presumed). }	0
38	2316			0 14 18.5	2.968	2	139 24 41.7	20.01	2	eF; S; R; gbM; 1st of 4	2
39	2317			0 14 28.2	2.967	2	139 25 13.7	20.01	2	eF; vS; R; 2nd of 4	2
40	2318			0 14 30.0	2.967	2	139 26 36.7	20.01	2	vF; S; R; gbM; 3rd of 4	2
41	2319			0 14 36.9	2.966	2	139 24 21.7	20.01	2	F; S; R; gbM; 4th of 4	2
42	19			0 15 9.8	3.088	1	80 17 48.7	20.01	1	F; pL; R; gbM	3
43	18			0 15 10.5	3.124	1	61 1 29.7	20.01	1	F; vS; R; gbM	1
44	2320			0 15 55.7	2.970	1	136 3 13.0	20.00	1	vF; pS; R; bM; r	1
45	2321			0 16 56.0	3.004	2	123 19 22.3	19.99	2	pB; pL; iE; *14, f.....	2
46	20			0 17 38.2	3.266	1	29 26 39.3	19.99	1	Cl; pS; pC; st11...18.....	1
47	2322		Δ. 18= 47 Toucani	0 17 47.4	2.721	4	162 51 33.3	19.99	4	⊕; !; vB; vL; vmCM	4†
48	21	III. 148		0 18 39.6	3.134	2	61 33 49.6	19.98	2	pF; pL; R; pslbM	3
49			D'Arrest, 2	0 18 49	3.117	[1]	68 58 20	19.98	[1]	vF; S; 3 st near, making quadr.	0
50	22			0 19 18.9	3.411	1	19 23 2.6	19.98	1	Cl; pR; iC; st9...12	1
51	2323			0 19 51.3	2.989	2	124 27 21.9	19.97	2	vF; pL; iE; D*2', np.....	2
52	2324			0 20 17.3	2.877	2	147 45 22.9	19.97	2	pB; S; R; mbM	2
53	2325			0 20 25.1	2.685	1	162 18 23.9	19.97	1	pB; pS; iE; vglbM	1
54	23	III. 869		0 21 41.4	3.123	2	87 56 3.2	19.96	2	vF; S; bM; D*vnr; p of 2 ...	3
55	23, a		R. nova	0 22 0.7	+3.123	...	87 57 38.5	-19.96	...	No descr (MS)	0

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
61	h. 23, b	II.	R. nova	h m s	s	...	87° 54' 8.5	-19.95	...	No descr (MS)	0
62	25	II. 854		0 22 4.7	3.077	2	87 54 38.5	19.95	2	pB; vS; IE, 0°±; bM; f of 2	3
63	24	VIII. 79		0 22 6.2	3.302	1	30 33 4.5	19.95	1	Cl; vL; pR; IC; st 9...13	2
64	23, c		R. nova	0 22 8.7	3.123	...	87 54 8.5	19.95	...	No descr (MS)	0
65	2326			0 22 42.6	2.979	2	124 2 2.5	19.95	2	F; pL; pmE; vgbm; p of 2...	2
66	26	II. 855		0 23 0.2	3.076	3	88 41 12.5	19.95	3	pF; cL; R; vglbm; r	6
67	2327			0 23 27.6	2.976	2	124 1 45.8	19.94	2	vB; L; vmE, 47°±5; psbM; f of 2; *10 327°9 45"	2†
68		VI. 35		0 23 48.3	3.232	1	29 16 1.8	19.94	1	⊕; vF; S; eC	1
69		II. 471		0 24 3.5	3.096	1	80 34 1.8	19.94	1	F; iF; lbM	1
70	{27=}			0 24 38.0	3.057	2	95 55 33.1	19.93	2	{ F; pL; vLE; vgbM; } *8.9, 75°±; 5'.	2
71	28			0 25 12.7	3.367	1	27 29 19.4	19.92	1	Cl; pL; IC; st 11...12; D*...	1
72	29			0 25 30.6	3.243	1	42 16 32.4	19.92	1	vF; vL; iR; g; smbM *11	1
73	2329			0 26 22.3	2.971	1	122 33 57.7	19.91	1	vB; S; IE 90°; smbM *11	2
74	{30=}	II. 478		0 26 57.8	3.042	2	100 28 30.7	19.91	2	pF; pL; IE 90°; vglbm	4
75	2331			0 27 3.7	2.513	1	163 53 11.0	19.90	1	vF; L; R; vglbm	1
76	31	III. 467		0 27 14.0	3.033	1	103 25 59.0	19.90	1	eF; vS; R	2
77	2332			0 28 2.9	2.818	2	146 33 23.3	19.89	2	vF; pS; R; glbm; 3stf	2
78		II. 3		0 28 32.5	3.044	2	99 32 4.3	19.89	2	F; L; mE; bet 2cBst	2*
79	32	III. 476		0 28 43.0	3.147	1	66 48 49.3	19.89	1	vF; vS; stellar; *7, 15°, 5' ...	2
80	32, a		R. nova	0 28		...	66 48		...	Makes Dneb with h. 32	0
81		III. 954		0 29 20.1	3.039	1	100 51 4.6	19.88	1	eeF; S	1
82		D'Arrest, 3		0 29 33	3.15	[1]	66 46 18	19.87	[1]	F; pL; R; *6, 3½ dist	0
83		III. 223??		0 29 41.2	3.008	1	109 44 5.6	19.88	1	vF; pL; iE; 2pBst sf	1
84	33	III. 871		0 30 1.5	3.076	2	88 48 44.9	19.87	2	vF; S; R; vgbM; *11, 225°±; 80'.	3
85	2333			0 30 2.4	2.968	3	120 14 30.9	19.87	3	eF; S; vLE; amBst	3
86	2334	III. 223		0 30 20.6	3.004	1	110 42 9.9	19.87	1	pB; pL; E; gbm; r	1
87	2335			0 30 21.5	2.446	2	163 56 28.9	19.87	2	eF; S; vLE; r; *8 near	2
88		III. 876		0 30 46.6	3.098	1	82 6 6.2	19.86	1	vF; pL; iR; *np inv	1*
89		III. 870		0 31 0.5	3.079	1	88 1 7.2	19.86	1	vF; S; iR; vgbM	1
90	35	II. 707		0 31 11.4	3.278	1	42 26 0.2	19.86	1	pB; vL; iR; vgmBm; r	2
91		D'Arrest, 4		0 31 15	3.08	[2]	87 36 6	19.85	[2]	F; S; R; lbM	0
92	34			0 31 40.6	5.151	1	5 26 27.6	19.84	1	Cl; vL; R; 150...200st 10... 18.	1
93	36			0 31 40.7	3.407	1	29 42 31.5	19.85	1	Cl; pL; R; st 11...15	1
94	37			0 31 49.4	3.081	1::	87 25 4.5	19.85	1	vF; L; p of 2; st 15 close	1
95	38	II. 479		0 31 53.4	3.039	1	99 46 25.5	19.85	1	pB; pL; iE 0°±	3
96	39	III. 872		0 32 2.3	3.072	5	89 54 28.5	19.85	5	F; pS; pmE; bM; 1st of 3...	6
97		II. 857		0 32 4.5	3.079	1	87 57 7.5	19.85	1	F; S; vgbM	1
98	40	II. 856		0 32 5.6	3.080	1	87 42 54.8	19.84	1	pB; S; R; vgbM	2
99	40, a		R. nova	0 32		...	87 42		...	(?=h. 37 or II. 857)	0
100	41	III. 595		0 32 6.8	3.072	1	89 51 23.8	19.84	1	F; pS; R; psmbM; f of 2	2
101	42	II. 860		0 32 14.4	3.081	1::	87 27 4.8	19.84	1::	pB; pS; R; vgbM; 2nd of 3	5
102	43	III. 873		0 32 21.4	3.072	2	89 55 7.8	19.84	2	vF; cL; E; vglbm; f of 3	3
103		D'Arrest, 5		0 32 22	3.08	[1]	87 37 24	19.84	[1]	F; vS; *8, p 27°7, ls	0
104		II. 858		0 32 22.5	3.079	1	87 52 7.8	19.84	1	pB; S; vgbM	1
105	44	V. 18	C.H.	0 32 45.4	3.243	1	49 4 49.8	19.84	1	vB; vL; mE 165°; vgvmbM	6†
106	45	V. 36		0 32 47.8	3.237	1	50 1 34.8	19.84	1	vF; vL; mE 0°	4†
107	46	II. 452		0 33 32.4	3.020	3	104 38 20.1	19.83	3	B; pS; R; psbM; r; *90°±	4
108	46, a		R. nova	0 33 32		...	104 38		...	E 0°±	0
109		III. 200		0 33 48.8	3.129	2	74 17 8.4	19.82	2	F; S; bet 2 Sst	2
110	2336			0 33 49.9	2.763	2	146 56 14.4	19.82	2	vF; S; R; p of 2	2
111	47	II. 209		0 34 5.4	3.166	1	65 16 6.4	19.82	2	pF; pS; gvlbm; r	4
112	2337			0 34 25.7	2.757	2	146 58 27.7	19.81	2	F; S; R; amst; f of 2	2
113	49	III. 244		0 34 28.9	2.990	1	111 48 59.7	19.81	1	eF; vS; IE 0°...90°	2
114	48	II. 480		0 34 30.2	3.033	1	100 47 1.7	19.81	1	F; S; IE 90°±; glbm	3
5058				0 35 1.0		...	89 50 6.6		...	See No. 5058.	
115	2338			0 35 2.1	+ 2.344	3	164 9 52.7	- 19.81	3	F; iR; vgbM; 1st of several	3

No. of Cata- logue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	"			
116	50		M. 31	0 35 3.9	+3.252	1	49 29 45.7	-19.81	1	{!!!eeB; eL; vmE; (An- drom. Gt. Neb.) Bifid (Bond)	13+
117	51		M. 32	0 35 5.3	3.250	1	49 54 12.7	19.81	1	lvvB; L; R; psmbMN	8+
118	2339			0 35 5.4	2.338	1	164 14 1.7	19.81	1	vF; R; 2nd of several.....	1
119			{D'Arrest, 6= Auw.N.4}	0 35 6	3.07	[3]	89 55 0	19.81	[3]	vF; pL; R (Bond, Jan. 1853)	0*
120	52	VIII. 78	C.H.	0 35 19.2	3.457	1	28 58 43.0	19.80	1	Cl; L; lC; st 9...10	3
121	53			0 35 24.0	3.204	1	58 10 55.0	19.80	1	eF; S; R; *13, 20" 180° ...	1
122		II. 444		0 35 33.8	3.064	1	92 18 11.0	19.80	1	F; pL; lbM.....	1
123	2340		Δ. 2??	0 35 50.6	2.329	1	164 7 16.6	19.78	1	i train of st and neb	1
124	54	III. 149		0 36 5.2	3.196	2	60 10 58.3	19.79	2	F; vS; R; lbM	3
125		II. 245		0 36 10.9	3.124	4	76 27 54.3	19.79	4	F; pS; ilE; bM	4
126	2341			0 36 53.3	2.803	1	140 56 42.6	19.78	1	eF; pL; R; gvlbM	1
127	2342			0 38 13.4	2.278	1	164 11 39.2	19.76	1	vF; R	1
128	2343			0 38 18.6	2.275	3	164 12 17.2	19.76	3	vF; S; bi-N.....	3
129	55	III. 485		0 38 45.1	3.005	1	106 20 58.5	19.75	1	vF; S; iR; r; *10, 5' s	3
130		II. 445		0 39 3.7	3.063	1	92 29 13.5	19.75	1	F; pS; iF; er	1
131	56	V. 25		0 40 1.4	3.019	2	102 38 24.1	19.73	2	vF; L; 4st in diff n.....	3
132	57	V. 20		0 40 4.2	2.979	1	111 30 57.1	19.73	1	F; vL; vmE 172°	2*
133	2344			0 40 13.4	2.240	2	164 8 36.1	19.73	2	F; S; E or bi-N; vglbM.....	2
134	2346		Δ. 19? 21?	0 40 17.9	2.255	3	163 51 1.1	19.73	3	F; pL; vLE; r	3
135	58	III. 204?		0 40 22.9	3.154	(2)	71 9 48.1	19.73	1	vF; S; R; lbM; 2vSstf; *inv	3
136	59	II. 609		0 40 32.9	3.195	5	63 8 35.1	19.73	5	pB; S; R; pmbM; r; * p ...	6
137	59, a		R. nova	0 40		...	63 8		...	One of R.'s novæ; the other = h. 60.	0
138	{61= 2345}	V. 1	C.H.	0 40 37.6	2.954	3	116 3 40.4	19.72	3	{!!; vvB; vvL; vmE 54.5 gbM; 4st.}	9*+
139	60			0 40 39.3	3.195	1	63 8 14.4	19.72	1	pF; R; bM	1
140	2347			0 40 40.7	2.920	3	122 11 28.4	19.72	3	vB; pS; lE; smbM; *8, 5' nf	3
141	62	II. 472		0 40 43.5	3.020	1	102 14 39.4	19.72	1	F; pS; R; gbM	2
142	2348			0 40 44.3	2.223	4	164 16 44.4	19.72	4	F; S; R; gbM; *9, 40" nf ...	4
143		II. 863		0 40 47.1	3.105	1	82 27 14.4	19.72	1	pL; lE; gbM; r	1
144	63			0 40 55.7	3.057	1	93 37 22.4	19.72	1	vF; Δ 2st & neb	1
145	64	{II. 703 = II. 621}		0 40 56.1	3.058	1	93 32 40.4	19.72	1	F; S; E 135°±; lbM.....	3*
146	2349		Δ. 3, 4, 21?	0 41 23.8	2.233	3	163 52 6.7	19.71	3	F; pL; R; gbM*13	3
147	2350			0 41 41.6	2.872	1	129 0 20.7	19.71	1	F; S; R; vsymbM*13	1
148	2351			0 42 3.8	2.198	4	164 15 4.0	19.70	4	F; pS; R.....	4
149	65	III. 153		0 42 13.6	3.226	4	58 29 23.0	19.70	4	pB; pS; lE; psbM; r; *8sf4',	5
150	2352			0 42 49.8	2.195	1	164 2 37.3	19.69	1	Cl; F; pL; stvS	1
151	66	III. 463		0 43 1.3	3.046	2	95 57 59.3	19.69	2	vF; pS; ilE; r.....	4
152	2353			0 43 20.8	2.169	1	164 18 1.6	19.68	1	vF; S; R.....	1
153	68	III. 955		0 43 28.7	3.030	(1)	99 25 39.6	19.68	2	pF; vS; iR; pgbM	3
154	67	II. 446		0 43 34.3	3.061	1	92 39 59.6	19.68	1	pF; S; lE; psbM; *8 f5.5...	3
155		III. 430		0 43 47.0	3.038	1	97 39 17.9	19.67	1	vF; vS.....	1
156	69	III. 429		0 43 56.8	3.037	1	97 49 22.9	19.67	1	pB; pS; smbM; sp of Dneb .	3
157	70			0 43 59.1	3.037	1	97 49 2.9	19.67	1	vF; S; R; nf of Dneb	2
158	71	I. 159		0 44 10.2	3.352	1	43 12 15.9	19.67	1	cB; pL; R; 2st10nr	4
159	73	III. 439		0 44 59.5	3.059	2	92 59 10.5	19.65	2	vF; S; iR; bM; stellar	4
160	72	III. 477		0 45 0.3	3.189	1	66 25 23.5	19.65	1	eF; S; R; *15, f30"	3
161	75			0 45 52.1	3.241	1	58 16 59.8	19.64	1	eF; S; R	1
162	{74= 2354}	VI. 20		0 45 52.1	2.932	2	117 20 41.8	19.64	2	⊕; B; L; lE; st 12...16.....	3
163	2355			0 45 57.6	2.903	3	121 57 53.1	19.63	3	vB; L; pmE; glbM; *11np	4
164	2357			0 46 16.9	2.135	1	163 54 48.1	19.63	1	eF	1
165	2356			0 46 33.5	2.115	2	164 6 33.1	19.63	2	Cl; F; eeL; R; st 12...18 ...	2*
166	2358		Δ. 5, 6?	0 47 14.8	2.102	1	164 8 28.7	19.61	1	vF; pL; R; vglbM; r.....	2
167		II. 214		0 47 51.9	+3.242	1	59 11 21.0	-19.60	1	F; E; aB*f, vnr	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
168	2360			0 48 23.0	+ 2.151	4	162° 57' 22.3	- 19.59	4	pB; vS; R; gvlbM; r.....	4
169	2359			0 48 24.5	2.844	3	128 27 8.3	19.59	4	pB; vL; vmiE; vgpmbM ...	5*†
170	76			0 48 59.6	3.131	1	78 40 51.6	19.58	1	Cl; S; sest	1
171	2361			0 49 15.8	2.132	1	163 0 0.9	19.57	1	F; vS	1
172	77			0 49 25.0	3.060	1	92 31 27.9	19.57	1	pF; S; E	1
173	78			0 49 55.7	3.240	2	60 28 21.2	19.56	2	pF; vS; R; gblM.....	2
174	2363			0 50 3.5	2.675	2	143 32 6.2	19.56	2	F; S; R; *12 f 90°	2
175	2362			0 50 10.7	2.883	2	122 43 13.2	19.56	2	eF; vS; R; pB* f 2'	2
176	79	II. 210		0 50 11.5	3.241	3	60 24 25.2	19.56	3	pB; pL; R; gblM; *9, 3' 135°	4
177	79, a, b		R. 2 novæ	0 50 22.3	3.241	...	60 21 31.2	19.56	...	{ F; S; R (ε of Lord R.). }	0*
5059										{ For b, see No. 5059. }	
178	4007			0 50 29.3	2.780	1	134 35 56.5	19.55	1	eF; vS; R; lbM	1*
179	4008			0 50 36.2	2.780	1	134 30 1.5	19.55	1	vF; vS; R; lbM; 3stp	1*
180	2365			0 50 36.7	2.668	2	143 43 58.5	19.55	2	pF; S; R; bM; p of 2	2
181	2364			0 50 40.4	2.810	1	131 12 36.5	19.55	1	(?)F; S; stellar	1
182	2366			0 50 51.5	2.667	1	143 39 58.8	19.54	1	vF; lE; vgbM; f of 2	1
183	2367		Δ. 23	0 51 27.2	2.078	5	162 13 35.1	19.53	5	⊕; vB; S; lE; st 13...15	5
184	2368			0 52 10.5	2.850	2	125 53 21.4	19.52	2	vF; S; R; gblM; 2st 1ls	2
185	80	II. 433		0 52 46.1	3.027	...	98 19 51.0	19.50	...	pF; L; E 0° ±; gblM; *10, f 20°.5.	2
186	2369			0 53 13.3	1.902	1	165 12 27.0	19.50	1	F; L; R; vgbM	1
187	2370		Δ. 25	0 54 19.9	2.043	5	162 56 9.3	19.49	5	B; L; viF; mbMD*; r	5†
188	2371			0 54 45.4	2.632	2	143 59 49.6	19.48	2	eF; S; R	2
189	81	III. 191		0 55 5.2	3.044	2	94 59 49.5	19.45	2	pF; S; iE; *8f97°	4
190	82	II. 434		0 56 16.4	3.032	1	97 5 49.1	19.43	3	F; S; iR; sbM; *14nf 20'...	4
191	2372			0 57 26.8	2.308	1	156 21 30.0	19.40	1	eF; vme 145°.4; vlbM	1
192	2374		Δ. 55 ??	0 57 28.7	2.022	2	162 22 33.0	19.40	2	vF; pL; vLE; vgbM	2
193	2375		Δ. 62	0 57 28.8	2.069	3	161 35 59.0	19.40	3	⊕; vB; vL; vC; vmbM; st 13...14.	4
194	2373			0 57 44.3	2.828	2	125 53 46.0	19.40	2	F; S; R; gblM	2
195	83			0 57 45.8	3.700	1	28 34 5.0	19.40	1	Cl; S	1
196	4012			0 57 57.6	2.745	1	134 1 50.3	19.39	1	eF; vS; *7.8 sp 3'	1*
197			D'Arrest, 7	0 58 53	3.28	[1]	58 20 18	19.37	[1]	vF; *13, s 15'; m diff	0
198	2376		Δ. 31 ??	0 58 56.4	1.969	5	162 48 27.9	19.37	5	Cl; F; L; R; pC; st 14...16.	5
199			D'Arrest, 8	0 59 21	3.29	[1]	57 57 18	19.36	...	F; S; bet 2 st 15.....	0
200	2378		Δ. 36 ??	0 59 27.3	1.904	2	163 34 23.2	19.36	2	⊕; B; S; R	2
201	2377			0 59 31.7	2.864	1	120 55 53.2	19.36	1	vF; S; R; gblM	1
202	84	II. 215		0 59 32.3	3.289	3	58 14 9.2	19.36	3	pF; S; R; bM; 1st of 3	4*
203	85	II. 216		0 59 34.3	3.289	3	58 16 41.2	19.36	3	pF; S; R; sbM; 2nd of 3 ...	4*
204		VIII. 64	C.H.	0 59 37.4	3.703	1	29 9 36.2	19.36	1	Cl; pC	1
205	86, a		R. nova	0 59 39.6	3.289	...	58 20 48.2	19.36	...	γ' in Lord R.'s diagram	0*
206	86	II. 217		0 59 40.5	3.289	3	58 20 16.2	19.36	3	pF; pL; R; gblM; 3rd of 3...	4*
207			D'Arrest, 9	0 59 42	3.29	[3]	58 26 0	19.36	[3]	p of Dneb; vF; pS; { Δ.R.A.=0 } { Δ.P.D.=93'' }	0*
208			D'Arrest, 10	0 59 42	3.29	[3]	58 25 18	19.36	[3]	vF; R; pS; f of D neb	0*
209	86, b		R. nova	0 59 46.3	3.288	...	58 23 20.2	19.35	...	δ in Lord R.'s diagram	0*
210	86, c		R. nova	1 0 2.3	3.288	...	58 27 16.2	19.34	...	θ in Lord R.'s diagram	0*
211			Auw. N. 9	1 0 15.1	3.071	...	89 48 55.8	19.34	...	F nebula (Bond, Jan. 1853).	0*
212	87	II. 218		1 0 39.6	3.299	1	57 37 10.1	19.33	1	F; vS; R; mbM; bet 2 st	2
213	87, a		R. nova	1 0		...	59 37		...	makes a D neb with h. 87 ...	0
214	88	I. 54		1 0 43.4	3.361	2	51 5 25.1	19.33	2	F; vS; vLE; gblM; 4Sstnr ...	4*
215			D'Arrest, 11	1 0 45	3.30	[1]	57 36 18	19.33	[1]	F; S; pos from h. 87=40°; dist 47''.	0
216	2379			1 0 45.8	1.940	1	162 44 38.1	19.33	1	vF; pL; R; gblM	1
217			D'Arrest, 12	1 1 29	3.30	[1]	57 59 48	19.31	[1]	vF; S; R; *10f 1°.8, s 80'' ...	0
218	89	II. 224		1 1 39.0	3.326	2	55 2 13.7	19.31	2	pB; eL; R; gblM; β Androm.nr	3
219	2380			1 2 5.5	2.679	1	137 25 34.0	19.30	1	eS; stellar; = *7m	1
220	2381			1 2 39.6	2.041	1	160 37 28.3	19.29	1	F; vL; R; vglbM	1
221		II. 219		1 2 44.2	3.306	(1)	57 34 0.3	19.29	(1)	eS; F; p of D neb	1
222		II. 220		1 2 44.2	+ 3.306	(1)	57 34 0.3	-19.29	(1)	pL; f of D neb	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and ll.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. II.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
223	2382			1 3 9.2	+2.800	1	126 31 25.9	-19.27	1	eF; S; R; vS* nr	1
224	2384			1 3 27.6	1.908	1	162 30 42.9	19.27	1	eF; pL; R; gvlbM	2
225	2383			1 3 37.2	2.801	2	126 14 35.2	19.26	2	vF; S; R; glbM	2
226	2386			1 3 39.2	1.861	4	163 6 23.2	19.26	4	F; pS; R; gbM	4
227	2385			1 3 57.5	2.849	3	120 57 48.5	19.25	3	F; pL; R; vglbM; p of 2	3
228	2387		Δ. 36 ?	1 4 4.2	1.814	1	163 37 59.5	19.25	1		4
229	90	III. 15		1 4 25.7	3.303	1	58 37 12.8	19.24	1	F; pS; R; bM	2
230		III. 15 ⁴		1 4 26.1	3.303	1::	58 36 22.8	19.24	1::	eF; vS	1
231	[162]			1 4 28.3	1.890	1	162 30 54.8	19.24	1	vF	1
232	2388			1 4 41.6	2.855	2	119 58 44.8	19.24	2	eF; S; E; glbM; f of 2	2
233	2389			1 4 57.6	2.768	1	128 49 39.1	19.23	1	vF; S; R; glbM	1
234	91	III. 59 ²		1 5 41.2	3.066	1	91 3 7.7	19.21	1	vF; vS; R	2
235	91, a		R. nova	1 5			91 3			Forms a Δ with h. 91 & 92...	0
236	2390			1 5 43.8	2.826	2	122 49 44.7	19.21	2	2 vSst + F neb	2
237	92	III. 593		1 5 46.4	3.065	1	91 4 22.7	19.21	1	eF; eS	2
238		II. 622		1 5 49.2	3.074	1	89 45 44.0	19.20	1	F; L; R; bM; er	1
239	93	II. 447		1 5 51.7	3.066	1	90 59 22.7	19.21	1	F; vS; R; vsbM*	3
240	95			1 6 16.1	3.324	1	57 1 50.0	19.20	1	F; S; vsbM	1
241	2391			1 6 16.3	2.340	1	152 20 39.0	19.20	1	F; S; R; gbM; *12 f	2
242	94			1 6 27.6	3.732	1	30 36 52.3	19.19	1	Cl; S; IC	1
243	2392			1 6 36.9	2.430	2	148 59 52.3	19.19	2	B; S; R; psbM	2
244		VII. 45		1 6 53.4	3.703	2	31 56 6.6	19.18	2	Cl; S; iF; pC	2
245	2393			1 7 7.0	2.761	2	128 38 59.9	19.17	2	pF; S; R; glbM	2
246	2394			1 7 12.4	2.823	2	122 29 25.9	19.17	2	pB; S; R; gbM	2
247	2396			1 7 13.1	2.423	2	149 1 44.9	19.17	2	F; vS; R	2
248	2395			1 7 16.6	2.823	2	122 32 55.9	19.17	2	pF; S; R; gbM	2
249			D'Arrest, 13	1 7 21	3.32	[1]	57 31 18	19.16	[1]	F; S; R; *15 p, 8 ^s .3, 270°	0
250			D'Arrest, 14	1 7 51	3.32	[2]	57 40 42	19.15	[2]	F; pL; bM; *11 nr	0
251	96, a		R. nova	1 8			59 43			vF; mE135°+; lbM; nph.96	0
252	96			1 8 29.3	3.306	1	59 42 41.8	19.14	1	vF; E; *9np; S*nf, vnr	1
253	2397			1 8 35.1	2.480	1	146 8 37.8	19.14	1	vF; S; R; bM	1
254		III. 440		1 8 48.8	3.061	1	91 35 19.1	19.13	1	vF; vL	1
255	2399		Δ. 7, 10 ?	1 9 50.6	1.668	3	164 2 15.0	19.10	4	pF; pL; iR; r; 1st of sev, neb and st.	3
256	97	VII. 42		1 10 24.8	3.722	1	32 24 35.3	19.09	1	Cl; B; L; pR; st 7, 8, 10	3
257	2401		Δ. 60 ?	1 10 40.8	1.799	2	162 17 13.6	19.08	2	pF; L; R; vglbM	2
258		III. 205		1 10 41.4	3.198	1	73 4 51.6	19.08	1	eF	1
259	2402		Δ. 8, 10 ?	1 10 48.6	1.650	4	164 2 41.6	19.08	4	F; pL; iR; gbM; r; 2nd of sev.	4
260	2400			1 10 54.4	2.794	1	124 6 11.6	19.08	1	pB; R; glbM; ? 1° in P.D.	1
261	2404		Δ. 9 ?	1 11 42.3	1.629	1	164 4 36.5	19.05	1	pB; pL; iF; 3rd of sev.	1
262	2403			1 11 43.3	2.365	1	149 38 46.5	19.05	1	vF; R; gbM; 30	1
263	99	III. 250		1 11 56.6	3.091	1	87 25 16.5	19.05	1	pB; pL; R; gmbM; p of 2	4
264		I. 108		1 12 1.0	3.091	1	87 30 53.5	19.05	1	eB; vL; iR; pB*f	1
265	98			1 12 2.9	3.335	1	58 1 55.5	19.05	1	vF; eS; stellar	1
266			D'Arrest, 15	1 12 41	3.33	[2]	58 2 0	19.03	[2]	eF; vS; *9p14 ^s ; v diffie	0
267		III. 206		1 12 41.2	3.192	1	74 14 55.1	19.03	1	eF; S	1
268	100	III. 577		1 13 19.9	3.428	1	50 14 18.7	19.01	1	vF; pS; vIE; vglbM	2
269		III. 251		1 13 21.4	3.092	3:	87 22 9.7	19.01	3:	pB; S; smbM; f of 2	3
270	101			1 13 47.2	3.119	1::	83 43 2	19.00	1	eF; pL; R; red *7.8, 225°	1
271	2405			1 14 3.5	2.691	1	131 42 52.3	18.99	1	eF; iE	1
272	102	III. 156		1 14 6.0	3.350	1	57 17 30.3	18.99	1::	vF; eS; 1st of 3	3
273	2406			1 14 7.3	2.353	2	149 15 33.3	18.99	2	vB; S; iE; psmbM	2
274			D'Arrest, 16	1 14 29	3.14	[1]	81 31 48	18.97	[1]	pB; S; E	0
275	103, a		R. nova	1 14 29	3.107		85 24 26.6	18.98		γ in Lord R.'s diagram	0
276	103	III. 252		1 14 31.2	3.107	2	85 28 25.6	18.98	2	pB; L; R; svmbM; *7 f1 ^m	3
277	103, b		R. nova	1 14 46.4	3.107		85 21 47.6	18.98		β in Lord R.'s diagram	0
278		III. 157		1 14 51.3	3.351	2	57 15 28.9	18.97	2	vF; S; 2nd of 3	2
279	2407			1 14 53.5	2.772	2	124 48 15.9	18.97	2	B; S; vIE; bM; vS*nr	2
280	103, c		R. nova	1 14 57.1	3.107		85 18 37.6	18.97		δ in Lord R.'s diagram	0
281	105	III. 594		1 14 59.5	3.075	1	89 47 7.2	18.96	1	vF; L; mE 60°±; lbM	2
282	104			1 15 4.4	+3.352	1	57 33 18.2	-18.96	1	vF; E; *s	1

No. of Cata- logue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and ll.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s						
.....	104, <i>a</i>		R. 5 novæ	1 15	+.....	...	57 33 "	-18°95	...	No deser Nos. 283...287 incl	0
288	106, <i>a</i>		R. nova	1 15		...	57 16	18°95	...	No description	0
289	106	III. 158		1 15 19.6	3.354	2	57 16 46.5	18°95	2	pB; pL; R; 3rd of 3	4
290	103, <i>d</i>		R. nova	1 15 23.2	3.107	...	85 20 17.6	18°95	...	ε in Lord R.'s diagram	0
291			D'Arrest, 17	1 15 35	3.35	[3]	57 31 18	18°94	[3]	vF; S; obs. with H. 157, 158, 159, 160.	0
292	107			1 15 36.6	3.352	1:	57 36 31.8	18°94	1:	No description	1
293			D'Arrest, 18	1 15 38	3.13	[2]	81 41 6	18°94	[2]	eB; S; R; bMN	0
294	108	III. 159		1 15 48.0	3.354	2	57 28 37.8	18°94	2	vF; pL; R; bM; p of 2	3
295	109	III. 160		1 15 49.0	3.354	1	57 26 52.8	18°94	1	vF; S; f of 2	2
296	110			1 16 7.8	3.362	1	56 49 53.1	18°93	1	vF; vS	1
297	111	III. 169		1 16 34.4	3.362	1	56 56 28.4	18°92	1	F; S; stellar	2
298	112	II. 252		1 16 35.8	3.169	2	77 49 10.7	18°91	2	F; L; lE; vglbM; * f.	3+
299	113	III. 167		1 16 45.3	3.360	1	57 16 37.7	18°91	1	Stellar; p of 2	2
300			D'Arrest, 19	1 16 48	3.14	[2]	81 10 6	18°91	[2]	eF; S; v diffie; I 151 f41 ^s	0
301	114	III. 168		1 16 48.8	3.360	1	57 18 37.7	18°91	1	pB; R; stellar; f of 2	2
302	114, <i>a</i>		R. nova	1 16		...	57 18		...	S; R; bM	0
303	116	III. 253		1 17 19.4	3.097	1	86 55 16.3	18°89	1	F; cL; E 135°+	3+
304	115	II. 461		1 17 19.8	3.078	2	89 0 21.3	18°89	2	F; pL; R; gbM	5
305			D'Arrest, 20	1 17 22	3.14	[2]	80 44 24	18°89	[2]	eF; pL; iF; ?Cl+neb	0
306			D'Arrest, 21	1 17 25	3.37	[1]	56 42 18	18°89	[1]	D neb; vF; 90° pos	0
307	117	I. 151		1 17 26.8	3.142	1	81 11 50.3	18°89	1	vB; pL; mbM; 4S st nr	2
308			D'Arrest, 22	1 17 31	3.14	[2]	81 1 12	18°88	[2]	vF; vS; * 11.12 p 5 ^s	0
309	2408			1 17 35.1	2.750	3	125 48 8.3	18°89	3	F; S; lE; p of 2	3
310	2409			1 17 37.5	2.748	3	125 51 0.3	18°89	3	F; S; lE; bM; f of 2	3
311	118			1 17 44.5	3.377	1	56 1 31.3	18°89	1	pB; vS; sbM; p of 2	1*
312	118, <i>a</i>		R. nova	1 17		...	56 1		...	One of 4 neb nr h. 120	0
313		III. 556		1 17 51.6	3.140	1	81 28 43.6	18°88	1	vF; pL; mE 15°+	1*
314	119			1 17 51.6	3.140	(1)?	81 29 30.6	18°88	1	Not vF; L; R; bM	2*
315	121	II. 462		1 18 19.5	3.081	2	88 58 4.9	18°87	2	pB; pL; R; gmbM	3
316	2410			1 18 22.9	2.708	2	128 52 11.9	18°87	2	eeF; S; R; vgbM; 1st of 4...	2
317	120			1 18 24.6	3.379	1	56 2 19	18°87	1	pB; pL; gbM; f of 2	1*
318	120, <i>a</i>		R. nova	1 18		...	56 2		...	One of 4, see h. 118, 120	0
319		III. 170		1 18 26.6	3.373	1	56 39 5.2	18°86	1	Stellar	1*
320	2411			1 18 51.2	2.707	2	128 48 57.5	18°85	2	eeF; S; R; vgbM; 2nd of 4...	2
321	2412			1 18 52.4	2.707	2	128 47 17.5	18°85	2	eeF; S; R; vgbM; 3rd of 4...	2
322		II. 448		1 18 55.2	3.056	1	92 3 54.5	18°85	1	Stellar; p of D neb	1
323		II. 449		1 18 55.2	3.056	1	92 3 54.5	18°85	1	Stellar; f of D neb	1
324	2413			1 19 6.8	2.707	1	128 44 40.8	18°84	1	eeF; S; R; vgbM; 4th of 4...	1
325		III. 171		1 19 27.2	3.383	1	56 4 7.1	18°83	1	Stellar	1*
326	122	II. 463		1 19 33.1	3.083	2	88 42 53.1	18°83	2	F; S; E90°; bM; r	5
327	123	III. 560		1 19 42.9	3.415	1	53 32 28.1	18°83	1	vF; S; E; vglbM; *13 nr	2
328		III. 172		1 19 56.5	3.371	1	57 16 7.4	18°82	1	vS; stellar; p of 2	1
329		III. 173		1 19 56.5	3.371	1	57 16 7.4	18°82	1	vS; stellar; f of 2	1
330	124	VII. 48		1 20 10.3	3.969	1	27 25 54.7	18°81	1	Cl; B; pL; pRi; st mm	2*
331			D'Arrest, 23	1 20 21	3.38	[1]	56 25 18	18°80	[1]	eF; pL; R	0
332		III. 441		1 20 34.8	3.051	1	92 40 8.0	18°80	1	vF; vS; iE; p of 2	1
333		III. 442		1 20 39.9	3.052	1	92 37 8.0	18°80	1	vF; vS; iF; f of 2	1
334	125			1 21 10.0	3.362	1	58 23 57.6	18°78	1	vF; S; R	1
335	2414			1 21 37.4	2.726	1	126 27 1.2	18°76	1	vF; S; R	1
336	2415			1 22 20.5	2.675	1	130 2 23.8	18°74	1	eF; S; att to S*; B*nr	1
337	2416			1 22 43.5	2.723	1	126 19 27.1	18°73	2	vS; *pos 225° inv	3
338	2417			1 23 15.2	2.453	1	142 18 52.7	18°71	1	F; S; R; bM; am st 11	1
339	2418			1 23 43.9	2.864	1	113 23 44.0	18°70	1	B; L; pmE; gpmbM	1
340	127			1 23 47.8	3.390	1	57 6 32.0	18°70	1	vF; pL; gbM	1
341	126		Σ.131=M.103	1 23 59.8	3.916	2	30 2 9.3	18°69	2	Cl; B; R; Ri; pL; st 10...11	5
342	128	I. 100		1 24 18.7	3.008	3	97 35 25.6	18°68	3	vB; pL; R; mbM; p of 2	4
343	128, <i>a</i>		R. nova	1 24		...	97 35		...	No description	0
344		III. 431		1 24 22.2	3.008	(1)	97 36 39.6	18°68	(1)	eF; S; f of 2	1
345	129			1 24 33.4	3.059	1	91 38 46.6	18°68	1	vF; S; R; bM	1
346	130			1 24 35.7	+3.008	1	97 36 47.9	-18°67	1	vF; vS; R	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. II.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
347			D'Arrest, 24	1 24 40	+3.41	[1]	55 26 18	-18.66	[1]	vvF; S; ?rr	0
348			D'Arrest, 25	1 25 0	3.35	[2]	60 5 0	18.65	[2]	F; M. 33, f65 ^a ; another f28 ^a ..	0
349			D'Arrest, 26	1 25 28	3.35	[2]	60 5 0	18.64	[2]	F; pL, f of 2.....	0
350	2419			1 25 53.4	2.737	2	124 13 32.1	18.63	2	F; S; R; bM	2
351	132	II. 4		1 25 54.1	3.006	1	97 44 13.1	18.63	1	pB; R; bM; r; *6, f47 ^a .5 ...	7
352	131	V. 17	M. 33	1 25 56.3	3.358	1	60 3 50.1	18.63	1	!; eB; eL; R; vRi; vgbMN; rr.	14†
353		II. 473		1 26 1.4	2.960	1	102 53 18.1	18.63	1	F; S; iF; er.....	1
354		III. 432		1 26 15.0	3.003	1	98 2 38.4	18.62	1	eeF	1
355	133	III. 150		1 26 17.0	3.361	1	59 57 47.4	18.62	1	vS; R; vvlbM	5
356	2421		Δ. 17?	1 26 24.6	1.327	2	164 16 37.4	18.62	2	B; S; R; psbM*; r.....	2
357			R. nova	1 26 33.5	3.355	::	60 31 53.9	18.61	::	S, a neb or Cl with 3stin v ...	0
358			{ D'Arr. = Auw.N.15 }	1 27 18.1	3.002	...	98 5 42.6	18.58	...	{ Nebulous *11, m (D'Arr. Resultate). }	0*
359	134			1 27 23.6	3.399	1	57 4 11.6	18.58	1	vF; psbM; stellar	1
360	2423			1 27 44.5	2.690	1	127 13 26.9	18.57	1	F; vS; R; *12, p	1
361	{ 139 = 2422 }	I. 281		1 27 47.4	2.780	3	120 7 37.9	18.57	3	{ vB; vL; vmE, 118°3; } sbM; *34°5, 6°5. }	5
362	135	III. 174		1 27 53.1	3.401	1	57 2 16.9	18.57	1	pF; psbM; stellar	2
363	137	II. 282		1 28 18.0	3.001	1:	98 2 43.5	18.55	1	pB; pL; iLE; gmbM; r; *8, np10'.	4
364	136			1 28 19.5	3.399	1	57 19 26.5	18.55	1	pB; pL; bM; *f, 2 ^m 51 ^s	1
365	2424			1 28 37.9	2.687	1	127 12 19.8	18.54	1	eeeF; vS; R; p of 2.....	1
366	138	III. 454		1 28 48.9	3.072	1	90 2 25.1	18.53	2	eF; pL; not bM	2
367	2425			1 28 50.4	2.686	1	127 12 40.1	18.53	1	F; S; R; f of 2	1
368	140	III. 471		1 28 55.4	2.976	1	100 43 34.1	18.53	1	eF; S; am vSst	2
369	2426		Δ. 479	1 28 57.2	2.612	1	132 9 23.1	18.53	2	pB; pL; mE; gpmbM	3
370	2427			1 29 2.6	2.647	1	129 51 50.1	18.53	1	pF; S; R; bM.....	1
371	141			1 29 9.2	3.405	1?	57 7 44.4	18.52	1::	vF; R; f of 2	1
372	142		M. 74	1 29 11.1	3.211	2	74 55 46.4	18.52	2	⊕; F; vL; R; vg, psmbM; rr	11†
373			Auw. N. 16	1 29 14.4	4.681	...	17 49 45.0	18.50	...	iF; 3st + neb (Struve, Σ. 2)...	0
374	2428			1 29 27.8	2.642	1	130 3 42.7	18.51	1	pF; S; R; bM.....	1
375	143			1 29 59.6	3.119	1	84 50 14.0	18.50	1	pB; S; R; psbM.....	1
376	2429			1 30 9.9	2.669	2	128 2 13.0	18.50	2	pB; S; R; gbM; *np.....	2
377	144	II. 283		1 32 7.6	2.997	2	98 13 12.4	18.42	2	pB; vS; R; mbM; r	4
378		VII. 49		1 32 8.2	4.132	1	26 40 32.4	18.42	1	Cl; pS; L & vSst.....	1
379	2430			1 32 32.0	2.759	3	120 38 10.7	18.41	3	vF; vS; p of 2	3
380	2432			1 32 38.7	2.575	2	133 14 25.0	18.40	3	F; S; R; gpmbM; p of 2 ...	3
381	2431			1 32 39.3	2.759	3	120 37 15.0	18.40	3	vF; pS; R; gbM; *f, nr.....	3
382	2435			1 32 40.7	0.929	2	166 16 4.0	18.40	2	vF; pS; R; vglbM	2
383	2433			1 32 50.9	2.573	2	133 18 22.0	18.40	2	F; S; vLE; glbM; f of 2	2
384	2434			1 32 55.1	1.907	1	155 36 37.3	18.39	1	vF; iR; vglbM	1
385			M. 76	1 33 28.5	3.334	1	39 8 52.4	18.37	1	vB; p of D neb	2
386		I. 193		1 33 37.5	3.734	1	39 7 27.4	18.37	1	vB; f of D neb	1
387	145	VII. 46		1 34 28.0	4.062	2	28 49 19.8	18.34	2	Cl; iF; Ri; one * 6.7; st 11 ...14.	4
388	146			1 34 41.6	3.856	1	34 49 52.1	18.33	1	Cl; pRi; st12, m.....	1
389		VIII. 65	C.H.	1 34 45.0	4.018	1	30 0 37.1	18.33	1	Cl; S; iRi; stL	1
390		II. 253		1 35 35.6	3.199	1	77 4 36.0	18.30	1	pB; pL; E; bM; r	1
391	147	II. 610		1 36 23.8	3.366	1	62 0 24.9	18.27	1	F; S; R; bM; r	3
392		VI. 31		1 36 29.2	4.055	1	29 27 40.9	18.27	1	Cl; B; L; eR; st pL	1
393	148			1 36 30.8	3.107	1	86 28 36.9	18.27	1	vF; S; R	1
394		II. 588		1 37 15.0	3.165	2	80 17 10.8	18.24	2	F; S; iE; bM; r	2
395	149	II. 611		1 39 33.1	3.365	2	62 48 58.2	18.16	2	F; S; iE 0°+	3
396	150	I. 157		1 40 0.8	3.360	2	63 16 25.8	18.14	3	F; E 0°...90°; bet 2st.....	4
397		II. 589		1 40 35.6	3.183	2	79 10 46.7	18.11	2	F; pL; E; lbM; *nf, 2'	2
398			D'Arrest, 27	1 41 29	3.30	[1]	68 21 3	18.08	[1]	pB; vmE; *14, f8 ^a	0
399		II. 228		1 41 34.5	3.299	2	68 42 50.6	18.08	2	pB; S; iR; mbM; 1st of 2 ...	2
400	151	IV. 42		1 41 37.9	3.126	2	84 47 12.6	18.08	2	vF; vmE, 165°±; sbM*9 ...	3†
401		III. 175		1 41 51.1	+3.482	1	54 53 51.9	-18.07	1	F; stellar	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. II.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s		° ' "	"			
402		II. 229		1 41 57.7	+3.299	2	68 44 49.2	-18.06	2	pB; S; iR; mbM; 2nd of 2...	2
403	152	II. 612		1 42 34.5	3.369	(1)	63 3 33.2	18.06	1	F; vIE, 90°; *15, nr	2
404	2436	II. 481		1 42 17.0	2.959	1	101 7 2.5	18.05	1	pF; cL; R; glbM; S*p, 90" ..	2
405	153			1 42 17.6	3.186	1::	79 0 17.5	18.05	1	eF	1
406	154	II. 501		1 42 18.3	2.910	1	105 39 54.5	18.05	1	cF; S; R; gvlbMN	2
407	2438			1 42 20.7	2.290	1	143 29 3.5	18.05	1	F; vL; R; vgvlbM	1
408	{ 155 = 2437 }	III. 459		1 42 21.3	2.808	1	114 29 31.5	18.05	1	vF; vS; R; gbM; er; 2stnr ..	4
409		III. 561		1 42 25.7	3.492	1	54 21 52.8	18.04	1	vF; stellar	1
410		II. 617		1 42 55.7	3.299	1	68 56 54.1	18.03	1	F; cL; vgblM	1
411	2439			1 43 2.1	2.393	2	139 19 59.1	18.03	2	B; S; R; gbM	2
412	156	II. 859		1 43 10.9	3.129	1	84 33 6.4	18.02	1	pF; S; E90°; vgblM; *10, nf	2†
413			D'Arrest, 28	1 43 13	3.30	[1]	68 42 18	18.01	[1]	F; S; R; bet 2 st 15	0
414		II. 618		1 43 29.0	3.309	1	68 7 54.7	18.01	1	vS; stellar	1
415	2440			1 43 32.7	2.652	1	125 38 52.7	18.01	1	F; S; R	1
416		III. 179		1 43 39.3	3.307	2	68 20 25.0	18.00	2	F; cL; E; mbM	2
417	2441			1 43 42.0	2.653	1	125 34 5.0	18.00	1	eF; S	1
418	{ 160 = 2442 }	I. 62		1 44 8.0	2.964	1	100 23 45.6	17.98	2	F; pL; E; vgvlbM; r	4*
419	158	III. 192		1 44 17.0	3.024	2	94 44 57.9	17.97	3	eF; vIE 0° ±, *13, s, 90"	4
420		III. 564		1 44 25.9	3.496	1	54 35 56.9	17.97	1	Stellar; 3rd of 4	1
421		III. 565		1 44 25.9	3.496	1	54 35 56.9	17.97	1	Stellar; last of 4	1
422	157	III. 562		1 44 27.3	3.497	1	54 31 58.9	17.97	1	vF; stellar; 1st of 4	2
423						...			...		
424	157, a		R. 3 novæ	1 44		...	54 32		...	Near h. 157, 159	0
425						...			...		
426	161	II. 596		1 44 30.7	3.131	1	84 24 17.9	17.97	1	F; S; bM; *13 1', n	3
427	159	III. 563		1 44 34.3	3.497	1	54 33 1.9	17.97	1	F; pL; bM; 2nd of 4	2
428	162		55 Androm.	1 44 55.9	3.575	1	49 57 41.5	17.95	1	Fine nebulous * with strong atm.	1*
429	163			1 44 56.2	3.509	1	53 52 9.5	17.95	1	vF; R; am pBst	1
430	164	II. 270		1 45 56.9	3.110	2	86 29 23.7	17.91	2	pB; S; iR; psmbM	3
431	{ 165 = 2443 }	I. 105		1 46 12.4	2.918	2	104 25 39.0	17.90	2	cB; pL; lE; psmbM	3
432			D'Arrest, 29	1 46 19	3.28	[1]	70 50 18	17.89	[1]	eF; R; *19, f	0
433			D'Arrest, 30	1 46 32	3.56	[1]	51 18 3	17.88	[1]	eF; pL	0
434			D'Arrest, 31	1 47 5	3.29	[3]	69 59 18	17.86	[3]	vF; vS; R; β Arietis in field	0
435	{ 166 = 2444 }	III. 460		1 47 13.9	2.797	2	114 26 46.2	17.86	2	pF; vS; R; vgblM	3
436	167			1 47 14.6	2.795	1	114 33 21.2	17.86	1	vF; pL; R; gbM; S*195° ..	1
437	2445			1 47 43.1	2.621	1	126 34 24.8	17.84	1	F; S; R; bM	1
438	168			1 47 47.9	3.110	1	86 29 14.8	17.84	1	Suspected neb	1
439	2446			1 47 50.1	2.621	1	126 32 0.8	17.84	1	eeeF; S; R	1
440		III. 266		1 48 6.0	2.969	1	99 38 5.4	17.82	1	eF; stellar	1
441	{ 177 = 2447 }	III. 265		1 48 17.0	2.968	1	99 44 29.4	17.82	1	vF; pS; vIE	3
442		II. 221		1 48 26.0	3.463	1	57 38 16.7	17.81	1	F; pL; mE; r	1*
443		III. 176		1 48 33.4	3.483	1	56 23 6.7	17.81	1	eeeF; stellar	1
444	169			1 48 33.9	3.463	3	57 38 53.7	17.81	3	pB; R; bM; *13, np ..	3*
445	169, a		R. nova	1 48 34.5	3.463	...	57 38 25.7	17.81	} { β, γ, δ of Lord R.'s diag. ε=II. 221 H.	} 0*	} 0*
446	169, b		R. nova	1 48 38.4	3.463	...	57 38 0.7	17.81			
447	169, c		R. nova	1 48 47.9	3.463	...	57 40 30.7	17.81			
448	172	II. 271		1 49 2.8	3.126	1	85 3 29.3	17.79	1	pF; S; R; p of 2; pos=102° 4	1
449	173	II. 272		1 49 3.2	3.126	1	85 3 41.3	17.79	1	vF; vS; R; sbM; f of 2	4
450	170			1 49 6.3	4.127	1	30 30 49.6	17.78	1	Cl; not Ri; *	1
451	171			1 49 7.6	+3.953	1	35 13 13.6	-17.78	1	Cl; pL; pRi; iF; st 11...13..	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
452	2449			1 49 12.4	+2.112	1	147° 22' 44.6	-17.78	1	pB; S; R; gbM	1
453	176	III. 193		1 49 20.4	3.017	2	95 9 13.6	17.78	2	eF; *9, 315°±	3
454	2448			1 49 21.5	2.707	3	120 36 23.6	17.78	3	pB; S; E; IM	3
455	175	II. 222		1 49 23.6	3.468	1	57 29 41.6	17.78	1	F; pL; mE; r; f of 2	3
456	175, a		R. nova	1 49		...	57 29		...	nf h. 175	0
457	174	VII. 32		1 49 25.8	3.541	1	53 1 14.9	17.77	1	Cl; vL; Ri; st L and sc ...	5
458	2450			1 49 27.7	2.107	1	147 26 49.9	17.77	1	vF; S; R; bM	1
459	{ 178 = 2451 }	III. 464		1 49 57.4	3.006	3	96 5 19.5	17.75	3	vF; S; lE; vglbM	4
460	180			1 51 17.8	3.158	1	82 20 13.3	17.69	1	vF; S; R; *10, 2' 285°	1
461			D'Arrest, 32	1 51 35	3.28	[2]	71 42 48	17.68	[2]	vF; S; R; nr I. 112 H.	0
462	179		50 Cassiop.	1 51 36.4	5.005	1	18 15 36.6	17.68	1	Suspected nebulous *	1*
463	181	112		1 51 39.0	3.283	2	71 40 12.6	17.68	2	B; cL; R; gbM; r	3
464	181, a		R. nova	1 51 39	3.283	::	71 45±		::	5 or 6' s of h. 181	0
465	2452	III. 468		1 52 0.8	2.936	1	102 10 50.9	17.67	1	cF; pL; E 0°±; glbM	2
466		III. 214		1 52 2.0	3.223	1	76 41 15.9	17.67	1	vF; stellar	1
467	2453			1 52 3.7	2.751	2	116 58 27.9	17.67	2	pF; S; R; glbM	1
468			D'Arrest, 33	1 52 5	3.34	[1]	67 2 18	17.66	[1]	F; pL	0
469	182	II. 223		1 52 10.0	3.451	1	59 15 3.2	17.66	1	pB; pL; R; glbM	2
470	183	I. 101		1 52 40.6	2.999	3	96 38 24.8	17.64	3	cB; L; mE 163°0; mbM ...	6
471		III. 215		1 52 43.3	3.209	1	77 59 17.8	17.64	1	eF; stellar	1
472	184	III. 583		1 52 52.3	3.406	1	62 26 29.1	17.63	1	vF; vS; E; 3 stp; *250° ...	2*
473	2454			1 52 57.3	2.039	1	148 28 9.1	17.63	1	pB; pL; lE; *12 att	1
474	185	II. 435		1 54 2.1	2.989	(1)	97 30 22.6	17.58	1	pF; pS; R; bM	2
475	186	III. 433		1 54 20.6	3.005	1	96 3 38.9	17.57	1	cF; cS; R; bM	3
476			D'Arrest, 34	1 54 25	3.16	[1]	82 10 18	17.56	[1]	vF; S; *14 f 90°; 11.65 ...	0
477	187			1 54 39.4	3.247	1	74 57 35.2	17.56	1	eF; S; R; *11 75°	1
478	188	III. 207		1 54 49.2	3.280	1	72 17 54.5	17.55	1	vF; cS; R; stellar	2
479	2455			1 54 55.4	2.100	1	146 30 26.8	17.54	1	pF; S; R; 2 st 11, nr	1
480	2456			1 55 1.3	0.699	3	164 54 18.8	17.54	3	eF; vS; R; *12, 25" 315° ...	3
481	189	III. 566		1 55 8.5	3.571	1	52 33 52.1	17.53	1	vF; S; iR; sbM; *nr	3
482	2457			1 55 46.1	1.428	1	158 32 40.7	17.51	1	eeF; vS; R; *13 p 100" ...	1
483	190	III. 208		1 56 8.7	3.253	1	74 38 1.3	17.49	1	vF; S; iR; glbM; *10 p 3.5	2
484	191	III. 151		1 56 54.9	3.428	1	61 40 47.2	17.46	1	vF; vS; iR; bet 2stn and sp..	3
485	{ 192 = 2458 }			1 57 28.5	2.780	2	113 58 18.1	17.43	2	vF; pS; vLE	2
486	2459			1 58 26.4	1.343	1	159 7 19.3	17.39	1	pF; S; R; gbM	1
487	193	I. 152		1 59 51.2	3.196	1	79 40 47.1	17.33	1	B; vS; vLE; svmbM; *10, 55" 320°	3*
488	194	II. 604		2 0 19.0	3.604	1	51 54 21.7	17.31	1	pB; cL; lE; mbM	2
489	195			2 0 49.6	3.238	1	76 18 58.6	17.28	1	F; R; vS; bM	1
490	2461			2 0 55.4	2.468	2	131 49 14.6	17.28	2	cF; vS; R; sbM; r	2
491	{ 196 = 2640 }			2 0 56.9	2.741	2	116 7 18.6	17.28	2	vF; vF *inv	2
492	2462			2 0 58.2	2.561	1	127 9 11.6	17.28	2	F; S; R; vsvmbM*13	2
493	198	III. 227		2 1 34.0	3.161	1	82 41 44.2	17.26	1	vF; S; R; bM; am st	2
494	197	II. 605		2 1 40.3	3.617	1	51 28 53.5	17.25	1	pB; S; iR; *f 15°	2
495	{ 199 = 2463 }	II. 482		2 2 30.1	2.942	3	100 47 38.7	17.21	3	F; S; R; 1st of 4	4
496		III. 567		2 2 32.5	3.591	1	53 0 41.7	17.21	1	vF; S; lE	1
497	{ 200 = 2464 }	II. 483		2 2 32.8	2.942	3	100 47 50.7	17.21	3	F; S; R; 2nd of 4	4
498	{ 201 = 2465 }	II. 484		2 2 47.1	+2.942	3	100 48 31.0	-17.20	3	vF; vS; R; 3rd of 4	4

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by I. and II.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s		° ' "	"			
499	$\left\{ \begin{array}{l} 202 \\ = \\ 2466 \end{array} \right\}$	II. 485		2 2 51.4	+2.941	2	100 51 11.0	-17.20	3	vF; pS; R; 4th of 4	4
500	203			2 2 55.0	2.971	1	98 25 7.3	17.19	1	vF; vS; R; psbM	1
501	204	III. 604		2 3 56.7	3.593	1	53 10 55.5	17.15	1	vF; iF; stellar	2*
502	2467			2 4 15.8	1.992	1	147 23 34.1	17.13	1	pF; pS; R; glbM; r	1
503		III. 259		2 4 27.5	3.047	1	92 9 47.4	17.12	1	eF; eS; iF	1
504		II. 486		2 4 45.7	2.951	1	99 58 5.7	17.11	1	F; S; E	1
505	2468			2 5 36.7	2.555	3	126 30 33.9	17.07	4	eF; pS; IE 0°; gbM	4
506		II. 613		2 5 51.5	3.435	1	62 46 51.2	17.06	1	F; S; IE 90°; bM	1
507	2469			2 6 30.8	2.623	2	122 36 11.1	17.03	2	eB; S; E; psmbM	2
508	2470			2 7 24.0	2.420	2	132 41 12.3	16.99	2	F; vS; svmbM	2
509	205	III. 260		2 7 24.4	3.056	1	91 24 41.3	16.99	1	vF; R; bM; stellar	2
510	206	III. 457		2 8 6.4	3.140	1	84 39 32.5	16.95	1	eF; cL; R; gbM; *12 sf att.	2*
511		III. 2		2 8 41.3	3.078	1	89 36 2.1	16.93	1	eF; vS; R; bM	1
512	207	VI. 33		2 9 15.3	4.166	2	33 29 55.0	16.90	2	l; Cl; vL; vRi; st7...14 ...	5
513	208	III. 201		2 9 33.1	3.252	1	76 5 52.3	16.89	1	vF; vS; E; *10 sf 4'	2
514	208, a		R. nova	2 9		...	76 5+		...	neb s of h. 208	0
515	$\left\{ \begin{array}{l} 209 \\ = \\ 2471 \end{array} \right\}$	II. 474		2 9 43.2	2.920	5	101 59 58.6	16.88	5	F; pL; R; vglbM	7
516	210	II. 246		2 10 22.6	3.253	1	76 6 11.5	16.85	1	pF; pL; IE; pgbM; *9, 185° ± 5' S* sf 1'	2*
517	210, a		R. nova	2 10		...	76 6		...	neb s of h. 210	0
518	211	II. 436		2 11 14.5	2.980	1	97 17 24.7	16.81	1	F; pS; E; bM; 2 or 3 st nr...	2
519	213			2 11 59.3	3.272	1	74 48 57.9	16.77	1	eF; R; gbM; *16 nr	1
520	215	II. 437		2 12 8.7	2.978	2	97 25 44.2	16.75	2	pF; pS; vLE; bM; * nr	3
521	212	VI. 34		2 12 34.2	4.188	2	33 32 49.5	16.75	2	l; Cl; vL; vRi; ruby * M ...	5
522	214			2 12 46.6	4.536	1	26 52 20.1	16.73	1	Cl; L; lC; sc st 9...13	1
523	216	III. 486		2 12 54.3	2.852	1	106 42 4.1	16.73	1	F; S; iR; pgbM	2
524	2473			2 13 20.1	1.774	1	150 30 8.7	16.71	1	eF; S; R; 2 or 3 vSst nr ...	1
525	2472			2 13 30.8	2.400	2	132 23 1.0	16.70	2	vF; vS; R; bM; *7 sf and 6 more.	2
526	217	II. 225		2 13 41.2	3.548	4	57 22 48.3	16.69	4	B; S; R; bM; 3S st sp	5
527	218	V. 19		2 13 50.4	3.737	1	48 16 47.6	16.68	1	l; B; vL; vmE 22° 3	5†
528	2474			2 14 21.4	2.404	2	132 2 58.2	16.66	2	pF; pS; R; lbM; *8 90°, 4'	2
529	219	II. 438		2 14 35.7	2.993	1	96 10 8.8	16.64	2	F; vL; iR; gbM	4
530	219, a		R. nova	2 14		...	96 10		...	E; F; bM; makes D neb with h. 219; both E.	0
531		III. 695		2 15 1.0	4.448	1	28 40 47.1	16.63	1	eF; pL; iF	1
532	2475			2 15 9.4	2.563	1	124 21 30.4	16.62	1	pB; S; R; psbM; *10 f 90° 0 35''	1
533		III. 570		2 15 14.0	3.732	2	48 42 17.4	16.62	1	eF; vS; IE	1
534	2476			2 15 26.4	2.779	1	111 27 12.0	16.60	1	pB; S; gbM; r; *p	1
535	2477	III. 224		2 16 34.0	2.778	1	111 20 33.5	16.55	1	F; S; E 90°; gbM	3
536		I. 153		2 16 36.1	2.770	1	111 52 23.5	16.55	1	eB; vL; E 0°...90°	1*
537		III. 571		2 16 44.2	3.736	1	48 49 21.5	16.55	1	eF; stellar	1
538	221			2 17 49.4	3.319	1	72 7 46.3	16.49	1	pF; L; R; *10 sf 3'	1
539	220			2 17 50.6	3.543	1	58 23 33.3	16.49	1	vF; S; R; 4 st nr	1
540	2478	III. 239		2 18 44.7	2.710	1	115 26 13.5	16.45	1	cF; pL; R; gpmbM	2
541		III. 474		2 18 50.8	3.350	1	70 7 27.8	16.44	1	eF; vS; iR	1
542	222	III. 177		2 18 55.7	3.571	2	57 3 25.1	16.43	2	cF; cL; E; vgbM; 2st13np..	3
543		II. 489		2 19 58.5	3.350	1	70 15 30.6	16.38	1	F; S; IE; 3 st inv	1
544	223	IV. 23		2 20 29.1	3.049	1	91 46 18.5	16.35	1	vB; vL; R; mbMN	2†
545	2479			2 20 59.9	2.295	2	135 4 17.1	16.33	2	vvF; S; R; gvlbM	2
546	224	III. 261		2 21 21.5	3.049	1	91 47 36.7	16.31	1	vF; cL; R; f of 2	2
547	$\left\{ \begin{array}{l} 225 \\ = \\ 2480 \end{array} \right\}$	II. 487		2 21 48.4	2.919	2	101 10 10.3	16.29	2	vF; L; iR; glbM	3
548	2481			2 22 1.5	+2.795	1	109 40 3.6	-16.28	1	pB; E; gbM	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
549	226	I. 154		2 22 18.6	+3.654	1	53 29 51.2	-16.26	1	eB; L; E; vgbM	3*
550	2482			2 23 20.3	2.366	4	132 1 30.7	16.21	4	vF; pL; lE; gbM; *8 sf 3'...	4
551	229	II. 278		2 23 24.6	3.050	1	91 43 1.7	16.21	1	pB; S; E; psbM	3
552	228			2 23 27.5	3.837	1	45 59 37.0	16.20	1	Cl; pRi; st 9...15	1
553	227			2 23 28.9	4.288	1	33 5 46.0	16.20	1	Cl; pL; pRi; st 13...15	1
554	230	II. 237?		2 23 38.8	3.024	1	93 33 56.3	16.19	1	pF; iE 0°±; bM	2
555	2483			2 25 21.5	2.482	2	126 39 0.0	16.10	4	pB; pS; mE 215°7	4
556	2484			2 25 38.1	2.817	2	107 49 36.8	16.09	2	F; S; iR; gbM	2
557	231			2 25 44.2	3.580	1	57 40 44.3	16.09	1	S; R; psbM; 1st of 3	1*
558	231, a		R. nova	2 25 47.7	3.580	...	57 38 59.3	16.09	...	} γ and δ of Lord R.'s diag.	{ 0*
559	231, b		R. nova	2 25 52.0	3.580	...	57 38 50.3	16.09	...		
560	232	II. 211		2 25 52.4	3.512	(2)	61 17 57.6	16.08	(2)	pB; eL; lE 0...90°; gmbM; 3st s.	3†
561	233			2 26 0.7	3.580	1	57 40 14.9	16.07	1	vF; R; bM; 2nd of 3	1*
562	2485	III. 472		2 26 16.8	2.912	1	101 22 37.5	16.05	1	eF; pS; R; vlbM; amscst...	2
563	234			2 26 21.2	3.580	1	57 46 23.5	16.05	1	pB; R; 3rd of 3	1*
564	2486			2 26 26.9	2.270	2	135 8 21.8	16.04	2	F; S; R; bet 2 st in par	2
565	235	III. 572		2 26 33.1	3.753	1:	49 47 23.8	16.04	1:	vF; pS; p of 2; 210°; 157°...	2
566	236	III. 573		2 26 37.4	3.754	1	49 44 28.1	16.03	1	F; S; f of 2; 210°; 337°...	2
567	2487		Δ. 519??	2 28 0.1	2.404	2	129 39 17.2	15.96	2	pB; L; pmE; smbM; bi-N...	2†
568	237	III. 161		2 28 23.1	3.595	2	57 17 16.5	15.95	2	F; S; vLE; bM; r; 2 st 14 np	4
569	238	III. 557		2 28 43.8	3.232	1	78 58 34.1	15.93	1	F; S; vLE; psbM; r	2
570	239	III. 434		2 28 46.2	2.963	1	97 46 30.4	15.92	1	vF; eL; iF; vlbM	2
571	240	II. 238	}	2 30 24.9	3.761	1	49 43 53.8	15.84	1	pF; L; E 90°±; mbM; r ...	4*
		=									
		III. 198									
572	241	III. 152		2 30 56.6	3.540	1:	60 27 46.0	15.80	1	F; pS; iR; bM; st inv	3†
573		II. 6		2 31 35.3	3.077	1	89 44 6.9	15.77	1	S; cometic	1*
574	244	I. 102		2 31 36.0	2.968	3	97 17 18.9	15.77	3	eB; pL; vR; mbM	5*
575	242	I. 156		2 31 38.3	3.731	1	51 32 45.9	15.77	1	vB; vL; vmE; vmbM	3†
576	243	II. 592		2 31 38.5	3.222	1	79 45 46.9	15.77	1	pF; S; lE; bM; *11, 25° 50°	2
577	2488			2 31 43.4	1.877	1	145 28 45.9	15.77	1	eF; S; R; p of 2	1
578		VIII. 66		2 32 1.1	4.572	2	29 3 23.5	15.75	2	Cl; L; sc st, one 10	2
579	245	III. 581		2 32 2.2	3.333	1:	72 34 8.5	15.75	1:	vF; iE	2
580	2490			2 32 8.3	1.874	1	145 28 16.8	15.74	1	F; S; R; gbM; *11, s 2' ...	1
581	246	II. 5		2 32 11.6	3.080	2	89 30 24.8	15.74	2	pB; S; vLE 0...90°; bm; 3st trap.	10
582	{ 249 = 2489 }	II. 284		2 32 36.5	2.945	2	98 44 27.7	15.71	2	pF; L; mE; r; *17, att sf ...	4
583	247	III. 475		2 32 42.1	3.354	(1)	71 19 38.7	15.71	1	F; S; R; lbM	2
584	248		M. 34	2 33 2.3	3.829	2	47 49 25.0	15.70	2	Cl; B; vL; lC; sc st 9	8
585	251	III. 228		2 33 36.3	3.193	1	81 52 7.2	15.66	1	vF; vS; p of 2; *10 p	2
586	{ 253 = 2491 }	II. 488		2 33 42.9	2.898	2	101 53 28.2	15.66	2	F; S; R; bM	3
587	252	III. 229		2 33 43.8	3.192	1	81 53 22.2	15.66	1	eF; vS; f of 2	2
588	2492			2 33 57.9	2.491	1	124 52 20.8	15.64	1	pB; S; R; stellar	1
589	{ 254 = 2493 }	I. 63		2 34 12.1	2.943	3	98 51 19.1	15.63	3	B; pL; R; mbM *12	4
590	256	III. 584		2 34 35.0	3.520	1:	62 1 17.0	15.60	1	F; S; R; psbM	2
591	258	I. 1		2 34 35.1	3.072	2	90 9 14.0	15.60	2	pF; eL; iE 80°; bM; pB*nr	8*
592	255	II. 633		2 34 39.9	3.701	1	53 16 21.0	15.60	1	pF; eL; R; glbM	3
593	259			2 34 44.5	3.337	1	72 35 27.0	15.60	1	eF; ?	1
594	257	III. 162		2 34 49.1	3.596	2	58 10 18.3	15.59	2	F; pL; R; lbM; *7.8 p 43°5	4
595	} 257, a		R. 3 novæ	2 34		...	58 10		...	{ 6 seen (including .- h.) 257, 260, 261). }	0
596											
597											
598	260	III. 163		2 35 23.6	3.598	1	58 7 34.9	15.57	1	vF; pL; R; lbM; sp of 2 ...	2
599	261			2 35 26.2	+3.599	1	58 4 35.2	-15.56	1	eF; S; nf of 2	1

No. of Cata- logues.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. II.'s Catalogues of Nebulæ.	Sir W. II.'s Classes and Nos.	Other Authorities.								
600	h. 262	H.	M. 77	h m s 2 35 30.2	s +3.065	1	90° 35' 56.2	-15.56	1	vB; pL; iR; sbMrrN; *130°, 2'	13†
601	263	II. 273		2 36 0.3	3.138	1	85 37 19.1	15.53	1	pF; S; iR; gbM	2
602		III. 455		2 36 33.3	3.085	2	89 13 22.0	15.50	2	vF; L; lbM; er	2
603	2494			2 37 43.7	2.589	1	119 35 49.8	15.44	1	B; pL; pmE; sbM	1
604	264	I. 64		2 39 6.9	2.950	1	98 9 52.5	15.35	1	vB; pL; E; gpmbM	3
605	265	II. 466		2 39 15.4	3.057	1	91 5 53.8	15.34	1	pB; cL; iE; mbM	4
606	266	II. 465		2 39 21.1	3.061	1	90 49 5.8	15.34	1	vF; pL; iR; bM	4
607		III. 582		2 39 22.7	3.313	1	74 25 31.8	15.34	1	vF; S; iF	1
608	267	III. 462		2 40 18.4	3.061	1	90 50 57.3	15.29	1	vF; S; R; 2Sst p	2
609	2496			2 40 20.8	1.552	1	150 30 14.3	15.29	1	F; pS; R; glbM	1
610	2495	V. 48		2 40 21.5	2.557	2	120 51 30.3	15.29	2	vB; L; vmE 151°1; vbMN	3
611	{ 269 = 2497	III. 449		2 42 35.2	2.796	2	107 34 23.5	15.15	2	pF; pL; pmE; glbM	3
612	268			2 42 39.0	3.839	1	48 55 4.5	15.15	1	neb or vScl of vSst	1
613	270	II. 601		2 43 44.5	3.858	1	48 22 21.3	15.09	1	eF; S; iR; vgbM; r	2
614			Bessel	2 44 6.5	3.738	...	53 4 14.9	15.07	...	? a comet	0*
615	272	III. 450		2 45 8.9	2.799	1	107 12 50.7	15.01	1	vF; S; iE; gbM	3
616	271	II. 602		2 45 25.7	3.847	1	49 0 2.3	14.99	1	eF; pS; iR; vglbM	2
617	{ 271, a		R. 2 novæ	2 45		...	49		...	h. 271 is D; another near ...	0
618											
619	273			2 45 44.4	3.044	1	91 51 24.6	14.98	1	eF; pL; gbM; *8f	1
620		II. 254		2 46 0.9	3.269	1	77 34 54.2	14.96	1	F; S; iR; r	1
621	2498			2 46 35.8	1.778	1	145 32 30.4	14.92	1	F; R; gbM	1
622	2499			2 46 44.5	1.772	1	145 38 31.4	14.92	1	F; R; gbM	1
623	274	III. 580		2 47 29.7	3.897	1	47 31 21.6	14.88	1	vF; vS; R; gbM; 2Sst Δ	2
624	{ 275 = 2500	II. 470		2 47 46.2	2.905	2	100 36 9.2	14.86	2	vF; S; R; stellar	4
625	2501			2 48 7.7	2.760	1	109 13 6.8	14.84	1	F; pL; vmE; 2Sst f	1
626	276	II. 274		2 50 52.8	3.118	1	87 11 7.6	14.68	1	F; vS; iE; sbM; er	2
627		II. 619		2 51 35.0	3.493	1	65 21 16.1	14.63	1	pB; cL; pmE 0°; r; *n 1'	1
628	277	III. 199		2 51 54.7	3.949	1	45 40 24.7	14.61	1	F; pS; iE; SbM; *p 6°5	4
629	277, a		R. nova	2 52 +		...	45 40 ±		...	R (nisi=H. II. 239)	0
630	2502	III. 469		2 52 16.6	2.863	1	102 57 33.3	14.59	1	F; R; glbM; stellar	2
631		III. 178		2 52 47.8	3.711	1	55 18 19.2	14.56	1	vF; pL; R; spmbM	1
632	278			2 52 51.3	3.908	1	47 58 42.2	14.56	1	eF; vS	1
633	2503			2 52 56.7	2.484	2	122 39 22.5	14.55	2	vF; pL; E; vlbM	2
634		II. 239?		2 53 0.1	3.977	2	45 35 20.5	14.55	2	pB; pL; iR; nbM	3
635	279	II. 620		2 53 28.9	4.025	(2)	44 10 52.4	14.52	(2)	pF; pS; iF; sbM	2
636	280	II. 502		2 55 0.2	2.818	1	105 23 8.1	14.43	1	pF; pL; R; psbM	2*
637		II. 607		2 55 21.2	3.906	1	48 13 26.7	14.41	1	F; cL; E	1
638		II. 704		2 55 44.9	8.608	1	9 42 40.3	14.39	1	F; pL; mE 90°...180°	2
639	281	IV. 43		2 56 20.1	3.925	1	47 43 32.5	14.35	1	F; mE; smbMS*	3*
640	2504	III. 245		2 56 25.0	2.669	1	113 25 4.8	14.34	1	pF; cL; pmE; gbM*16; r...	2
641		II. 608		2 56 32.2	3.975	1	46 10 29.0	14.40	1	F; cL; er	1
642	2505			2 56 55.1	2.864	1	102 38 19.0	14.30	1	vF; sp of 2	2
643	282	II. 503		2 57 5.8	2.802	1	106 8 59.3	14.29	1	eB; pS; iR; smbM	2
644	2506	II. 475		2 57 14.1	2.865	1	102 33 6.3	14.29	1	pF; cL; iR; bM; nf of 2	3
645	283	I. 109		2 58 3.7	2.603	1	116 35 45.8	14.24	1	eB; pS; vIE 0°; r; S*nr	4
646	284	III. 578		2 59 19.7	3.810	1	52 9 24.2	14.16	1	eF; vS; R; psbM	2*
647	{ 285 = 2507	II. 285		2 59 25.2	2.905	3	100 5 8.5	14.15	3	pB; S; iE 80°±; lbM	7
648	286	II. 504		2 59 30.2	2.799	1	106 8 39.5	14.15	1	B; S; eE; psbM	2
649	2508			3 0 48.0	2.288	1	129 34 26.9	14.07	1	pF; S; R; psbM	2
650	287			3 1 29.2	4.337	1	37 11 56.1	14.03	1	Cl; vS of Sst	1
651	2509	II. 258		3 3 27.9	2.702	3	111 7 0.0	13.90	3	pB; cL; R; gbM; r	7
652	288	III. 262		3 4 20.1	3.022	(1)	93 3 4.5	13.85	1:	stellar; difficult	2
653		III. 164		3 4 23.3	+3.634	1	59 57 41.8	-13.84	1	eF; vS; ? vSst	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	"			
654	$\left\{ \begin{array}{l} 289 \\ = \\ 2510 \end{array} \right\}$	II. 286		3 4 26.8	+2.912	3	99 27 23.8	-13.84	3	F; pL; R; vglbM; *9np ...	5
655	$\left\{ \begin{array}{l} 289, a \\ 291 \\ = \\ 2511 \end{array} \right\}$		R. nova	3 4 31.4	2.912	...	99 26 25.8	13.84	...	No description.....	0*
656	$\left\{ \begin{array}{l} 291 \\ = \\ 2511 \end{array} \right\}$	III. 591		3 4 36.2	2.912	2	99 29 11.6	13.78	2	eF; vS; R; stell; sf of 2.....	3*
657	2512		Δ. 205??	3 4 41.9	0.771	2	157 18 47.1	13.83	2	F; S; pmE; gbM	3
658	290	VI. 25		3 5 6.3	4.106	1	43 17 22.3	13.79	1	Cl; pL; Ri; C; iR; st 12...15	3
659	2513			3 5 18.2	0.747	2	157 29 13.6	13.78	2	pF; S; R; glbM	2
660		II. 900		3 5 38.6	2.884	1	101 0 7.2	13.76	1	F; pL; E 80°+	1
661	292	III. 443		3 5 49.8	2.975	3	95 45 4.5	13.75	3	cF; S; iE; bM; *9, n 5' ...	4
662	2514			3 5 57.7	1.775	1	143 52 15.8	13.74	1	B; L; vmE 80°; vgbM	2
5060				3 6 55.2		...	89 4 27.5		...	See No. 5060.	
663	2515			3 7 7.5	1.474	1	148 40 18.9	13.67	1	Cl of 18 or 20 st.....	1
664		IV. 17		3 7 7.6	3.016	1	93 27 10.9	13.67	1	* with neb att 90° 1	1
665	2516			3 7 46.9	2.667	1	112 30 46.1	13.63	1	F; S; E; alm stell; *8, np...	1
666	2517		Δ. 337	3 8 25.7	1.635	2	145 44 50.6	13.58	2	⊕; B; L; R; rr	2
667		III. 194		3 8 43.9	3.020	1	93 8 18.9	13.57	1	eF; eS	1
668			D'Arrest, 35	3 9 31	3.93	[2]	49 2 30	13.49	[2]	F; vS; R; stellar; 1st of 7...	0
669			D'Arrest, 36	3 9 31	3.93	[2]	49 1 42	13.49	[2]	eF; S; iE; cometary; 2d of 7	0
670	2518			3 9 41.5	2.198	1	131 36 25.7	13.51	1	vB; R; gmbM	1
671			D'Arrest, 37	3 9 45	3.93	[2]	49 1 54	13.48	[2]	vF; S; R; 3rd of 7.....	0
672			D'Arrest, 38	3 10 8	3.93	[2]	49 1 36	13.46	[2]	F; S; R; 4th of 7	0
673			D'Arrest, 39	3 10 11	3.94	[2]	48 58 30	13.45	[2]	vF; vS; 5th of 7	0
674	293	II. 603		3 10 32.7	3.937	1	49 0 5.5	13.45	1	pB; pS; R; bM; 6th of 7 ...	2*
675			D'Arrest, 40	3 10 35	3.97	[2]	49 0 3	13.42	[2]	F; S; *17; 7th of 7	0
676	2519	III. 956		3 11 2.0	2.884	1	100 48 41.4	13.42	1	eF; vS; 2 st 2' or 3' s.....	2
677	293, a		R. 6 novæ	3 11 ±		...	49 ±		...	6 of 15 (including probably h. 294, 295)	0
678											
679											
680											
681											
682											
683	2520			3 11 34.2	2.425	1	123 5 43.3	13.39	1	vF; L; R; vglbM	1
684		III. 195		3 11 43.7	3.017	1	93 14 30.6	13.38	1	eeF; eS	1*
685	2521		Δ. 487	3 12 15.0	2.189	2	131 36 47.1	13.33	2	⊕; vB; pL; R; mbM; er ...	2
686	294	III. 574		3 12 23.9	3.938	1	49 11 42.1	13.33	1	vF; R; bM; p & sm of 2 ...	2
687	295	III. 575		3 12 25.0	3.938	1	49 10 4.4	13.32	1	vF; R; bM; f of 2; 100'', 352°-4.	2
688	296	II. 287		3 13 15.0	2.954	4	96 45 57.9	13.27	4	vF; S; vIE; gbM; er	7
689	2522			3 13 22.5	2.710	2	110 55 8.9	13.27	2	cB; vL; vmE; psvmbM.....	2
690		III. 444		3 14 7.5	2.983	1	95 8 11.7	13.21	1	eF; vS	1
691		III. 568		3 15 25.7	3.016	1	93 15 44.1	13.13	1	eF; S; iF; am 3 or 4 st	1
692	2523	I. 106		3 15 36.2	2.786	1	105 53 51.7	13.11	1	pB; cL; iR; gbM; *7, f 7°-5, 211°-0.	3
693	2524			3 15 41.5	2.295	3	127 38 38.7	13.11	3	⊕; vF; pL; R; vglbM.....	3
694	2525			3 16 0.3	1.749	1	142 41 0.3	13.09	1	F; pL; mE 37°-3; gbM	1
5061				3 16 29.0		...	89 19 3.0		...	See No. 5061.	
695	2528		Δ. 206	3 16 34.1	0.698	1	156 59 37.2	13.06	1	pB; L; iE; vgbM; r	1
696	2526			3 16 54.8	2.666	1	111 52 19.1	13.03	1	pB; S; R; gbM	1
697	2527		Δ. 548	3 17 20.4	2.288	2	127 43 40.0	13.00	2	vB; cL; vIE; vsvmbMN ...	2
698	2529		Δ. 547	3 17 23.1	2.291	2	127 36 24.0	13.00	2	pB; pS; psbM.....	2
699	2533			3 17 45.6	2.662	1	112 1 0.9	12.97	1	F; S; R; bM; p of 2	1
700	$\left\{ \begin{array}{l} 298 \\ = \\ 2530 \\ 297 \\ = \\ 2531 \end{array} \right\}$	III. 197		3 17 48.3	3.011	1	93 31 34.9	12.97	1	vF; S; R; bM; 1st of 3.....	3
701	$\left\{ \begin{array}{l} 297 \\ = \\ 2531 \end{array} \right\}$	III. 196		3 17 48.8	3.011	1	93 30 34.9	12.97	1	vF; vS; E; ? neb *; 2nd of 3	3
702	2532			3 17 50.9	+3.012	1	93 25 54.9	-12.97	1	F; vS; R; bM; 3rd of 3.....	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. II.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	"			
703	297, a		R. nova	3 17 53.6	+ 3.011	...	93 16 41.9	-12.97	...		0
704	299	III. 445		3 18 4.9	2.962	2	96 14 11.2	12.96	2	vF; pS; pmE	3
705	2534	IV. 77		3 18 18.0	2.661	2	112 1 49.8	12.94	2	F; mE; 239° 1; com; *9, 10 att.	3†
706	2535			3 18 37.6	2.305	1	126 58 12.7	12.91	1	O? pS; vsymbMN	1
707	2536			3 19 39.5	2.739	1	108 4 59.5	12.85	1	F; pS; R; glbM	1
708		III. 959		3 19 53.5	2.662	2	111 51 1.8	12.84	2	vF; vS; sf of 2	1*
709		I. 60		3 20 15.0	2.662	2	111 50 36.0	12.80	2	vB; S; E90°...180°; smbMN; np of 2.	2*
710			Anw. N. 17	3 20 41.7	3.690	...	59 6 31.2	12.76	...	F; L; *10f 4'; n 2' 5 (Schönfeld, 1858).	0*
711			D'Arrest, 41	3 20 47	3.98	[1]	48 39 6	12.72	[1]	eF; pL; lbM	0
712	2537			3 21 10.0	2.319	3	126 12 33.5	12.75	3	vF; S; vLE; gbM	3
713	2538			3 22 29.8	2.407	2	122 46 15.5	12.65	2	cB; pS; R; psbM; *p	2
714	2542	I. 257		3 22 38.1	2.437	(1)	121 34 40.8	12.64	1:	cB; pL; iR; vgbM	2
715	2539			3 22 38.5	2.441	1	121 23 12.8	12.64	1	vB; pS; lE; psbM	1
716	2540			3 22 39.6	2.275	1	127 38 11.8	12.64	1	F; S; R; *12, sf	1
717	301	VIII. 88		3 22 41.0	3.852	1	53 9 36.8	12.64	1	Cl; vL; ab 60 st	3
718	300	III. 694		3 22 54.4	6.312	2	17 54 18.1	12.63	2	F; vS; iR; gbM; *vnr	3
719	2541			3 23 10.3	2.732	1	108 16 10.7	12.61	1	vF; S; R; psbM	1
720		VIII. 84		3 23 42.3	4.361	1	39 3 16.6	12.58	1	Cl; lRi; stL	1
721	2545		Δ. 591	3 24 11.0	2.365	1?	124 12 43.8	12.54	1:	B; L; mE; vmbMRN	1
722	2544			3 25 8.3	2.332	1	125 20 9.9	12.47	1	pB; pS; R; psbM	1
723	2543			3 25 16.8	2.698	1	109 45 38.2	12.46	1	eF; psbM; v diff *8, sf	1
724	2546	III. 246		3 25 52.1	2.665	2	111 17 46.1	12.43	2	pB; cL; iE; mbM	5
725	2547	III. 487		3 26 2.0	2.778	1	105 41 32.4	12.42	1	vF; S; lE; glbM	2
726	2548	II. 290		3 26 44.0	2.808	2	104 9 24.9	12.37	2	pF; pL; R; lbM; pL*nf 5'	5
727	302	III. 446		3 26 44.3	2.971	2	95 33 27.9	12.37	2	vF; S; bet 2st	3
728	2549			3 26 30.7	1.789	2	140 45 56.6	12.38	2	vF; pL; iR; gbM; *nr	2
729	2550			3 27 28.6	2.690	1	109 58 38.4	12.32	1	F; L; R; vglbM	1
730	2551	III. 960		3 27 38.5	2.674	2	110 46 3.7	12.31	2	vF; S; R	3
731	2552			3 28 18.0	2.289	2	126 36 31.9	12.27	2	!! vB; vL; mE; rN in vLE Halo.	2†
732	2553	III. 857		3 28 18.2	2.421	(1)	121 40 37.9	12.27	(1)	vF; S; iF; lbM	2
733	2554	III. 559		3 29 0.5	2.670	3	110 50 45.4	12.22	3	vF; S; R; bet 2st 14	4
734	2555	II. 262		3 29 1.6	2.570	1	115 24 21.4	12.22	1	pB; pL; vLE; psbM	2
735	2556			3 29 50.6	2.310	1	125 42 24.2	12.16	1	eF; vS; p of 3	1
736	2557			3 30 0.6	2.311	1	125 41 4.5	12.15	1	vB; pL; lE; gmbM; 2nd of 3	1
737	2558			3 30 0.6	2.309	1	125 43 44.5	12.15	1	B; S; lE; pmbM; 3rd of 3	1
738	303	II. 288		3 30 7.0	2.971	2	95 30 0.1	12.13	2	eF; pL; iR; bM; r	5
739	2559		Δ. 574	3 30 9.5	2.316	1?	125 28 56.8	12.14	1:	vB; L; R; psbM	1
740	2560	III. 961		3 30 28.7	2.657	2	111 21 44.4	12.12	2	F; S; R; gbM	3
741	2561			3 30 42.7	2.307	1	125 55 1.7	12.11	1	⊕; B; pL; R; gpmbM	2
742	2562			3 31 19.8	2.710	1	108 48 18.5	12.05	1	pF; S; R; psmbM	1
743	2563	II. 263		3 31 26.9	2.576	1	114 58 4.5	12.05	1	pB; pS; R; gpmbM	2
744	2564			3 31 34.7	2.298	1	125 58 33.5	12.05	1	⊕; vB; pL; R; gmbM	2
745	2565	III. 451		3 32 19.8	2.706	1	108 53 34.6	11.98	1	F; S; R; glbM	2
746	2566	I. 58		3 32 24.1	2.608	2	113 28 55.6	11.98	2	B; pS; E; psmbM	4
747	2567	II. 593		3 33 11.0	2.700	1	109 9 29.4	11.92	1	eB; pS; R; psmbM	2
748	2569			3 33 11.6	2.296	2	125 54 53.1	11.93	2	⊕; vB; pL; psbM; rr	3
749	2568	III. 247		3 33 14.1	2.613	1	113 10 47.4	11.92	1	vF; vS; R	2
750	2571			3 33 43.1	2.291	1	126 2 48.0	11.90	1	vB; pL; R; psmbM	2
751	2572			3 33 49.8	2.407	1	121 46 25.6	11.88	1	F; cL; vME; vglbM; *7np	1
752	2570	I. 107		3 33 55.4	2.702	1	109 2 21.9	11.87	1	vB; L; R; svmbMN	3
753	304	III. 263		3 34 4.2	3.040	(2)	91 45 34.5	11.85	(2)	eF; stellar or lE	2
754	304, a		R. nova	3 34		...	91 45		...	makes D neb with h. 304	0
755	2573			3 34 5.2	2.014	2	134 33 18.9	11.87	2	B; pS; R; smbM	2
756	305	III. 569		3 34 32.3	2.977	1	95 7 5.4	11.82	1:	eF; lE; er; 1st of 3	2
757	2574			3 34 36.5	2.531	1	116 40 8.4	11.82	1	F; S; E; gbM; *sf 2'	1
758	306	II. 455		3 34 41.5	2.976	1	95 8 51.7	11.81	1	pF; pL; lE; lbM; *sf; 2d of 3	3
759	2575	II. 267		3 34 51.1	+ 2.615	3	113 0 43.7	-11.81	3	pB; S; lE; pglbM; *sf 2'	4

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	"			
760	307	II. 456		3 35 2.3	+ 2.976	3	95 9 45.6	-11.78	2	vF; S; E; B* 135°, 1'; 3d of 3	4
761	2576			3 35 29.9	2.229	2	127 58 34.9	11.77	2	pF; pS; R; psbM	2
762	2577	II. 291		3 35 56.9	2.804	2	103 56 56.1	11.73	2	F; eL; mE 0° ±; r.....	4
763	307, a		R. nova	3 35 58.6	2.976	::	95 9 45.6	11.73	::	No description	0
764		II. 852		3 36 30.1	2.438	1	120 21 11.3	11.69	1	F; pL; iR; gbM	1
765	2578	III. 248		3 36 42.1	2.622	2	112 33 21.6	11.68	2	pF; S; lE; bM	3
766	2579			3 37 0.0	2.288	1	125 50 50.2	11.66	1	pF; S; R; psmbM	1
767	2580		Δ. 426	3 37 37.8	1.879	2	137 40 33.4	11.62	2	vB; L; pmE; vsymbM*10...	2
768			Auw. N. 18	3 37 52.3	3.542	...	66 40 12.9	11.57	...	!!!B; vL; iF; VAR. (Tempel)	0*
769	2581		Δ. 562	3 38 7.9	2.264	1?	126 34 12.6	11.58	1?	⊕; vB; pmE; pgbM	2
770	2582			3 38 20.0	2.272	1	126 18 1.9	11.57	1	F; vL; R; glbM	1
771	2584	III. 249		3 38 43.4	2.623	3	112 21 40.1	11.53	3	F; pS; gpmbM	5
772		II. 597		3 38 44.4	2.936	2	94 31 51.1	11.53	2	vF; S; iE; *nr	1
773	2583	II. 458		3 38 44.4	2.703	2	108 43 32.1	11.53	2	pB; pS; R; smbM*13	3
774		II. 594		3 38 46.3	2.682	2	109 41 52.1	11.53	2	pB; vS; bM	1*
775	308	VIII. 80		3 38 52.0	4.490	2	37 46 18.6	11.48	2	Cl of ab 30st 12...14	3
776	2585			3 39 4.9	1.977	1	135 5 20.4	11.52	1	pB; L; vmE 221° 6	1
777		II. 459		3 39 12.4	2.696	1	109 0 25.3	11.49	1	F; R; lbM	1
778	309	I. 155		3 39 32.4	2.989	1	94 24 37.9	11.47	1	pB; S; R; *17M	3*
779	2586			3 39 56.6	1.974	3	135 5 22.2	11.46	3	pF; pL; eE 42° 3; vgpmbM	3
780	2587			3 40 59.9	2.241	1	127 7 51.9	11.37	1	F; S; R; *att	1
781	2588	II. 460		3 42 24.8	2.739	1	106 49 22.2	11.26	1	pB; S; lE; mbMN	2
782	2589			3 43 32.5	+1.139	2	150 14 27.4	11.22	2	eF; S; R; glbM; am 7Bst	2
783	2590			3 45 15.1	-0.360	1	162 6 36.1	11.13	1	pF; pS; iR; glbM; *7f	2
784	2592			3 46 48.5	+0.224	2	158 38 24.7	11.01	2	eF; pL; R; gvlbM	2
785	2591			3 47 34.5	1.955	2	134 56 47.0	10.90	2	eF; S; E 90°; gbM	2
786	2593			3 48 18.8	2.644	1	110 52 29.8	10.84	1	eF; S; R; 2Bstf; p of 2	1
787	2594	III. 962		3 48 28.0	2.643	3	110 54 59.4	10.82	2	F; S; vLE; 2st 10nr; f of 2	4
788	2595		Δ. 427? 428?	3 48 31.1	1.830	2	137 53 54.8	10.84	2	eF; pL; R; vglbM	2
789	2596			3 49 9.1	2.212	1	127 24 18.6	10.78	1	vF; L; E; vglbM	1
790	2597		Δ. 480	3 51 4.2	2.029	3	132 46 40.8	10.64	3	pb; pL; R; gbM; 2st Δ	3
791			Auw. N. 19	3 52 1.3	3.444	...	71 49 55.8	10.54	...	*12 inv in neb (Markree Cat. Nov. 24, 1854).	0
792	2599			3 52 22.9	0.478	2	156 25 42.3	10.59	2	pB; S; vLE; pmbM	2
793		I. 258		3 52 32.8	4.482	1	39 1 19.4	10.52	1	vB; S; iF; bM; r; *inv	1
794	2598			3 52 59.7	2.251	1	125 51 46.3	10.49	1	vF; vS; R	1
795	2600		Δ. 438	3 53 5.8	1.870	2	136 36 53.0	10.50	2	F; eL; R; vglbM	2
796	2601			3 53 36.3	1.748	2	139 18 42.9	10.47	2	F; L; R; vglbM; 3st n	2
797	2602			3 53 47.2	1.721	3	134 52 57.5	10.45	3	eF; S; lE 90°; vglbM	3
798	310			3 53 47.4	4.551	1	37 45 45.6	10.38	1	Cl; segment of a ring	1
799		VII. 3		3 53 52.2	2.821	1	102 25 27.4	10.42	1	Cl; S; C	1
800	2603		Δ. 369?	3 54 33.6	1.573	1	142 43 35.3	10.39	1	F; vS; R; pmbM; *8 np	1
801		IV. 53		3 54 57.0	5.109	2	29 27 30.6	10.28	2	O; pB; pS; vLE; 1' diam	2
802		VII. 47		3 55 9.9	5.232	2	28 3 51.2	10.26	2	Cl; pRi; eC; iF	2
803	2604			3 55 25.3	0.456	1	156 25 44.2	10.36	1	eF; pS; R; *10 np	1
804	2605			3 56 44.9	1.552	1	142 58 5.1	10.23	2	eeeF; S; R; bet 2st 12 & 13.	2
805		II. 279		3 57 19.5	3.021	2	92 35 1.5	10.15	2	vF; pL; mE; vlbM; er	2
806	2606			3 59 0.1	1.965	2	133 47 35.5	10.05	2	F; pL; R; vgmbM	2
807	2608			3 59 3.1	0.218	3	158 1 23.7	10.09	3	pB; pS; mE 121° 5; gbM	3
808	2607		Δ. 466	3 59 20.3	1.966	3	133 44 23.1	10.03	3	⊕; B; eL; R; bM; rr	3
809		VII. 60		3 59 35.9	4.416	1	40 51 49.5	9.95	1	Cl; L; vRi; pC; st vL	1
810	311	IV. 69		4 0 28.6	3.755	2	59 35 29.2	9.96	2	*8m in neb 3' diam	3*†
811	2609		Δ. 348	4 0 39.4	1.448	2	144 29 25.8	9.94	2	B; L; vmE 10°; bM	2
812	2610	III. 499		4 1 24.0	2.884	2	99 12 21.5	9.85	2	eeF; S; E; psmbM; er	3
813	2611			4 1 44.0	+2.613	1	111 33 0.1	9.83	1	B; L; pmE; gbM; *8 sp	1
814	2615			4 2 8.1	-2.026	1	167 13 0.4	9.92	1	Cl; pL; lRi; st 9...10	1
815	2612			4 2 10.3	+2.616	1	111 25 36.0	9.80	1	pB; R; bM	1
816	2613			4 2 35.2	1.525	2	143 2 40.3	9.79	2	eF; vS; R; vlbM	2
817	2614			4 2 48.1	1.440	1	144 28 35.9	9.77	1	vF; R	1
818	2617			4 4 8.8	0.425	2	156 12 37.0	9.70	2	eF; vS; R; glbM	2
819	2616			4 4 17.7	+1.761	2	138 15 52.5	- 9.65	2	pB; pS; E 77°; vsmbMRN...	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
820		VII. 61		4 4 37.1	+4.520	1	39° 7' 14.2	-9.56	1	Cl; B; vRi; cC	1
821	2619			4 5 33.4	0.746	1	153 16 3.3	9.59	1	vF; S; R; gbM	1
822	2620			4 6 34.6	2.302	3	123 12 36.2	9.46	3	pB; pL; R; bM; np of 2 ...	3+
823	2621		Δ. 600	4 6 39.6	2.301	3	123 14 8.2	9.46	3	B; vL; vmE 32° 2; psmBm ...	3+
824	2622			4 6 51.0	1.297	2	146 29 15.9	9.47	2	vB; vL; R; smbM; 2st *10nf	2
825	2623			4 6 59.8	0.749	2	153 9 21.9	9.47	2	F; S; R; vS* ² d sf	2
826	2618	IV. 26		4 7 50.8	2.792	A	103 5 32.2	9.36	1	⊕; vB; S; R; ps, vsbM; r...	5*+
827	2625			4 8 4.3	1.267	1	146 50 24.6	9.38	1	vF; R; pL; vlbM	1
828	2624			4 8 17.9	2.337	1	121 54 41.8	9.34	1	vB; pS; IE; psvmbM	1
829	2626			4 9 32.7	2.420	2	118 50 16.1	9.23	2	vF; vS; E; gvlbM; r	2
830	2627			4 9 58.6	1.170	1	148 5 33.1	9.23	1	B; pL; E; smbMN=*11 ...	1
831		VIII. 85		4 10 25.3	4.489	1	40 5 39.7	9.11	1	Cl; pRi; IC; stL	1
832	2628			4 11 37.5	1.282	2	146 24 57.7	9.11	2	pB; IE; gbMEN; *p	2
833	312			4 11 45.8	3.959	1	53 25 50.1	9.03	1	Cl; vL; lRi; IC; st10...12...	1
834	2629			4 12 9.4	1.313	1::	145 55 46.2	9.06	1	B; pS; R	1
835		D'Arrest, 42		4 12 21	3.11	[1]	87 56 12	8.99	[1]	vF; S; R; *13nr	0
836		II. 464		4 12 22.5	3.097	1	88 55 34.0	9.00	1	F; vS; R	1*
837	313	III. 490		4 13 10.4	3.052	2	91 2 17.1	8.93	2	eF; pS; IE; vgbM; *11sf ...	3
838	2630			4 13 11.1	1.295	1	146 7 50.6	8.98	2	vB; pS; R; gmbM; am 3st...	2
839			Anw. N. 20	4 13 47.7	3.488	...	70 48 46.0	8.87	...	!!!; vF; S; variable (Hind)...	0*
840	2631			4 13 56.5	+1.622	2	140 30 9.7	8.91	2	cF; S; R; vglbM	2
841	2633			4 14 0.8	-0.349	1	160 46 26.2	8.96	1	Cl; vL; ab 20 sc st	1
842	2632			4 14 56.3	+1.858	1	135 21 58.1	8.83	1	pF; S; E; gbM	2
843	2634			4 15 56.3	0.704	2	153 7 47.6	8.78	2	vB; vL; mE; vgpmbM; *14	2
										attn.	
844	2635		Δ. 338??	4 16 52.1	1.337	2	145 16 34.3	8.69	2	B; vL; vg, svmbM; 15 ^a d	2
										in R.A.	
845	2636			4 17 9.1	1.707	1	138 35 23.2	8.66	1	F; S; R; bM	2
846	2637			4 17 42.3	1.916	1	133 47 30.7	8.61	1	F; S; R; gbM	1
847		II. 768		4 17 43.5	5.621	1	25 27 13.7	8.51	1	pB; S; IE; bNM; pB* ⁿ ...	1
848	2638			4 17 45.5	1.910	1	133 57 0.7	8.61	1	vF; S; R; gbM; *nf	1
849	2639			4 17 59.3	2.026	1	130 55 4.6	8.58	1	pF; S; R; *13 nf 1'	1
850	2640			4 19 13.4	1.190	1	147 17 51.7	8.51	1	pB; S; R; pgbM; 2S st sf ...	1
851	314	III. 587		4 19 24.3	2.989	1	93 56 58.5	8.45	1	eF; bM; bet 2 st	2*
852	2641			4 20 12.4	1.526	3	141 55 9.4	8.42	3	pF; S; R; bM	3
853	315	I. 217		4 21 2.8	3.925	3	55 2 25.3	8.29	3	pB; vL; iR; mbM; *8,	6+
										350°, 2'.	
854	2642			4 21 37.2	1.321	2	145 15 47.7	8.31	2	F; S; E; glbM	2
855		VIII. 70		4 22 9.3	4.233	1	46 27 35.0	8.20	1	Cl; vL; pRi; IC; stL	1
856	2643			4 22 58.1	1.957	1	132 27 41.3	8.19	1	pF; S; R; gbM; *12, 287° 8	1
857		D'Arrest, 43		4 23 5	3.05	[1]	90 50 6	8.15	[1]	vF; iF; vlbM; bet * & *14..	0
858	316	II. 8		4 23 29.1	3.081	1	89 39 14.4	8.12	1	F; pS; R; r; p of D neb	5
859	317	II. 9		4 23 32.6	3.081	1	89 38 36.7	8.11	1	F; vS; R; r; f of D neb	5
860	318	II. 7		4 23 33.5	3.085	1	89 26 33.7	8.11	1	F; pL; IE 132°; *42°, 80'' ...	5
861	2644			4 23 48.7	2.450	3	117 0 49.0	8.10	3	pF; pS; R; gbM	3
862	2645			4 23 52.7	2.443	1	117 15 57.0	8.10	1::	vF; vS; R	1
863	2646			4 24 24.8	1.710	1	138 7 6.6	8.08	1	vF; S; R; bM	1
864	2648			4 24 35.0	1.305	2	145 20 17.6	8.08	2	B; pL; mE 15° 0; smbM; p	2
										of 2.	
865	2647			4 24 36.3	1.711	1	138 5 22.2	8.06	1	F; S; R; bM	1
866	319	I. 158		4 24 45.0	2.958	4	95 23 3.4	8.02	4	pB; pL; R; gmbM	6
867	319, a		R. 3 novæ	4 24 ±		...	95 23 ±		...		0
868											
869											
870	2649			4 24 48.1	1.303	2	145 22 1.2	8.06	2	eF; pL; IE; f of 2	2
871		VI. 26		4 24 59.0	4.298	1	45 3 48.9	7.97	1	Cl; vF; pS; C; steS	1
872		III. 585		4 25 45.2	2.973	1	94 38 56.8	7.94	1	Susp in hazy weather	1
873		III. 586		4 26 10.8	2.975	2	94 33 58.0	7.90	2	eF; S; E 90° ±	2
874	2650			4 28 20.2	1.883	3	134 0 32.2	7.76	3	F; S; E; vglbM	3
875	2651		Δ. 339??	4 28 33.6	+1.321	2	144 54 14.2	-7.76	2	B; L; mE 105° 8; } vg, vsmbMN5''.	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and m.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
876	320	II. 524		4 29 6.8	+2.999	1	93 26 37.9	-7.67	1	F; S; iF; lbM; 2st sf	2
877	321	II. 514		4 29 26.0	3.064	2	90 25 50.1	7.63	2	vF; pL; mE0°...90°; B *nf..	3
878	320, a		D'Arrest, 44	4 29 34	3.00	[2]	93 28 48	7.63	[2]	vF; S; *20, 270°, 5°; II. 524 p (R).	0
879		V. 49		4 29 47.5	4.558	1	39 50 13.9	7.57	1	F; cL; iF; 6 or 7 st+neb ...	1
880	322			4 30 7.9	2.995	1	93 35 45.6	7.58	1	vF; E90°...180°; sbM; B*p 40°.	1*
881	322, a		R. nova	4 30 19.9	+2.995	::	90 30 45.6	7.58	::	MS	0
882	2653			4 31 14.0	-0.756	1	162 8 7.3	7.59	1	vF; pL; R; glbM	1
883	2652			4 32 20.0	+2.598	1	110 55 34.4	7.42	1	Neb. No description	1
884	323	III. 952		4 32 36.3	3.227	2	82 56 7.6	7.38	2	eF; S; R; *8 sp; p of D neb	3
885	324	III. 953		4 32 36.5	3.226	1	82 57 2.6	7.38	1	eF; vS; f of D neb	2
886	325	II. 515		4 33 1.4	3.055	1	90 49 38.5	7.35	1	F; S; R; bM; *9 nf 12° 5 ...	2
887	{ 326 = 2654 }	II. 522		4 33 57.4	2.878	2	98 52 48.9	7.27	2	vF; pS; R; vgbM; r; *nf 1'.	4
888	327	I. 122		4 34 26.6	3.005	2	93 8 53.1	7.23	2	cB; L; R; vgbM; er	3†
889	2655			4 34 36.9	2.686	1	107 16 3.1	7.23	1	eF; vS; R; bet 2 st	1
890		II. 525		4 35 29.3	3.028	1	92 2 39.5	7.15	1	F; pL; lE	1
891	2656			4 35 30.1	0.266	1	156 4 46.4	7.22	1	Cl; pL; pRi; pnc; st11...16	1
892			D'Arrest, 45	4 35 45	3.08	[1]	89 39 1	7.12	[1]	F; R; cometary; Δ with 2 st 18, f.	0
893	328	III. 588		4 36 51.6	2.950	1	95 35 42.8	7.04	1	eF; vS; iR; bM	2
894	2657			4 37 7.2	0.206	3	156 27 51.3	7.09	3	F; S; R; gbM	3
895	329	II. 523		4 37 41.4	2.879	1	98 47 13.9	6.97	1	F; vS; iR; bM; *7 np	2
896		VIII. 8		4 37 54.5	+3.500	2	71 11 21.8	6.94	2	Cl; vL; stL, sc	2
897	2660			4 38 37.3	-0.201	1	159 5 7.6	6.98	1	F; pS; R; gbM	1
898	2662			4 38 42.3	-0.531	3	160 51 32.6	6.98	3	pF; L; vLE; vglbM	3
899	2661			4 38 43.9	-0.177	3	158 56 26.9	6.97	3	vF; S; R; glbM	3
900		II. 526		4 38 49.7	+3.015	1	92 38 59.9	6.87	1	F; cS; R; lbM	1
901	330			4 39 0.9	2.954	1	95 24 18.2	6.86	1	eF; iF; ?	1
902	2658			4 39 28.8	1.952	1	131 45 5.5	6.85	1	F; pS; pnc; glbM	1
903	331	III. 589		4 39 36.7	2.962	1	95 2 37.7	6.81	1	pF; pS; iE90°+; bM	2
904	2659			4 39 37.6	1.950	1	131 46 55.1	6.83	1	vF; S; iE; glbM	1
905	332	VII. 1		4 40 44.9	3.310	1	79 19 18.7	6.71	1	Cl of L & S sc st	3
906		VIII. 7		4 40 45.1	3.361	2	77 6 7.7	6.71	2	Cl; iRi; st L & S	3
907		VIII. 59		4 41 2.4	4.268	1	46 33 17.5	6.65	1	Cl; iRi; iC; pL	1
908	333	II. 547		4 41 23.7	2.947	1	95 41 12.2	6.66	1	eF; pL; R; lbM	3*
909	2663			4 41 59.7	1.811	1	135 2 6.8	6.64	1	eF; R; att to *14	1
910	2664			4 42 20.4	0.236	1	156 4 5.2	6.66	1	eF; S; R	1
911		III. 501		4 42 46.2	3.007	1	93 0 18.8	6.54	1	vF; vS	1
912	2665		Δ. 296 ??	4 43 32.0	+0.930	2	149 30 3.8	6.54	2	B; L; smbMN	2
913	2667			4 43 34.7	-0.404	1	160 4 2.6	6.58	1	vF; S; att to *10	1
914	2669			4 44 19.2	-0.228	1	159 4 32.7	6.51	1	vF; pL; iR; r	1
915		III. 502		4 44 35.4	+3.010	1	92 51 28.0	6.40	1	vF; S	1
916	2666			4 44 39.1	2.276	2	122 12 45.4	6.42	2	vB; L; iR; 4st inv	2
917	2668			4 44 41.5	1.665	3	138 3 31.4	6.42	3	vF; S; R; r or st inv	3
918	334, a		R. nova	4 45 16.3	2.999	::	93 14 46.4	6.32	::	R, MS	0
919			D'Arrest, 46	4 45 22	3.00	[2]	93 21 0	6.33	[2]	vF; vS; II. 527, f 12°+	0
920	334	II. 527		4 45 32.3	2.999	2	93 20 46.4	6.32	2	pF; S; R; bM; *7, 225°±...	4
921	334, b		R. nova	4 45 44.3	2.999	::	93 20 46.4	6.32	::	MS } No description	0
922	334, c		R. nova	4 45 44.3	2.999	::	93 8 46.4	6.32	::	MS }	0
923	2670			4 46 9.1	2.212	2	124 10 25.0	6.30	2	vF; S; R; vglbM	2
924		II. 528		4 46 12.6	2.999	1	93 18 35.2	6.26	1	F; S; lbM	1
925	2671			4 46 18.6	0.875	3	150 2 6.7	6.31	3	pB; pL; iR; pgmbM	3
926	335			4 47 6.7	+3.104	1	88 35 49.3	6.19	1	vF; vS; am vSt; L*sp	1*
927	2672			4 48 23.2	-0.339	2	159 35 14.9	6.17	2	F; S; R	2
928	2673			4 48 30.2	0.345	2	159 37 5.2	6.16	2	F; S; R	2
929	2674			4 48 47.0	0.146	1	158 27 26.8	6.14	1	vF; E; vlbM	1
930	2675			4 49 1.9	0.203	5	158 47 31.4	6.12	5	⊕; pB; L; R; rr	5
931	2677			4 49 42.0	-0.302	1	159 20 44.2	-6.06	1	pB; pS; R; glbM	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s		° ' "	"			
932	336	IV. 32		4 50 1.9	+2.959	1	95 5 43.5	-5.95	1	eB; S; mbM*.....	3
933	2676			4 50 25.3	+2.333	3	120 6 11.1	5.93	3	F; S; vIE; glbM; *10, 75".....	3
934	2680		Δ. 73?	4 50 27.0	-0.443	1	160 5 32.0	6.00	1	Cl; vF; S	1
935	2678			4 50 48.7	+0.866	2	149 58 0.1	5.93	2	F; L; R; vglbM; * att.....	2
936	2683			4 50 53.7	-0.427	2	159 59 14.9	5.97	2	F; pS; IE; r	2
937	2679			4 51 8.5	+1.342	3	143 35 14.3	5.89	3	pF; S; R; pmbM	3
938	2682			4 51 15.3	0.544	1	153 13 27.7	5.91	1	F; pS; R; vglbM	1
939	338			4 51 15.6	3.253	1	81 58 39.1	5.83	1	S; R; rrr	1
940	337			4 51 32.4	+4.753	1	37 19 51.9	5.77	1	Cl; vL; pRi; IC; stL and S	1
941	2684		Δ. 76?	4 51 44.5	-0.473	2	160 13 0.0	5.90	2	⊕; B; S; iR; rrr; st14	2
942	2685			4 51 46.1	-0.370	1	159 39 49.3	5.89	1	Cl; pB; S	1
943	339	II. 516		4 51 47.1	+3.057	2	90 42 45.3	5.79	2	F; S; R; bM; p of 2	2
944	339, a		R. nova	4 51 ±			90 42 ±			No description	0
945	2686			4 52 0.4	0.043	5	157 8 55.2	5.86	5	vB; S; Eorbi-N; bM; spof2	5
946	2687			4 52 2.6	0.046	3	157 7 48.2	5.86	3	vF; S; R; sbM; 2stnr; nfof2	3
947	2681			4 52 16.4	2.592	1	110 34 42.9	5.77	1	pF; pL; R; glbM	1
948	340			4 52 18.9	+3.063	1..	90 28 7.5	5.75	1..	pF; S; iR; psbM	2
949	340, a		R. nova	4 52 ±			90 28 ±			No description	1
950	2688			4 52 20.4	+0.022	3	157 16 42.8	5.84	3	F; pS; R; vglbM	3
951			D'Arrest, 47	4 52 39	+2.89	[2]	98 4 0	5.73	[2]	pF; pL; lbM; h. 341 nr.....	0
952	2689			4 52 41.7	-0.366	3	159 37 11.4	5.82	3	Cl; pF; S; R; 2nd of 3	3
953	341			4 52 57.2	+2.893	1	97 58 11.7	5.71	1	F; R; *13, s	1*
954	2690			4 53 0.9	-0.358	3	159 34 7.3	5.79	3	Cl; pB; pS; pmE; st12	3
955		III. 503		4 53 21.2	+2.994	1	93 32 9.9	5.67	1	vF; pL; 2B st v nr.....	1
956	2691			4 53 22.6	+0.038	3	157 8 36.5	5.75	3	Cl; pL; lRi; IC; st10...15...	3
957	2694			4 53 43.2	-0.238	2	158 52 23.4	5.72	2	S; R; close * in M	2
958	2693			4 53 49.2	+0.073	1	156 53 44.7	5.71	1	eF; pL; R; gbM.....	1
959	2695			4 54 8.4	-0.259	1	158 59 17.0	5.70	1	pB; L; R; gmbM	1
960	2696			4 54 18.5	+0.009	1	157 18 56.9	5.67	2	pF; pS; R; 2st att	2
961	2697			4 54 19.0	-0.143	1	158 17 20.9	5.67	1	B; R; r	1
962	2698			4 54 31.9	-0.335	1..	159 24 10.2	5.66	1..	vF; S; 1st of 4	1
963	2699		Δ. 114	4 54 49.4	-0.339	2	159 25 17.8	5.64	2	B; pL; R; gbM; r; 2nd of 4	2
964	2692			4 54 51.1	+2.440	1	116 14 38.2	5.56	1	F; vL; vmE; vglbM	1
965	342			4 54 58.2	+2.994	1	93 30 26.1	5.53	1	eF; vS; *12, sf	1
966	2702			4 55 7.9	-0.335	1..	159 23 11.7	5.61	1..	F; S; 3rd of 4	1
967	2701			4 55 12.5	-0.006	2	157 23 32.0	5.60	2	Cl; pS; lRi; stvS	2
968	2704			4 55 13.8	-0.340	1	159 25 6.0	5.60	1	pB; vS; R; 4th of 4	1
969	2703			4 55 16.5	-0.166	1	158 24 26.3	5.59	1	vF; R; p of 2	1
970		VIII. 43		4 55 18.9	+3.630	1	66 33 18.3	5.49	1	Cl; stL, vsc	1*
971	2705			4 55 19.4	-0.456	1	160 1 59.0	5.60	1	eF; pL; iR	1
972			D'Arrest, 49	4 55 26	+2.88	[2]	98 26 48	5.49	2	F; pL; pmE; 2 or 3st11nf...	0
973	2708			4 55 36.9	-0.581	1	160 39 49.9	5.57	1	F; S; R; *13att, 135°	1
974	2707			4 55 37.8	-0.350	1	159 27 21.9	5.57	1	vF; S; R.....	1
975	343			4 55 45.0	+2.962	1	94 55 35.9	5.47	1	vLdiff neb in zigzags??	1*
976	2706		Δ. 167	4 55 49.5	0.266	1	155 25 43.1	5.53	1	vB; pL; R; gbM; f of 2	1
977		VII. 21		4 55 55.7	3.634	1	66 25 21.1	5.43	1	Cl; pC; stL and S	1
978	2700			4 56 5.5	2.037	1	128 55 2.2	5.46	1	vF; pL; vglbM	1
979	2709			4 56 15.6	0.087	1..	156 43 53.0	5.50	1..	vF; S; 3vSst inv.....	1*†
980	2710			4 56 19.0	0.093	3	156 41 5.0	5.50	3	Cl; L; mC; * 9	3†
981		III. 453		4 56 24.5	3.104	1	88 34 31.3	5.39	1	vS; vF	1*
982	2711			4 56 31.9	+0.102	5	156 37 13.3	5.49	5	vB; vL; vimE.....	5†
983	2713			4 56 42.0	-0.092	1	157 54 38.9	5.47	1	vF; S; R.....	1
984	2712			4 56 51.7	+0.631	2	152 14 11.8	5.44	2	cF; S; R; glbM	2
985	2717			4 57 15.1	-0.543	2	160 26 16.8	5.44	2	cF; S; gbM	2
5062				4 57 18.4			159 36 32.1			See No. 5062	1
986	2718			4 57 22.2	-0.185	1	158 28 4.4	5.42	1	F; S; R; gbM	1
987	2716			4 57 29.6	+0.091	5	156 39 56.0	5.40	5	B; L; iR; vsmbM * 10.....	5†
988	2715		Δ. 169	4 57 39.7	-0.213	2	158 37 42.3	5.39	2	Cl+neb; pL; pRi; st11...18	2
989	2720			4 57 48.0	+0.509	1	153 20 44.2	5.36	1	vF; mE; glbM; *7, 8np	1
990	2722			4 57 48.9	-0.416	3	159 46 2.3	5.39	3	pB; pS; iR; rr	3
991	2721			4 57 50.3	+0.105	1	156 33 59.9	-5.37	1	pF; pL; iR; 2 or 3Bst nr	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. II.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
992	2723			4 58 11.0	-0.027	3	157° 27' 6.5	-5.35	3	B; S; R; smbM; *+neb ...	3
993	2725			4 58 18.6	-0.590	1	160 38 26.5	5.35	1	eF; pL; iR	3
994	2724			4 58 23.2	+0.093	1	156 38 1.4	5.32	1	vF; S; R; gbM	1
995	2728			4 58 28.2	-1.562	1	164 29 44.2	5.36	1	eF; E; * 9att, f	2
996	344	VIII. 61		4 58 35.7	+4.041	1	53 8 20.3	5.19	1	Cl; pC; lRi; iF; stL	2
997	{ 345 = 2714 }	III. 500		4 58 37.6	+2.860	3	99 20 30.1	5.23	3	pB; S; R; gpmbM	4
998		III. 268		4 58 42.9	+2.643	1	108 22 53.0	5.31	1	eF; vS; stellar	1*
999	2727			4 58 43.5	-0.390	2	159 36 16.7	5.31	2	⊕; pB; S; R; pmbM; rr ...	2
1000	2726			4 58 44.0	+0.153	2	156 11 37.6	5.28	2	cB; L; R; vgpmB; r	2
1001	2719			4 58 56.6	+2.797	1	102 4 3.7	5.21	1	pB; pL; vLE; vgbM; am st ..	1
1002	(147)			4 59 12.2	-0.294	1	159 3 36.9	5.27	1	No description	1
1003	2729			4 59 20.7	-0.109	3	157 57 2.5	5.25	3	vB; pS; R; vsymbM*9 ...	3
1004	2731			4 59 52.1	+0.170	1	156 2 39.3	5.19	1	Cl; vL; pRi	1
1005		V. 32		4 59 55.9	+2.993	2	93 32 59.7	5.11	2	B; cL; R; bM * 15; *10, 318°.	4
1006	2733			4 59 57.5	-0.926	1	162 5 37.4	5.22	1	vF; pS; R; vglbM	1
1007	346			5 0 12.6	+4.723	1	38 7 8.8	5.04	1	Cl group of 8 or 9 st10	1
1008	2734			5 0 20.2	-0.542	1	160 21 49.6	5.18	1	eF; S; R	1
1009	2730		Δ. 531?	5 0 23.9	+2.056	2	128 11 33.0	5.10	3	vB; vL; mE314°; glbM; rr.	3
1010	2736			5 0 36.4	-0.426	2	159 45 22.5	5.15	2	F; S; R; glbM	2
1011	2738		Δ. 81	5 0 58.7	-0.472	1	159 59 38.4	5.12	1	F; pL; iE	1
1012	2735			5 1 1.0	+0.710	2	151 19 31.6	5.08	2	pB; pS; pmE; vglbM	2
1013	2732			5 1 10.9	+2.260	1	122 8 40.1	5.03	1	pB; pmE; gpmbM; *13f ...	1
1014	2739			5 1 34.1	-0.440	4	159 48 27.9	5.07	4	F; pL; R; vglbM; p of 2 ...	4
1015		VIII. 41		5 1 44.4	+3.647	1	66 4 49.8	4.94	1	Cl; st c sc	1
1016	2737			5 1 51.5	+1.544	1	139 45 48.3	4.99	1	F; S; R; vglbM; *11sf; ? neb	1
1017	2742			5 1 51.9	-0.344	2	159 16 55.8	5.04	2	F; S; R; bM	2
1018	2741		Δ. 233?	5 2 1.9	+0.128	5	156 18 4.4	5.02	5	B; vS; vsmbM; st+neb	5
1019	2745			5 2 31.5	-0.159	1	158 11 17.6	4.98	1	pB; L; gbM	1
1020	348			5 2 37.5	+3.453	1	73 39 38.9	4.87	1	Cl; pRi; stL and S	1
1021	2740		Δ. 549	5 2 50.1	+2.071	2	127 41 47.0	4.90	2	B; L; E; psbM	2
1022	2747			5 3 4.7	-0.448	1	159 49 11.8	4.94	1	pF; S; R; gbM; 2nd of 2 ...	1
1023	2746		Δ. 235	5 3 9.1	+0.086	5	156 34 15.4	4.92	5	eF; S; R; lbM; ⊕f	5
1024	2743			5 3 16.9	+2.340	2	119 27 56.5	4.85	2	cF; S; iE; p of 2	2
1025	2744			5 3 27.3	+2.341	2	119 26 7.1	4.83	2	F; S; R; glbM; f of 2	2
1026	2752			5 3 54.8	-0.586	1	160 30 43.6	4.88	1	vF; S; R; r	1
1027	2748			5 3 55.4	-0.054	2	157 29 41.2	4.86	2	vF; R; s of 2 in Cl	2
1028	2753			5 3 56.0	-0.648	2	160 48 33.6	4.88	2	F; vS; R; vlbM; am st	2
1029	2750			5 3 58.7	-0.048	1	157 27 2.5	4.85	1	vF; R; 2nd neb in Cl	1
1030	349	VII. 4		5 3 59.1	+3.458	2	73 28 42.5	4.75	2	Cl; L; Ri; iC; st11...14 ...	5*
1031	2749		Δ. 236	5 4 2.0	+0.076	6	156 37 3.5	4.85	6	⊕; vB; pL; R; vmC; rr ...	6
1032	2754			5 4 22.1	-0.049	1	157 26 48.4	4.82	1	Cl; pL; Ri; C; iF	1
1033	2756			5 4 47.9	+0.104	1	156 23 46.6	4.78	1	vF; S; p of 2	1
1034	2758			5 4 51.8	-0.595	1	160 31 53.0	4.80	1	Cl; pF; L; iF; st12...15 ...	1
1035	2755			5 4 53.3	+0.834	2	149 54 12.5	4.75	2	vF; pL; vmE162°0	2
1036	(199)			5 5 2.2	-0.328	1;	159 7 51.0	4.80	1;	No description	1
1037	2757			5 5 10.7	+0.101	1	156 24 57.5	4.75	1	vF; S; f of 2	1
1038	2751			5 5 12.8	+2.088	1	127 9 17.3	4.69	1	vF; vmE; long ray; *11inv.	1
1039	2761			5 5 16.9	-0.411	4	159 34 33.2	4.76	4	F; S; R; 1st of 3	4
1040	2760			5 5 22.7	-0.177	1	158 14 14.8	4.74	1	F; pL; R; r	1
1041	2762			5 5 33.9	-0.403	4	159 31 45.1	4.73	4	F; pS; R; 2nd of 3	4
1042	2759		Δ. 246	5 5 35.8	+0.275	2	155 6 40.7	4.71	2	B; L; R; glbM; r	2
1043		II. 292		5 5 52.9	+2.702	1	105 53 13.4	4.62	1	pB; iR; mbM; cSnfl'	1
1044	2765			5 5 56.2	-0.679	1	160 55 5.7	4.71	1	vF; pL; 1st of sev	1
1045	2764			5 6 3.3	0.381	1	159 23 56.5	4.68	1	○? B; eS; iE	1
1046	2763			5 6 3.7	0.415	4	159 35 14.6	4.68	4	cB; S; R; gmbM; 3rd of 3..	4
1047	2766			5 6 12.5	0.278	2	158 48 40.9	4.67	2	st+neb; 1st of sev	2
1048	2769			5 6 28.0	0.677	2	160 53 56.2	4.66	2	Cl; L; Ri; st sc	2
1049	2767		Br. 895	5 6 32.8	-0.246	1	158 37 28.8	-4.64	1	Cl; L; vLC	1

No. of Cata- logue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
1050	2768		Δ. 170?	5 6 38.4	-0.278	2	158 48 37.5	-4.65	2	st+neb; pB; iF; 2nd of sev	2
1051	2771			5 6 46.2	0.909	1	161 56 6.8	4.64	1	F; R; bM; r(? min. of R.A.)	1
1052	2788			5 7 1.6	9.684	1	174 12 38.6	4.88	1	pF; L; iR; vsbM; r	1
1053	2772			5 7 20.8	0.060	2	157 27 15.9	4.57	2	vvF; R; p of 2	2
1054	2773			5 7 36.6	0.068	4	157 29 56.8	4.54	5	pF; pL; R; gbM; f of 2	5
1055	2770			5 7 38.3	0.651	2	160 44 41.2	4.56	2	Cl; vICM; st 9, 11...16	2
1056	2774			5 7 45.1	0.089	3	157 37 57.8	4.54	4	pB; eL; R; vglbM; r	4
1057	2775			5 7 56.4	0.343	2	159 8 54.1	4.53	2	B; S; iE; * in M	2†
1058	2776			5 8 5.8	-0.785	1	161 21 52.1	4.53	1	Cl; vIC; st 9, ...	1
1059	2778			5 9 20.8	+0.078	1	156 28 44.3	4.39	1	vF; S; iE; glbM	1
1060	2780		Δ. 170?	5 9 25.2	-0.309	6	158 55 47.0	4.40	6	⊕! vB; L; iE; vmCM; rr	6
1061	2777		Δ. 508	5 9 29.8	+1.970	2	130 12 20.1	4.33	2	⊕! vB; vL; R; vsvvbM; rrr.	3
1062	2781			5 9 41.5	-0.145	4	157 56 58.9	4.37	4	F; pL; R; vglbM	4
1063	2779			5 9 51.6	+1.019	3	147 33 40.1	4.33	3	F; S; mE 45°; vglbM; *11 nf.	3
1064	2782			5 10 2.3	-0.327	5	159 1 12.5	4.35	5	⊕; eB; S; R; gbM; 2d of 3	5
1065	2783			5 10 8.1	0.326	1	159 0 38.8	4.34	1	Cl; vB; L; R; st12	1
1066	2784			5 10 20.3	-0.379	3	159 17 59.1	4.33	3	B; pL; R; gbM; 12' diam R.A.	3
1067	350	VII. 33		5 10 26.0	+4.138	1	50 48 30.6	4.18	1	Cl; pRi; pC; st 7, ...	2
1068	2785			5 10 37.2	-0.337	7	159 3 44.0	4.30	7	B; L; iE; biN; Cl+neb	7
1069	2786			5 10 59.4	+0.220	1	155 24 46.2	4.26	1	F; S; R; vgbM; * 7 nf 6'	1
1070	2787		Δ. 172?	5 11 21.6	-0.316	1	158 55 38.1	4.23	1	F; pL; R; vgbM	1
1071	2790			5 12 2.3	-0.708	1	160 56 56.6	4.18	1	eF; pL; R; vglbM	1
1072	2789			5 12 6.1	+0.094	1	156 18 45.2	4.16	1	pF; L; iR; vgbM; r	1
1073	2791		Δ. 173?	5 12 19.1	-0.312	5	158 53 22.5	4.15	5	vB; vS; R; r or stellar	5
1074	2792			5 12 56.3	0.130	2	157 47 15.6	4.08	2	F; pS; iR; bM; r or stellar	2
1075	2794		Δ. 173??	5 13 6.0	-0.321	3	158 55 37.6	4.08	3	vF; pL; R; vglbM	3
1076	2793		Δ. 247? 248?	5 13 11.3	+0.185	1	155 37 35.2	4.06	1	vB; L; R; vgmBM; r	1
1077	2795			5 13 27.1	+0.071	1	156 27 4.8	4.04	1	eF; pL; R	1
1078	2796			5 13 41.5	+0.376	2	154 6 36.7	4.01	2	pB; pL; R; vglbM	2
1079	2798		Δ. 210	5 13 58.4	-0.092	3	157 32 7.0	4.00	2	Cl; L; pRi; st sc	2
1080	2799			5 13 59.0	0.386	2	159 16 29.7	4.01	2	B; S; R; glbM	2
1081	2800			5 14 5.5	0.105	2	157 36 46.0	4.00	2	Cl; iRi; 2nd of sev	2
1082	2802			5 14 7.8	0.423	4	159 28 12.0	4.00	4	pB; R; gbM; 1st of group	4†
1083	2801			5 14 8.8	0.086	1	157 29 23.3	3.99	1	Cl; 3rd of sev	1
1084	2803			5 14 9.6	0.434	1	159 31 55.0	4.00	1	neb & Cl; biN	1†
1085	2804			5 14 16.2	0.432	4	159 31 1.6	3.98	4	pB; iR; biN; 2nd in group	4†
1086	2805			5 14 19.0	0.434	1	159 31 50.6	3.98	1	vF; 3rd of group in Cl	1†
1087	2807			5 14 26.2	-0.651	1	160 37 52.6	3.98	1	vF; iE; vglbM; r	1
1088	2797			5 14 37.2	+2.244	1	122 17 45.6	3.88	1	vF; L; R; vglbM; *12p	1
1089	2808			5 14 37.9	-0.435	1	159 31 47.2	3.96	1	4th N, of neb in Cl	1†
1090	2810			5 15 3.1	-0.421	1	159 27 2.4	3.92	1	vF; * p	1†
1091	2809			5 15 15.7	+0.091	1	156 16 34.6	3.88	1	pF; R; vglbM; r	1
1092		VII. 34		5 15 28.7	4.452	1	43 35 54.8	3.74	1	Cl; vF; pRi; pC; iF	1
1093	2812			5 15 45.0	0.084	1	156 18 45.8	3.84	1	eF; pL	1
1094	2814			5 15 47.4	0.362	1	159 7 24.2	3.86	1	pB; vS; R; bM	1
1095	2813			5 15 53.7	0.061	1	156 28 16.1	3.83	1	vF; vS; R; *p25"	1
1096	{ 352 = 2806 }	II. 289		5 16 3.7	+2.802	2	101 38 3.8	3.74	2	pB; pL; R; r	2
1097	352, α		R. nova	5 16		...	101 38		...	Makes a close D neb with h. 352.	0
1098	2816			5 16 5.8	-1.017	1	162 13 55.5	3.85	1	vF; S; R; glbM	1
1099	2811			5 16 20.8	+2.124	1	125 51 33.8	3.74	1	Cl; L; Sc; * taken	1
1100	2815			5 16 31.9	+0.245	2	155 6 43.9	3.77	2	eF; pL; E 90°±; vglbM	2
1101	351			5 16 32.3	+3.935	1	56 44 33.9	3.67	1	Cl; L; Ri; iC	1
1102	2818			5 16 56.7	-0.457	1	159 36 42.2	3.76	1	F; pL; R; sbM; r; st inv	1
1103	2817			5 17 3.0	-0.092	4	157 28 36.8	3.74	4	pF; pL; R; vglbM	4
1104	353	VIII. 4		5 17 8.1	+3.802	1::	69 58 11.1	-3.63	1::	Cl; vIRi; vIC; st 9...12	3

No. of Cata- logue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by H. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
1105	2820			5 17 44.0	-0.112	1	157 35 20.6	-3.68	1	eF; S; R	1
1106	2822			5 17 51.1	-0.495	1	159 47 44.6	3.68	1	F; pS; R	1
1107	2821			5 17 58.4	-0.180	1	158 1 7.2	3.66	1	F; pS; R; vglbM; 3st 10p ...	1
1108	2819			5 18 1.2	+0.475	2	153 10 14.8	3.64	2	F; pL; iE; vglbM; *7np ...	2
1109	2824			5 18 7.8	-0.302	1	158 44 6.2	3.66	1	Cl; BM; iRi; st 7	1
1110	2823			5 18 14.1	+0.013	5	156 46 1.1	3.63	5	⊕; pB; pL; R; pmbM; rr ...	5
1111	2825			5 18 22.1	-0.436	5	159 28 39.8	3.64	5	vB; S; R; gmbM	5
1112			M. 79	5 18 25.6	+2.469	A	114 39 39.5	3.55	A	⊕; pL; eRi; eC; rrr	4
1113	4016			5 18 36.3	-0.087	1	157 25 24.7	3.61	1	F; S; R; r	1
1114	354	VII. 39		5 18 42.6	+4.001	3	54 48 28.6	3.48	3	Cl; pRi; pC; R; st 9...12 ...	5
1115		V. 33		5 18 52.3	3.011	1	92 39 22.0	3.30	1	v diffused neb. susp	1
1116		V. 38		5 19 10.4	+2.881	1	98 15 15.9	3.47	1	eL; strongly susp (2° in P.D.).	1
1117	2827		Δ. 129	5 19 10.4	-0.416	4	159 21 30.9	3.57	4	Cl; L; pRi; iR; st 11...16 ...	4
1118	2826			5 19 13.2	-0.010	1	156 54 4.5	3.55	1	F; R; gB; am st	1
1119			M. 38	5 19 17.0	+4.020	(2)	54 17 36.1	3.43	(2)	Cl; B; vL; vRi; iF; st L & S	7
1120	(356)			5 19 33.6	-0.480	1	159 41 31.8	3.54	1	No description	1
1121	2830			5 19 35.7	0.833	2	161 23 42.8	3.54	2	F; L; iE	2
1122	2828			5 19 36.7	0.016	1	156 56 22.4	3.52	1	eF; pL	1
1123	2829			5 19 39.1	0.453	3	159 32 47.1	3.53	3	B; S; R; vgvmbM; r	3
1124	2831			5 19 53.3	0.379	1	159 8 15.0	3.50	1	vF; L; R; vglbM	1
1125	(369)			5 20 10.1	0.496	1	159 46 33.6	3.48	1	No description	1
1126	2832			5 20 15.1	0.030	1	157 0 57.2	3.46	1	Cl; eF; L; iR; mC; rr	1
1127	2832			5 20 33.0	0.014	7	156 54 44.8	3.44	7	pB; pL; R; vgbM	7
1128	2834			5 20 33.9	0.527	1	159 55 8.5	3.45	1	vF; pS; iE; r	1
5063				5 20 52.9	-0.469	1	159 36 35.4	3.42	1	(See No. 5063)	1
1129	2835			5 21 6.5	+0.168	1	155 36 58.6	3.38	1	vF; pS; R	1
1130		III. 447		5 21 8.1	2.947	2	95 26 56.0	3.30	2	vF; pL; iR; st nr	2
1131	2837			5 21 22.2	+0.113	1	156 0 17.2	3.36	1	Cl; vRi; iC; st 10	1
1132	2838			5 21 40.7	-0.480	3	159 39 51.5	3.35	3	pB; pL; iR; r; in diff n	3
1133	356			5 22 0.7	+2.875	1	98 29 41.4	3.22	1	Diffused nebosity	1*
1134	2839		Δ. 131	5 22 1.5	-0.473	2	159 36 55.4	3.32	2	pF; pL; R; gB	2
1135	2840			5 22 1.9	-0.198	1	158 3 33.7	3.31	1	F; p of group	1†
1136	2836			5 22 3.0	+1.666	2	136 51 6.2	3.26	2	pF; S; R; bM; 4B st p	2
1137	355	I. 261		5 22 10.0	3.968	3	55 52 3.6	3.18	3	vB; L; R; b * in M	4†
1138				5 22 14.9	+0.074	5	156 16 28.3	3.29	5	{ pB; S; R; smbM } D neb	5*
1139	2841			5 22 21.3	-0.197	1	158 2 52.6	3.28	1	{ eF; R; stellar } 26°, 80"	1†
1140	2842			5 22 24.9	0.204	4	158 5 19.6	3.28	4	2nd neb of group	4†
1141	2843			5 22 41.6	0.207	6	158 6 22.5	3.25	6	pF; S; R; 3rd of group	6†
1142	2844		Δ. 175	5 22 43.6	0.195	1	158 1 49.5	3.25	1	pB; S; R; 4th of group ...	1†
1143	2845			5 22 45.6	0.564	4	160 4 39.2	3.26	4	vF; pL; follows a group	4
1144	2848		Δ. 89?	5 22 54.5	-0.082	2	157 18 47.1	3.23	2	{ pB; pS; R; gB } D neb	2
1145	2847			5 22 55.3	+0.099	1	156 30 33.4	3.22	1	{ F; S; R; gB } 339°-1, 50"	1
1146	2846			5 23 51.4	+0.358	2	154 4 25.8	3.14	2	pB; vS; R; bM; 2st 9 & 10 f	2
1147	2849			5 23 53.2	-0.608	3	160 16 48.9	3.17	3	vS; neb+st	1
1148	2850		Δ. 90	5 24 26.6	-1.138	1	162 36 15.8	3.14	1	eF; stell; *14+ neb	2
1149	2850			5 24 48.3	+0.025	1	156 34 43.9	3.07	1	pF; pS; iR; vglbM; *15, 190°-6, 60"	3
1150	2852			5 25 8.5	0.034	1	156 30 29.8	3.04	1	pB; pL; R; bM	1
1151	2851			5 25 37.5	0.255	1	154 52 39.3	2.99	1	eeeF; vvL; irr diff	1
1152	2854		Δ. 237?	5 25 40.3	+0.051	1	156 23 6.3	2.99	1	pF; R; gB; r	1
1153	2855			5 25 47.4	-0.296	1	158 35 32.3	2.99	1	pB; L; R; gB; *9np	1
1154	2856			5 25 50.5	-0.561	2	160 1 23.3	2.99	2	Cl; cL; Ri; st 13	1
1155	2857			5 26 3.9	+3.605	1	68 5 10.5	2.85	1	pB; S; R; psbM	1
1156	2859			5 26 4.1	+0.003	2	156 42 31.2	2.96	2	The 1st of a group of 7 !	2†
1157	357		M. 1	5 26 16.7	-0.362	2	158 57 2.5	2.95	2	vB; vL; E135°±; vglbM; r	12†
1158	2858			5 26 24.5	+2.738	1	104 9 58.5	2.85	1	B; iE; sbM * 10 & 11	2
1159	2862			5 26 31.0	-0.137	1	157 37 22.4	2.92	1	pB; S; R; gB	2
1160	2853	III. 590		5 26 39.9	-3.085	2	167 51 34.3	-2.99	2	vF; S; R; smbM	2
1161	2863		Δ. 211							Cl; Ri; 2nd of sev	1
1162	2874										2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of used. Obs.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and m.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1163	h. 2864	II.		h m s 5 26 46.8	s -0.550	3	159 57 22.7	-2.91	3	F; pL; iR; vgbM; 2nd of group!	3+
1164	2865			5 26 50.2	-0.569	1	160 2 51.7	2.91	1	F; vL; vgbM; 3rd of group!	1+
1165	2866		Δ. 136?	5 27 15.3	-0.363	1::	158 57 7.3	2.89	1	vF; pL; R; 1st of 4!	1*
1166	358		M. 36	5 27 3.1	+3.966	3	55 57 28.2	2.76	3	Cl; B; vL; vRi; iC; st 9...11sc	9
1167		III. 747		5 27 11.2	+6.681	1	20 35 59.9	2.67	1	eF; pL; iF; mbM; er; * inv (? P.D.).	1*
1168	2867		Δ. 136?	5 27 32.7	-0.358	2::	158 55 26.2	2.86	2::	F; S; 2nd of 4!	2*+
1169	2861			5 27 17.2	+2.094	1	126 28 48.3	2.79	1	Cl; st 8...11	1
1170	2860	IV. 21		5 27 26.0	+2.536	1	112 2 39.9	2.77	1	F; vS; R; vsymbM*12; 3st inv.	2
1171	2868		Δ. 136	5 27 37.6	-0.361	4	158 56 0.7	2.84	4	{ Cl; pL; iF; 1st 9; } + group of 4 neb pB; R; psbM; 3rd of 4	3*+
1172	(456)			5 27 43.2	0.411	2:	159 12 33.1	2.83	2:	No description.....	2
1173	2870			5 27 44.2	0.132	1	157 34 9.4	2.82	1	Cl; Ri; 3rd of sev	1
1174	2872			5 27 46.7	0.553	1	159 57 36.7	2.82	1	F; S; 4th of gr of 7	1+
1175	2869			5 27 50.4	0.361	1::	158 56 34.5	2.84	1::	4th of 4	1*+
1176	2875			5 28 1.0	0.556	1	159 58 24.7	2.81	1	5th of gr of 7	1+
1177	2876			5 28 2.0	0.555	1	159 57 34.7	2.81	1	6th of gr of 7! D; a vS neb np	1+
1178	2877		Δ. 213	5 28 16.4	-0.129	1	157 32 34.9	2.77	1	Cl; L; irr.....	1
1179	360		{ M. 42= } 6 ¹ Orionis	5 28 24.0	+2.945	B.A.C.	95 29 10.9	2.68	B.A.C.	!!!; 6 ¹ Orionis & the great neb	Mon.*†
1180		V. 30	42, c ¹ Orionis	5 28 29.2	+2.958	B.A.C.	94 56 2.3	2.67	B.A.C.	!!; c ¹ 42 Orionis & neb.....	2
1181	2878		Δ. 238??	5 28 32.1	-0.019	3	156 20 18.8	2.74	3	vB; vL; iE; vgpmbM	3
1182		III. 240		5 28 32.9	+2.498	1	113 26 9.6	2.68	1	vF; vS; stellar.....	1
1183	361	V. 31	44, i Orionis	5 28 35.3	2.933	B.A.C.	96 0 21.0	2.66	B.A.C.	vF; vL; i 44 Orionis inv	3
1184	362			5 28 36.5	2.969	1	94 26 50.5	2.65	1	Cl; vB; iRi; stL, sc	1
1185		III. 1??	{ M. 43= } 144 Bo. Orionis	5 28 38.4	+2.948	...	95 21 48.7	2.65	...	{ i vB; vL; R, with tail; } mbM*8-9	Mon.*†
1186	2881			5 28 39.5	-0.393	1	159 5 49.5	2.75	1	Cl; vL; pRi; iF	1
1187	2882			5 28 39.8	-0.419	1	159 14 12.5	2.75	1	Cl; place of *	2
1188	359	III. 865		5 28 41.1	+3.897	1	58 6 38.4	2.62	1	eF; S; R; psbM	2
1189	2883			5 28 57.6	-0.581	4	160 4 57.4	2.72	4	B; pL; R; gbM	4
1190	2885			5 28 59.3	-0.739	2	160 50 53.1	2.73	2	F; L; iR; 3stp	2
1191			Chacornac	5 29 4.0	+3.581	...	68 52 20.4		...	!!!; variable (Chacornac)	0*
1192	2871			5 29 5.5	2.279	3	120 53 59.1	2.63	3	vF; S; R; lbM; st nr	3
1193	363	V. 34	ε Orionis	5 29 6.6	+3.042	1	91 17 44.7	2.61	1	!!!; eL; ε Orionis inv	2
1194	2884			5 29 12.9	-0.130	1	157 32 21.3	2.69	1	Cl; 4th of sev	1
1195	2873			5 29 13.9	+2.276	1	120 59 31.4	2.62	1	eeF; vS	1
1196		III. 269		5 29 20.4	+2.644	1	107 55 16.0	2.60	1	eF; vS; stellar.....	1*
1197	2887			5 29 21.2	-0.423	2	159 15 1.3	2.69	2	Cl; eS; st 11...16	2
1198	2879			5 29 24.4	+1.559	1	138 46 56.4	2.62	1	eeF; R; bM; diffie; p of 2...	1
1199		VIII. 42		5 29 32.9	3.711	2	64 15 45.5	2.55	2	Cl; L; iC; lRi.....	2
1200	2886			5 29 35.3	0.437	1	153 18 40.5	2.65	1	eF; eS; R	1
1201	2880			5 29 37.1	1.559	2	138 47 31.0	2.60	2	vF; R; gbM; st s; f of 2	2
1202		IV. 33		5 29 37.6	+2.914	3	96 48 42.9	2.57	4	B* inv in N.....	4
1203	2889			5 29 45.0	-0.997	1	161 58 34.9	2.67	1	F; pL; R; vlbM	1
1204	2888		Δ. 178??	5 29 51.1	0.350	2	158 50 55.5	2.65	2	Cl; st 13m	2
1205	2890		Δ. 214?	5 30 25.2	-0.048	2	156 59 13.3	2.59	2	vB; S; R; * + neb in vLCl..	2
1206	2981			5 30 49.7	+0.014	1	156 34 8.5	2.55	1	B; S; stellar; r	1
1207	2893		Δ. 215	5 30 54.5	-0.110	6	157 23 18.5	2.55	6	⊕; B; pL; pRi; C; st 12	6
1208	(509)			5 31 22.8	0.541	1::	159 51 17.4	2.52	1::	No description.....	1
1209	2895			5 31 26.5	-0.061	2	157 4 8.3	2.49	2	Cl; eL; vRi; vBvSNM	2
1210	2892			5 31 36.8	+1.433	1	141 1 12.1	2.43	1	eF; pL; R	1
1211	2894			5 31 44.8	+1.431	1	141 2 52.1	2.43	1	eF; pL; R; vlbM	1
1212	2897			5 31 59.0	-0.434	4	159 16 53.2	2.46	4	pF; pS; R; glbM; in Cl.....	4
1213	2898			5 32 15.9	0.764	1	160 55 58.8	2.44	1	F; eL; R; vglbM	1
1214	2899			5 32 40.9	0.149	1	157 37 1.3	2.39	1	vB; S; R; psmbM	1
1215	2907			5 32 41.8	-4.411	1	169 57 24.4	-2.52	1	vF; S; iE; bM; 2 st 9nf	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s						
1216	364			5 32 43.1	+6.02	1	24 16 50.7	-2.21	1	Cl; vLri; st 11.....	1
1217	2900			5 32 48.0	-0.177	3	157 47 24.3	2.39	3	Cl; pL; pC; iF; st 9...15 ...	3
1218	2901			5 32 52.0	0.448	1	159 21 16.3	2.39	1	Cl; vL; Ri; vIC	1
1219	2902			5 32 57.5	-0.583	1	160 3 5.6	2.38	1	F; vL; iR; gbm	1
1220	2896			5 33 7.7	+2.643	1	107 55 45.9	2.27	1	Cl of Lst	1
1221	2904			5 33 16.8	-0.819	1	161 10 19.2	2.36	1	pB; pL; R; pglbm; *10pinv	1
1222	2905		Δ. 98	5 33 29.3	0.624	2	160 15 3.1	2.33	2	B; pL; gbm	2
1223	2903		Δ. 218	5 33 31.3	0.180	2	157 48 16.7	2.31	2	F; vL; vLE; vglbm	2
1224	2906			5 33 46.9	-0.139	1	157 32 58.0	2.30	1	vF; S; R; in pLCl	1
1225	365	IV. 34		5 34 26.4	+3.283	2	80 59 3.8	2.14	2	○; pB; vS; vLE; r?	4+
1226		IV. 24		5 34 40.0	3.019	1	92 18 43.1	2.13	1	B* in M of L, lE neb	1*
1227		V. 28		5 34 47.2	+3.028	2	91 55 43.7	2.11	2	lirr; B; vvL; black sp incl ...	2
1228	2909			5 34 47.8	-0.964	1	161 47 23.1	2.23	1	vB; vS; lE; gmbM; r	1
1229		VIII. 28		5 34 48.5	+3.557	1	69 57 43.0	2.10	1	Cl; lRi; lC; st pL	1
1230	2908		Δ. 241	5 35 4.3	-0.058	3	157 0 29.6	2.18	3	Cl; vL; Ri; st 9...11	3
1231	2912		Δ. 100?	5 35 11.2	0.585	1	160 2 26.6	2.18	1	vF	1
1232	2911		Δ. 240	5 35 20.3	-0.158	2	157 38 43.5	2.15	2	pB; pL; R; gbm; in cLCl...	2
1233	2910			5 35 26.6	+0.072	2	156 6 50.8	2.14	2	pB; L; iR; gbm; 1st of 3 ...	2+
1234	2915			5 35 33.9	-0.804	2	161 5 22.5	2.15	2	⊕; B; pL; R; gbm; rr	2
1235	2913		Δ. 219?	5 35 39.8	0.162	5	157 39 47.1	2.13	5	B; L; E; 2nd of 3	5+
1236	(579)			5 35 45.4	0.553	1::	159 52 28.8	2.14	1::	Cl; no description	1
1237	2914			5 35 51.8	0.055	1	156 58 42.7	2.11	1	Cl; vL; Ri	1
1238	2916		Δ. 220	5 35 52.8	0.163	3	157 40 20.7	2.11	3	B; L; R; bM; 3rd of 3	3+
1239	2917			5 35 58.0	0.416	1	159 8 38.7	2.11	1	vF; pL; R; gbm	1
1240	(593)			5 36 18.4	0.553	1::	159 51 30.6	2.08	1::	Cl; no description	1
1241	2920			5 36 23.5	-0.710	3	160 38 22.6	2.08	3	pB; S; R; gbm; *9, np 5'...	3
1242	366			5 36 25.7	+3.273	1	81 25 37.9	1.97	1	Cl; vL; lRi; lC	1
1243	2918			5 36 27.9	-0.160	5	157 39 3.2	2.06	5	F; L; iR; gbm; r	5+
1244	2919			5 36 35.9	0.070	4	157 4 23.8	2.04	4	B; S; R; vglbm	4
1245	2922			5 37 1.6	0.391	1	159 0 17.4	2.02	1	Cl; vL; Ri; st 12...15	1
1246	(608)			5 37 7.7	-0.444	1::	159 17 32.7	2.01	1::	Cl; no description	1
1247	367		Lal. 10842	5 37 8.3	+3.375	1	77 10 43.0	1.90	1	*8, 9, with Fneb	1
1248	2923			5 37 11.1	-0.644	1	160 18 59.7	2.01	1	vF; R; gbm; 1st of 7	1+
1249	2925			5 37 27.0	0.635	1	160 16 15.3	1.99	1	F; S; lE; 2nd of 7	1+
1250	2926			5 37 35.8	-0.522	1	159 41 40.9	1.97	1	vF; L; pmE	1
1251	2921			5 37 53.2	+2.300	2	120 8 42.2	1.86	2	vF; S; R; bM	2
1252	2928			5 37 54.6	-0.479	1	159 28 11.8	1.94	1	Cl+neb; mC; iF; st vS	1
1253	2930			5 38 2.8	0.810	1	161 5 21.8	1.94	1	pB; S; R; gbm	1
1254	2929			5 38 3.3	0.557	1:	159 52 11.8	1.94	1	eF; vvS; vglbm	1
1255	2927			5 38 3.9	0.139	1	157 30 8.4	1.92	1	F; pL; lE; gbm	1
1256	2931			5 38 14.7	0.485	1	159 29 59.7	1.91	1	Cl; vL; Ri; st 10...15	1
1257	2932			5 38 17.9	0.755	1	160 45 15.4	1.92	1	pB; R; bM; p of 2; *9 bet...	1
1258	2935			5 38 24.3	0.652	1	160 20 40.7	1.91	2	pF; S; R; gbm; 4th of 7 ...	2+
1259	2933		Δ. 102	5 38 24.6	0.631	2	160 14 33.7	1.91	2	vB; pL; R; gbm; 3rd of 7...	2+
1260	2936			5 38 27.3	0.624	1::	160 12 26.7	1.91	1	vF; 5th of 7	2+
1261	(642)			5 38 42.9	-0.438	1::	159 14 36.6	1.88	1::	neb; no description	1
1262	2924			5 38 49.3	+2.174	1	124 1 5.3	1.79	1	Cl; L; lC; st 13	1
1263	2937			5 38 57.3	-0.055	1	156 56 50.8	1.84	1	vF; pS; E; gbm; 2st 10, s...	1
1264		VIII. 2		5 39 6.6	+3.276	1	81 16 35.1	1.73	2	Cl; poor; S sc st	3
1265	2938		Δ. 103?	5 39 10.9	-0.646	2	160 18 26.8	1.84	2	B; R; 6th of 7	2+
1266	2939			5 39 21.0	-0.634	1::	160 15 5.4	1.82	1::	vF; S; E; 7th of 7	1+
1267	368		M. 78	5 39 34.1	+3.072	2	90 0 15.7	1.61	2	B; L; wisp-sh; vgmbN; 3st inv; r	8+
1268	2940		Δ. 143	5 39 37.8	-0.409	1	159 4 43.3	1.79	1	F; L; E	1
1269	2941		Δ. 142	5 39 40.7	-0.423	8	159 10 18.3	1.79	8	!!!vB; vL; looped	8+
1270		IV. 36		5 39 49.8	+3.077	3	89 46 49.6	1.68	3	* with vF, lchev	3
1271	2934	III. 241		5 39 59.5	+2.532	1	112 3 59.6	1.68	1	eF; vS; R; gbm	2
1272	2942			5 40 1.0	-0.499	1	159 33 35.2	1.76	1	pB; pL; mE; 5st inv	1
1273	2943			5 40 9.0	-0.740	1	160 45 20.2	1.76	1	B; R; bM; rr; f of 2	1
1274		III. 267		5 40 35.0	+2.670	1	106 48 12.4	1.62	1	vF; pS; iE; bM	1
1275	2947			5 40 49.6	-0.534	1	159 43 48.3	-1.69	1	F; R; p of D neb	1

No. of Cata- logue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. II.'s Catalogues of Nebulæ.	Sir W. II.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s						
1276	2948			5 40 53.3	-0.550	1	159 48 52.3	-1.69	1	neb; np of gr of 4	1†
1277	2949		Δ. 152??	5 40 54.2	0.556	1	159 50 42.3	1.69	1	neb; sp of gr of 4	1†
1278	2950			5 40 56.7	0.532	1	159 43 22.6	1.68	1	B; R; f of D neb	1†
1279	2951			5 41 10.5	-0.484	1	159 28 20.2	1.66	1	Cl; vF; mC; st+neb	1†
1280	2945			5 41 11.3	+0.297	3	154 21 38.1	1.63	3	pF; L; R; glbM	3
1281	2952			5 41 11.8	-0.549	1:	159 48 24.2	1.66	1:	neb; nf of gr of 7	1†
1282	2953			5 41 15.6	0.554	1:	159 49 51.5	1.65	1:	neb; sf of gr of 7	1†
1283	2954			5 41 18.8	0.537	1:	159 44 57.5	1.65	1:	vF; R; *10vnr	1†
1284	2956			5 41 33.6	-0.536	1:	159 44 30.1	1.63	1:	B; pS; R; lbM; *10, p	1
1285	2946			5 41 38.9	+1.127	1	145 35 32.9	1.57	1	eF; pS; R; vlbM	1
1286	2955			5 41 39.6	-0.314	2	158 31 43.4	1.62	2	vF; S; R	2
1287		III. 270		5 41 43.7	+2.648	1	107 39 20.4	1.52	1	vF; eS; stellar	1*
1288	2944		Δ. 594	5 41 56.8	+2.163	2	124 18 21.4	1.52	2	⊕; B; pL; iR; gbM	2
1289	2957			5 42 5.1	-0.495	1	159 31 7.6	1.58	1	vF; S; mE; glbM; ?D	1
1290	2962			5 42 38.5	0.450	1	159 16 34.1	1.53	1	vF; pL; R; rr	1
1291	2963		Δ. 184??	5 42 43.0	0.397	1	158 59 18.4	1.52	1	vF; S; R	1
1292	2959			5 42 45.6	0.298	1	158 25 39.4	1.52	2	vF; S; R	1
1293	2961			5 42 47.9	0.128	2	157 23 26.7	1.51	2	Cl; F; eS; irr	2
1294	(725)			5 43 2.4	-0.319	1::	158 32 44.3	1.49	1::	neb; no description	1
1295	369		M. 37	5 43 7.5	+3.922	3	57 29 38.3	1.49	3	Cl; Ri; pCM; st L & S	8
1296	2960			5 43 6.1	+0.474	3	152 50 34.2	1.46	3	vF; pS; iR; psbm*16	3
1297	2965		Δ. 185??	5 43 7.3	-0.282	1	158 20 10.3	1.49	2	⊕; B; S; rr	2
1298	2966		{ Δ. 147? 151? 154? }	5 43 9.6	-0.450	5	159 16 26.3	1.49	5	⊕; B; pL; irrR; rr	5
1299	2958			5 43 9.8	+1.358	1	142 8 17.1	1.43	1	eF; pS; R; 3st10 sf	1
1300	(730)			5 43 24.9	-0.503	1::	159 33 29.2	1.46	1::	neb; no description	1
1301	2968			5 43 44.7	-0.888	1	161 23 47.8	1.44	1	pB; L; pmE; gbM*13	1
1302	2964			5 43 49.9	+1.390	1	141 36 16.9	1.37	1	pB; pS; R; glbM	1
1303	2969			5 44 23.6	-0.064	1	156 58 0.9	1.37	2	F; pS; R; gbM	2
1304	2967			5 44 49.2	+2.544	1	111 36 16.5	1.25	1	vF; S; vIE; gbM	1
1305	2971			5 44 54.6	-0.736	2	160 42 13.8	1.34	2	pB; pS; R; gbM	2
1306	2970		Δ. 153?	5 44 58.2	0.444	1	159 14 0.1	1.33	1	eF; pL; IE	1
1307	2972			5 45 6.7	-0.331	3	158 35 56.7	1.31	3	F; pS; R; vglbM	3
1308	370	{ III. 448 = III. 510 }		5 45 30.3	+2.897	(4)	97 30 3.2	1.16	1	eF; eS; IE; psbm; er	5
1309	2973			5 46 27.5	-0.817	2	161 2 46.7	1.21	2	vF; S; R; gbM	2
1310	371	VII. 24		5 46 37.1	+3.080	1	89 38 56.6	1.08	1	Cl; pL; lRi; pC; stS	3
1311	2975			5 46 40.1	-0.560	4	159 49 44.6	1.18	5	Cl; F; S; iF; vIC; rr	5
1312	2974			5 46 44.7	-0.246	1	158 5 35.9	1.17	1	eF; pL; iR	1
1313	2976			5 47 56.7	+1.446	1	140 37 4.7	1.01	1	eeF; vS; 3st10 sp	1
1314	2977			5 47 57.0	-0.326	1	158 33 11.2	1.06	1	F; S; R; *11p	1
1315	2978			5 48 7.4	0.149	5	157 29 33.8	1.04	5	F; pL; iR; vlbM; rrr	5
1316	2979			5 48 39.2	-0.438	2	159 10 22.7	1.01	2	⊕; vB; vS; vsbmM; rr	2
1317	2980			5 49 41.3	+0.369	2	153 42 41.3	0.89	2	cF; pL; R; vglbM	2
1318	2982			5 50 15.5	-0.922	1	161 30 44.6	0.88	1	vF; cL; vgbM	1
1319	2981		Δ. 106	5 50 16.8	-0.619	5	160 6 17.6	0.88	5	Cl; pB; iF; gvmCM; st15 ...	5
1320	2983			5 51 17.8	+0.162	3	155 20 50.2	0.76	3	pB; vS; R; gbM	3
1321		III. 225		5 51 46.8	+2.584	1	110 3 13.8	0.64	1	eeF; pS; E; r	1
1322	2985			5 51 59.6	-0.505	1	159 31 16.4	0.72	1	vF; pS; R; gbM	1
1323		VIII. 68		5 52 7.3	+4.659	1	40 6 11.5	0.55	1	Cl, not Ri; 1*7m	1
1324	2986			5 52 28.5	-0.481	3	159 23 33.9	0.67	3	pB; vS; R; gmbM	3
1325	372	VIII. 26		5 52 34.6	+3.647	1	66 42 23.8	0.54	1	Cl; pL; 40 or 50 st 8...15 ...	2
1326	2987			5 52 41.1	-0.131	7	157 21 35.8	0.64	7	F; pS; R; glbM	7
1327	2984			5 53 11.9	+2.401	1	116 39 52.1	0.53	1	vF; pS; R; gbM	1
1328	2988			5 53 22.1	+0.765	1	149 55 53.5	0.55	1	Cl; vIC; st L & S	1
1329	2989			5 53 26.1	-0.854	2	161 12 17.0	0.60	2	F; pL; R; gpmbM	2
1330	2991			5 53 53.9	0.837	4	161 7 52.5	0.55	4	⊕; B; pL; R; gmbM; r ...	4
1331	2990			5 53 54.3	0.146	5	157 27 0.8	0.54	5	F; pS; R; r; am st	5
1332	2992		Δ. 160	5 54 7.4	-0.504	4	159 30 51.1	-0.53	4	⊕; pB; R; gmbM; rr; st 14...16.	4

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
1333	2994			5 54 23.3	-0.503	2	159 30 24.0	-0.50	2	vF; S; R; f of 2	2
1334	2993			5 54 34.6	+0.090	1	155 51 30.9	0.47	1	eF; S; R	1
1335		II. 264		5 54 55.9	+2.482	1	113 49 28.9	0.37	1	F; S	1
1336	2995			5 55 1.4	-0.342	3	158 37 0.8	0.44	3	pF; pS; iR; bM	3
1337	373			5 55 15.5	+2.822	1	100 36 14.8	0.34	1	* (3 Monoc) inv in pL, F, n	1
1338	374			5 55 39.0	+3.206	1	84 16 44.3	0.29	1	Cl; L; pRi; vLC; st10,	1
1339	3009			5 56 0.0	-6.638	2	172 9 27.8	0.54	2	F; pS; iR; bM	2
1340	2998			5 56 13.4	0.798	1	160 56 36.5	0.35	1	F; lE; r	1
1341	2997			5 56 22.5	-0.274	2	158 13 12.4	0.32	2	eF; S; R; bM	2
1342	2996			5 56 50.7	+0.836	1	149 7 34.5	0.25	1	eF; S; R; *12 vnr	1
1343	3000			5 57 0.8	-0.519	1	159 34 54.6	0.28	1	F; vS; R; vsmbM; stellar	1
1344	3001			5 57 13.7	-0.416	1	159 2 0.5	0.25	1	F; pS; R; bM	1
1345	2999			5 57 34.0	+1.437	2	140 44 5.6	0.18	2	eeF; R; *15 att	2
1346	3002			5 57 46.1	+0.010	1	156 24 55.3	0.19	1	eeF; lE; *16 att	1
1347	3003			5 57 52.5	-0.119	4	157 16 2.3	0.19	4	F; pL; R; vglbM	4
1348	3004			5 58 9.8	+0.141	3	155 28 41.2	0.16	3	F; pL; R; vglbM	3
1349	3005		Δ. 196	5 58 32.0	-0.316	5	158 28 3.8	0.14	5	pB; S; R; gbM; 1st of 3	5
1350	3006		Δ. 161?	5 58 35.9	-0.447	4	159 11 59.1	-0.13	4	⊕; vB; S; R; vgvmbM; rr.	4
1351	375	VI. 17		5 58 49.3	+3.670	1	65 53 46.0	0.00	1	Cl; pS; mC; vRi; nrΔ; steS.	3
1352	3007		Δ. 193	5 58 50.0	-0.345	4	158 38 5.4	-0.12	4	pF; S; R; gbM; *15 att nf.	4
1353	3008			5 58 50.7	0.287	1	158 17 48.7	0.11	1	pF; pS; R; gbM	1
1354	3013			5 59 33.7	-1.702	1	164 21 22.3	0.09	1	F; pL; R; gpmbM	1
1355	3010			5 59 36.3	+0.365	3	153 43 19.4	0.02	3	F; pL; R; vglbM	3
1356	3011		Δ 194	5 59 39.1	-0.325	5	158 31 0.8	-0.04	5	⊕; vB; R; mCM; rr	5
1357	376			5 59 59.5	+4.765	1	38 17 55.2	+0.14	1	Cl; pL; poor; st11	1
1358	3012		Δ. 223?	6 0 4.7	-0.229	1	157 56 54.0	0.00	1	F; S; R; gbM	1
1359	378	IV. 44		6 0 8.7	+2.927	1	96 11 46.0	+0.10	1	Nebulous *7; am 3 st	2
1360	377		M. 35	6 0 12.5	3.677	1	65 39 16.9	0.13	1	Cl; vL; cRi; pC; st 9...16...	8
1361	379	VIII. 24		6 0 33.4	3.405	1	76 1 38.5	0.15	1	Cl; S; lRi; pmC; *Σ. 848...	3
1362		IV. 19		6 0 44.0	+2.923	3	96 22 39.5	0.15	3	*9 in vF, pLneb; E 170°	3
1363	3016			6 0 44.6	-0.750	1	160 43 11.2	0.04	1	eF; L; R; glbM	1
1364	3015			6 0 52.1	0.348	2	158 38 48.1	0.07	2	F; cL; R; lbM	2
1365	3018			6 0 56.3	-1.295	1	162 58 41.2	0.04	1	pF; pL; R; gmbM	1
1366			Auw. N. 21	6 1 19.1	+3.569	...	69 29 42.5	0.01	...	*8m in neb (Bruhns)	0
1367	3017			6 1 27.2	-0.056	1	156 51 19.9	0.13	1	eeF; pL; R; gbM	1
1368	3020			6 1 42.0	-0.192	2	157 43 40.5	0.15	2	F; vS; iR; lbM; r	2
1369	3019			6 1 56.1	+0.359	1	153 45 54.4	0.18	1	eF; vS; R	1
1370	3014			6 2 6.2	2.539	1	111 43 53.8	0.26	1	F; pS; vmE; glbM	1
1371	380	VIII. 6		6 2 12.0	3.182	1	85 15 56.7	0.29	1	Cl; pRi; lC; st L & S	4
1372	3021			6 2 28.9	0.171	1	155 15 14.6	0.22	1	vF; S; R	1
1373	381	IV. 38		6 2 41.6	2.924	1	96 18 57.6	0.32	1	pB *; L*neb; E 90° ±	3
1374	382			6 3 59.9	2.990	1	93 30 5.2	0.44	1	Cl; L; vLC	1
1375	383	IV. 20		6 4 17.9	2.927	2	96 12 6.1	0.47	2	*11&4 S st in vF, L neb	5
1376	384	VII. 25		6 4 39.8	+3.200	1	84 31 53.0	0.50	1	Cl; pL; pRi; pC; st L & S	2
1377	3025			6 5 3.8	-0.516	3	159 33 49.6	0.42	3	{ pB; pS; R; gbM } D neb; { vF; R; glbM } 12° 5' } 3	3
1378	3022			6 5 5.6	+2.168	1	124 4 30.3	0.51	1	pF; pL; vmE; gvlbM	1
5064				6 5 7		...	88 50 39		...	See No. 5064.	1
1379	3027			6 5 7.1	-1.817	1	164 42 26.7	0.39	1	vF; pL; R; glbM	1
1380	3023			6 5 16.5	+1.331	1	142 29 20.0	0.50	1	pB; vS; E; vsbM; *9 p 5...	1
1381		VII. 57		6 5 25.2	4.189	1	50 6 21.0	0.60	1	Cl; cL; C; iF; st vS	1
1382	3026			6 5 47.6	0.196	1	155 4 9.3	0.51	1	F; iF; glbM; 2 or 3 st inv	1
1383		VI. 5		6 6 2.3	3.377	2	77 9 55.9	0.63	2	Cl; L; Ri; gvmCM	2
1384	3024	II. 265		6 6 14.4	+2.539	4	111 46 26.6	0.62	4	pF; pS; vLE; pmbM; st nr...	5
1385	3028			6 6 21.3	-0.087	2	157 4 22.5	+0.55	2	vF; pS; R; gbM	2
5065				6 6 40.9		...	88 58 11.2		...	See 5065.	1
1386	3031			6 7 57.9	-1.404	1	163 22 10.8	+0.66	1	F; vS; R; bM	1
1387	3029			6 9 5.0	+1.799	2	133 37 23.5	0.85	2	eF; pS; R; vlbM; ?134° PD.	2
1388	3030			6 9 18.6	1.798	2	133 39 27.1	0.87	2	eF; S; R; psbM; ?134° PD.	2
1389	385		Σ. 885	6 9 23.0	+3.213	1	83 58 10.6	0.92	1	* Chief of Cl	1
1390	3035			6 9 26.6	-2.061	1	165 24 21.8	+0.76	1	pB; pL; iR; vgpmbM; r	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	IL		h m s	s						
1391		VII. 13		6 9 33.3	+2.622	1	108 36 47.3	+0.91	1	Cl; L; pRi; IC	1
1392	3034			6 10 13.2	0.506	1::	152 29 50.0	0.90	1	pF; S; R; bM	1
1393	3032			6 10 22.8	2.551	1	111 19 46.4	0.98	1	pB; pL; mE, 87°; psbMRN	1
1394	3033			6 10 23.9	+2.399	2	116 43 35.4	0.98	2	F; pS; vLE; psbM	3
1395	3037			6 12 5.0	-1.526	1	163 47 55.3	1.01	1	vF; cL; R; gvlbM	1
1396	3036			6 12 33.1	0.423	3	159 5 2.4	1.08	3	vB; pL; R; mbM; r	3
1397	3038			6 12 45.1	0.917	1	161 29 30.7	1.09	1	vF; S; l; glbM; ** p	1
1398	3039		Δ. 201	6 13 34.4	-0.270	2	158 13 12.4	1.18	2	B; pS; lE; gbM; rrr	2
1399	386	VII. 20		6 14 5.0	+2.902	2	97 14 22.3	1.31	2	Cl; cL; pRi; pC; st 11...15	5
1400	3040			6 15 36.9	2.532	1	112 0 58.2	1.44	1	vF; pL; R; vglbM	1
1401	3041			6 16 4.8	2.387	2	117 10 38.1	1.47	2	vB; S; R; psmbM; r	2
1402			Anw. N. 22	6 16 25.2	3.539	...	70 35 28.2	1.54	...	F Cl (Markree Obs. Jan. 13, 1853).	0
1403	387			6 16 28.9	2.964	1	94 37 19.9	1.53	1	Cl; P; vIC; st 6, 11...12	1
1404	3042			6 17 5.4	1.753	1	134 41 54.5	1.55	1	Cl; B; P; st 8, ...	1
1405	3044			6 18 1.5	0.980	1	147 29 35.0	1.60	1	vF; lE; vgbM; p of 2	1
1406	3045			6 18 1.9	0.983	1	147 27 30.0	1.60	1	vF; lE; vglbM; f of 2	1
1407	3043			6 18 43.9	2.513	3	112 46 8.0	1.70	3	F; pL; R; vglbM; 2st inv...	3
1408		VII. 35		6 19 44.5	3.372	1	77 16 45.6	1.82	1	Cl; pC; with neb?	1
1409	388	VII. 26		6 19 57.6	2.847	1	99 34 30.9	1.83	1	Cl; P; lCM; st 12...15	2
1410	3046			6 20 1.4	2.536	1	111 55 24.9	1.83	1	eF; R; * p 270°, 90''	1
1411	3047			6 20 33.0	0.293	1	154 23 21.3	1.81	1	F; S; R; glbM	1
1412	3048			6 20 49.4	0.231	1	154 52 40.6	1.82	2	eF; vS; R; 1st of 3	2
1413	3049			6 20 55.3	+0.226	1	154 55 25.2	1.84	1	eF; S; lE; 2nd of 3	1
1414	3050			6 21 0.8	-0.136	3	157 27 7.2	1.84	3	F; pL; R; gvlbM; * f	3
1415		VIII. 25		6 21 1.6	+2.963	1	94 40 44.6	1.92	1	B* (10 Monoc) + Cl	1
1416	3051			6 21 11.0	0.220	1	154 58 9.5	1.85	1	eF; S; 3rd of 3	1
1417	389	VIII. 9		6 21 13.5	3.473	(1)	73 13 33.5	1.95	1	Cl; eL; pRi; lC; st L & S...	2
1418	3052			6 21 49.5	0.235	1	154 51 45.3	1.91	1	vF; S; R; * 12 nr	1
1419	390	VII. 5		6 22 11.0	3.233	1	83 4 23.2	2.04	1	Cl; pRi; pC; st 10, 12...15	2+
1420	392			6 23 28.8	3.189	1	84 57 32.2	2.14	1	* 8 in L; P; BCl	4
1421	391	VIII. 49		6 23 31.7	+4.013	2	54 42 17.1	2.17	2	Cl; pL; P; vIC; st 7, 10...15	3
1422	3054			6 23 42.4	-0.366	3	158 50 33.8	2.06	3	vF; pL; R; glbM	3
1423	3053		Δ. 616?	6 24 13.8	+2.267	2	121 11 28.4	2.18	2	pB; cL; R; vglbM; 4''	2
1424		VII. 2	12 Monoc. B.A.C.	6 24 53.4	3.189	B.A.C.	85 2 14.5	2.27	B.A.C.	Cl; beautiful; st sc	
1425	393	IV. 3		6 24 58.0	3.312	2	79 44 43.1	2.27	2	pL; com; mbNsf alm*; * 7.8 nf.	7*+
1426			Anw. N. 23	6 25 51.9	+3.729	...	63 35 16.9	2.36	...	Small cluster (Markree Obs. Dec. 23, 1853).	0
1427	3055			6 26 38.1	-0.360	4	158 50 0.3	2.31	4	pB; pL; R; vgbM; * p	4
1428	394			6 27 5.0	+2.956	1	94 57 53.2	2.44	1	Cl; pRi; lC; iF; st 8, 12...14	1
1429	395	VIII. 3		6 27 8.4	3.269	(2)	81 32 16.8	2.46	(2)	Cl; vL; E; Ri; lC	4
1430	396	VIII. 50		6 27 23.7	3.199	(1)	84 32 14.7	2.49	1::	Cl; vL; pRi; lC; st S	3
1431		VII. 54		6 27 56.0	6.043	1	24 2 15.6	2.62	1	vF; st eS	1
1432	397	VII. 22		6 28 25.1	3.253	1	82 13 44.4	2.58	1	Cl; S; pC; iF; st 11...15	2
1433	3056			6 28 59.1	2.153	2	124 42 52.7	2.59	2	eF; S; lE; vlbM	2
1434	3057			6 29 25.9	0.325	3	154 13 23.4	2.58	3	F; cL; R; vglbM; r; 17.0d	3
1435		VI. 28		6 30 47.9	3.329	1	79 0 33.4	2.78	1	Cl; cRi; eC; iF; st eS	1
1436	398	VIII. 48		6 31 2.3	3.041	1	91 20 42.7	2.79	1	Cl; vL; P; vIC; st L & S	2
1437	399	IV. 2		6 31 31.4	3.278	3	81 8 20.5	2.85	3	B; vmE 330°; Ncom = * 11	7+
1438	400	VII. 37		6 32 24.0	3.101	1	88 43 47.6	2.92	1	Cl; vC; iR; bM; st eS	2
1439	3058			6 32 41.8	2.462	1	114 43 55.6	2.92	1	pF; lE; bet 2 vS st; psbM...	1
1440	401	V. 27	15 Monoc.	6 33 16.0	3.305	1	79 59 24.7	2.99	1	15 Monoc; Cl; *; ? neb	6*
1441	402	VIII. 5		6 33 40.5	3.354	1	77 57 3.2	3.04	1	Cl; P; 30 or 40 st 12...13	1
1442	403	VI. 21		6 34 32.6	3.748	1	62 53 35.6	3.12	1	Cl; pS; eC; Ri; st 11...15	2
1443	3059			6 35 31.5	2.236	1	122 20 49.8	3.16	1	pB; S; R; 2 or 3 st v nr	1
1444	404	VI. 3		6 36 26.7	3.180	(1)	85 17 41.1	3.27	1	Cl; vmC; not Ri; st vS	2
1445	405	VII. 36		6 36 35.5	3.154	1	86 24 51.4	3.28	1	Cl; lC; not Ri	2
1446	3060			6 37 3.1	2.503	3	113 20 15.0	3.30	3	pF; S; R; gbM; amst	3
1447	3061			6 37 7.0	+2.390	3	117 19 37.3	+3.31	3	pF; pS; vLE; bM; r	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
1448	406	II. 615		6 38 2.9	+3.950	2	56 17 42.9	+3.43	2	F; S; bM.....	4
1449	407	II. 614		6 38 2.9	3.951	2	56 15 56.9	3.43	2	eF; vS.....	4
1450	3062			6 39 16.6	2.386	1	117 30 7.7	3.49	1	pF; pL; lE; gbM	1
1451		VIII. 71		6 39 30.5	4.223	1	48 47 17.8	3.56	1	Cl; pRi; vLC; st pL	1
1452		III. 271		6 39 46.2	2.642	1	108 3 23.8	3.56	1	3 or 4 S st + neb.....	1*
1453	408	VIII. 31		6 40 39.4	3.002	1	93 1 23.9	3.63	1	Cl; L; C; ab 100 st 9...15...	3
1454	411		M. 41	6 41 0.3	2.578	1:	110 36 2.2	3.64	1	vL; B; IC; st 8,...	3*
1455	410, a		R. nova	6 41 34.1	3.944	...	56 23 51.2	3.74	...	No desc.; β of Lord R.'s diag.	0
1456	410, b		R. nova	6 41 35.9	3.944	...	56 22 58.2	3.74	...	No desc.; γ of Lord R.'s diag.	0
1457	410	III. 898		6 41 39.3	3.944	1:	56 25 16.2	3.74	1	eF; vS.....	2
1458	409	III. 897		6 41 39.4	3.946	1:	56 21 16.2	3.74	1	eF; vS.....	2
1459	3063			6 41 44.2	2.414	1	116 35 49.0	3.70	1	{pB; R; gbM} D neb; am st	1
1460	410, c		R. nova	6 41 54.0	3.944	...	56 19 55.2	3.74	...	No desc.; ϵ of Lord R.'s diag.	0*
1461	3064			6 42 3.3	2.414	1	116 34 19.9	3.73	1	eF; S; R; bet st; D neb p ...	1
1462	3066			6 43 23.6	0.428	1	153 34 16.4	3.78	1	vF; S; R; vglbM	1
1463	3065		Δ 578	6 44 1.7	2.124	4	125 50 56.4	3.88	4	$\oplus$; B; pL; iR; gbM; rr.....	4
1464	412			6 44 17.3	2.915	1	96 49 35.2	3.94	1	Cl of 30 or 40 st.....	1
1465	413	VI. 27		6 44 35.1	3.086	2	89 22 44.1	3.97	3	Cl; Ri; L; iF; st L & S	5
1466	414	VIII. 39		6 45 3.1	2.912	3	96 55 13.0	4.00	3	Cl; L; P; IC	6
1467	415	VI. 2		6 46 55.4	3.501	2	71 49 12.7	4.09	2	Cl; pL; Ri; mC; st vS	5+
1468	3067			6 47 45.6	0.375	1	154 6 43.8	4.16	1	vF; vS; R; 2 st Δ	1
1469	416	VIII. 51		6 47 47.1	2.910	1	97 1 26.2	4.24	1	Cl; P; vLC	4
1470	3068			6 47 58.3	0.369	2	154 9 52.4	4.18	2	vF; pS; vLE 90°	2
1471	417	VI. 18		6 49 15.4	2.911	3	97 1 24.8	4.36	3	Cl; pL; pRi; mC; st 13... ..	7
1472	3069			6 49 23.0	1.949	2	130 41 29.2	4.34	2	pB; pL; vME 44° 8; psbM..	2
1473	418	VIII. 60		6 50 51.9	2.971	1	94 24 15.0	4.50	1	Cl; IC; not Ri.....	2
1474	419			6 51 6.4	3.311	1	79 33 25.9	4.53	1	Cl; P	1
1475			D'Arrest, 50	6 51 23	2.89	[3]	97 45 24	4.51	[3]	F; vS; R	0
1476	420			6 51 49.9	4.665	2	39 12 51.9	4.63	2	eF	2
1477	421	II. 304		6 52 55.6	2.898	3	97 35 39.1	4.67	3	pF; S; R; r; S st inv	6+
1478	421, a		R. nova	6 52		...	97 35		...	Makes a close D neb with h. 421.	0
1479	{ 422 = 3070 }	VII. 14		6 53 4.4	2.759	2	103 30 44.4	4.68	2	Cl; L; Sc; st 8...9	3
1480	423	VIII. 1B		6 53 48.2	3.145	2	86 44 53.5	4.75	2	Cl of sc st; st 8, 9,...	3*
1481		III. 874		6 54 46.7	4.658	1	39 15 39.4	4.88	1	vF; vS; lE	1
1482	424	II. 861		6 54 57.4	4.661	1	39 13 10.3	4.81	1	pB; S; iR; gbM; *8, 120°...	2
1483	425		M. 50	6 56 12.5	2.886	4	98 8 46.5	4.95	4	Cl; vL; Ri; pC; E; st 12...16	8
1484	427	VII. 38		6 56 57.0	3.100	2	88 44 31.6	5.02	3	Cl; L; Ri; cC; st 12...16 ...	5
1485	3071			6 57 7.4	2.368	1	118 30 13.3	5.01	1	pB; pL; lE; gbM	1
1486	426	II. 734		6 57 28.6	4.663	1	39 5 50.6	5.02	1	vF; pL; iR; psmbM; st p ...	2
1487	428	IV. 25		6 57 33.2	2.817	1	101 6 47.8	5.06	1	pB* inv in S, vF, neb	3
1488	3072			6 58 6.8	1.912	2	131 51 55.4	5.08	2	vF; S; vLE; bM; am st	2
1489	429 {	II. 735 =		6 58 33.7	4.547	1	41 10 42.0	5.20	1	vF; vS; stellar.....	4
1490	432	III. 875		6 58 47.5	3.741	2	62 35 40.7	5.19	2	Cl; L; vLC; Sel inv.....	3
1491	430	VIII. 40		6 58 49.2	4.631	1	39 36 2.6	5.22	1	F; S; R; psbM	2
1492	430, a	II. 862	R. nova	6 58 ±		...	39 36 ±		...	Several near h. 430 (? 426, 433 & 1 nov).	0
1493	431	III. 899		6 58 54.3	3.986	(1)	54 39 36.3	5.21	1:	vF; S; R; bM.....	2
1494		VIII. 32		6 59 54.6	2.847	1	99 52 4.8	5.26	1	Cl; L; IC	1
1495	435			7 0 6.6	2.949	1	95 24 37.7	5.29	1	Cl; vLC	1
1496	434	II. 769		7 0 9.7	3.515	1	71 0 17.0	5.30	1	pB; pL; R; glbM	2
1497	433	II. 736		7 0 27.9	4.627	2	39 36 26.8	5.36	2	pF; S; R; glbM; r.....	4
1498		VIII. 33		7 1 35.7	2.834	1	100 26 14.0	5.40	1	Cl; cL; P; IC	1
1499	3073			7 1 52.0	2.775	1	102 57 4.9	5.43	1	Cl; pL; pRi; gbM; st 10...14	1
1500		IV. 65		7 2 14.0	3.055	2	90 30 8.1	5.47	1	*9 aff with S, vF, neb	1
1501		III. 746		7 2 47.5	+5.827	1	24 57 49.7	+5.59	1	vF; S; R; lbM	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1502	h. 3074	H.		h m s 7 3 7.1	+0.009	1	157 11 8.8	+5.46	1	Cl; P; IC; 30 st ±	1
1503	436	VII. 27	C. H.	7 3 16.3	2.881	1	98 23 59.5	5.55	1	Cl; cL; P; cC	3
1504	437			7 6 55.9	2.817	1	101 15 9.5	5.85	1	Cl; IC; * taken	1
1505		VII. 15		7 7 48.2	2.506	1	113 51 10.6	5.92	1	Cl; pRi; pC	1
1506		VIII. 34		7 7 50.5	2.845	1	100 3 44.9	5.93	1	Cl; L; IC; one vB*	1
1507	438	VII. 16		7 8 28.2	2.463	1	115 29 29.4	5.98	1	Cl; cRi; IC	2
1508	439	VI. 6		7 9 3.8	3.390	1	75 58 53.5	6.05	1	Cl; pS; pRi; mC; st 15...16	2*
1509		VII. 6		7 9 12.4	3.394	1	75 46 51.8	6.06	1	Cl; IC	1
1510		VIII. 45		7 10 31.8	2.687	2	106 47 59.8	6.16	2	Cl; P; IC	1
1511	3075	V. 21		7 11 3.0	2.778	1	102 57 54.0	6.20	1	ll; vF; vL; viF	3†
1512	{ 440 = 3076 441 = 3077 }	VII. 12	C. H.	7 11 23.4	2.721	2	105 23 19.9	6.23	2	Cl; vL; Ri; pC; st 9...12	5
1513	{ 442 = 3077 }	VII. 17	A.S.C. 905	7 12 54.6	2.488	2	114 42 14.2	6.34	2	Cl; pL; Ri	4
1514	442			7 13 59.7	2.908	1	97 18 9.5	6.05	1	Cl; pC; st pL; bifid	1
1515		III. 748		7 14 2.3	6.410	1	20 42 16.2	6.54	1	vB; pL; R; mbM; r; vS* inv.	1
1516		VIII. 27		7 14 8.9	2.568	1	111 40 18.5	6.45	1	Cl; S; P; IC	1
1517	443			7 14 20.5	2.845	1	100 7 38.4	6.48	1	Cl; S; pRi; st 15	1
1518	3078			7 15 0.5	0.679	2	152 5 45.1	6.47	2	pB; pL; iE; glbM	2
1519	444	II. 316		7 16 42.9	3.792	1	60 14 41.0	6.70	1	B; S; R; bMN; p of D neb, 45°, 60°	3†
1520	445	II. 317		7 16 44.7	3.792	1	60 14 21.0	6.70	1	pB; S; R; bMN; f of D neb	3†
1521	3080	VIII. 35		7 17 31.6	2.781	1	102 59 33.2	6.74	1	Cl; vL; pRi; IC; st L	5
1522	3079			7 18 18.9	2.423	1	157 15 54.7	6.79	1	pF; pS; R; vsmBM; am st	1
1523	3084			7 18 27.0	0.617	1	112 48 48.5	6.75	1	vF; vS; R; am st	1
1524	3082			7 18 32.8	2.427	1?	117 6 24.0	6.80	1:	pF; S; R; bM	1
1525	3081			7 18 39.7	2.597	1	110 39 57.6	6.82	1	Cl; pS; pmC; st 12	2
1526	3083			7 18 59.7	2.595	2	110 44 57.5	6.85	2	Cl; IC; bifid; *	2
1527	446			7 19 17.6	3.920	1?	55 54 27.3	6.91	1:	neb; 1st of 4	1*
1528	447	III. 703		7 19 23.1	3.921	1	55 53 27.3	6.91	1:	vF; vS; R; bM	3*
1529		II. 820		7 19 44.8	4.017	1	52 58 44.5	6.95	1	pB; S; stellar	1
1530	448	III. 900		7 19 45.1	3.920	1	55 55 8.2	6.95	1	vF; S; R; bM	2*
1531	449	III. 901		7 19 57.6	3.921	1	55 51 47.1	6.97	1	vF; S; R; psbM	2*
1532	450	IV. 45		7 20 54.4	3.557	1	68 48 33.2	7.04	1	B; S; R; *8 M	4†
1533		VIII. 44		7 21 2.5	3.232	1	82 41 1.2	7.04	1	Cl; L; P; vLC; st L	1*
1534		VIII. 11		7 21 11.7	3.384	1	75 56 51.5	7.05	1	Cl; pRi; C	1
1535	451	VIII. 36		7 21 38.5	+2.818	1?	101 27 32.1	7.07	1?	Cl; vL; vLC	2
1536	3085			7 21 42.1	-0.151	2	158 43 56.0	7.00	2	pB; cL; cE 117°, lbM	2
1537			Auw. N. 24	7 22 32.1	+3.070	::	89 55 49.5	7.15	...	Two B neb (Bond, Feb. { 1853).	0
1538			Auw. N. 25	7 22 32.1	3.070	::	89 55 49.5	7.15	...		0.
1539	454	VII. 65		7 22 57.2	2.767	1	103 41 23.1	7.17	1	Cl; S; cRi; cC; st vS	2
1540	453	III. 19		7 23 6.9	3.290	2	80 3 25.0	7.20	3	eF; S; R; lbM; * inv.	4
1541		V. 44		7 23 18.6	5.864	1	24 0 3.0	7.30	1	ll; cB; cL; vmE; vgmBMN 7'	2
1542	452			7 24 35.7	6.898	1	18 1 37.9	7.43	1	Cl; vLC	1
1543	3086			7 25 21.7	2.693	1	106 54 16.1	7.37	1	Cl; S but B; st 8...10	1
1544		VIII. 52		7 26 49.5	2.789	1	102 47 49.7	7.49	1	Cl; vL; P; vLC	1
1545	455	VIII. 37		7 26 49.5	2.735	1	105 8 46.7	7.49	1	Cl; P; IC; st 9, &c	3
1546	456	II. 821		7 27 43.4	3.954	1	54 28 22.7	7.59	2	pB; cS; R; vgvbM; r; almO	3
1547	3087			7 28 25.2	0.743	1	151 57 47.1	7.57	1	vF; L; R; gbM; r	1
1548	457	I. 218		7 28 37.8	4.077	1	50 48 47.4	7.68	1	pB; pL; lE 90°; vgbM; *7, 8, 19°	3
1549	458	VI. 1		7 30 5.7	3.566	1	68 7 10.4	7.78	1	Cl; cL; Ri; C; st 11...18	10
1550	3089	VII. 67		7 30 10.0	2.615	1	110 18 0.8	7.76	1	Cl; L; cRi; st 11...13	3
1551	{ 459 = 3088 }	VIII. 38		7 30 10.8	2.760	2	104 10 31.8	7.76	2	Cl; B; vL; pRi; st L & S	4
1552	3090	VII. 28		7 30 38.0	2.774	1	103 32 55.0	7.80	1	Cl; vL; Ri; pC; st vS	2
1553		VIII. 87		7 31 51.7	+2.751	1	104 35 14.0	+7.90	1	Cl; P; S; st vS	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1554	h. 460	H. 822		h m s	s	1	37° 20' 50.0	+8.00	1	eF; R; vgbM; r; *8 p	2
1555	3091			7 32 26.8	+4.676	1	137 18 28.9	7.93	1	eF; L; pmE; gmbM; 2 st inv	1
1556		VIII. 47		7 32 31.5	1.739	1	106 11 49.4	7.98	1	Cl; vL; vIC.....	1
1557		VIII. 46		7 32 55.5	2.715	1	106 1 50.0	8.00	1	Cl; vL; vIC.....	1
1558		III. 829		7 33 4.6	2.719	1	36 34 59.8	8.16	1	eF; vS; R; bM	1
1559	3092	VI. 36		7 34 26.7	4.715	1	108 45 38.6	8.12	1	Cl; pL; pC; E 0°; st L & S...	3
1560	462			7 34 41.6	2.656	1	80 24 50.1	8.17	1	eF; *15, 300° 0, 90"	1
1561	3096			7 35 4.7	+3.277	1	158 58 3.7	8.09	5	pB; S; R; pmbM; 3 st 11 n...	5
1562		II. 616		7 35 9.9	-0.117	5	57 59 59.0	8.20	1	F; S; lbM	1
1563	461			7 35 11.2	+3.833	1	37 35 56.6	8.22	1	vF; vS; R; bM	1
1564	463		M. 46	7 35 19.3	4.653	1	104 29 50.4	8.18	1	l; Cl; vB; vRi; vL; inv O...	4
1565	{ 464 = 3093 }	IV. 39		7 35 24.3	2.755	1	104 24 39.4	8.18	3	O; pB; pS; eIE; r; 3*75 d...	4†
1566	3094			7 35 25.4	2.757	2	121 19 45.1	8.17	1	Cl; B; pRi; pL; IC; st 9, 12...14.	1
1567	3095	IV. 64		7 35 26.2	2.328	1	107 53 22.3	8.21	1	O; cB; not v well def	3†
1568	3097			7 35 41.2	+2.677	1	159 12 49.3	8.21	4	{ cL; vF; R } D neb; 40°; { pL; vF; R } * inv M ...	4
1569				7 36 43.9	-0.149	4	35 3 14.3	8.41	1	F; am 4 st	1
1570	465			7 37 36.4	+4.803	1	113 32 43.2	8.44	1	Cl; L; pRi; IC; st 8...13 ...	2
1571	3098		M. 93	7 38 39.2	2.542	1	114 21 12.2	8.44	1	Cl of 18 or 20 st 11...13 ...	1
1572	466		Lal. 15134	7 38 41.4	2.522	1	127 38 15.8	8.56	1	Cl; vL; vIC; 1* 4.5 m	1
1573	3099			7 40 19.2	2.138	1	117 0 6.7	8.69	2	O; F; S; IE; am 60 st	2
1574	3100			7 41 47.4	2.457	2	116 54 21.0	8.70	1::	Cl; S; pRi; pC	1
1575	3101			7 41 56.8	2.459	1::	110 57 4.1	8.77	1	Cl; cL; pRi; IC; st 12	1
1576	3102			7 42 52.2	2.611	1	34 9 26.9	8.83	1	vF; R; vgbM	1
1577	467			7 42 56.7	4.838	1	80 5 29.5	8.91	2	vF; S; rr group + neb	3*
1578	468	III. 479		7 44 24.4	3.280	2	32 57 44.1	9.07	1	eF; R; p of 2	1
1579	469			7 45 47.6	+4.908	1	161 3 33.8	8.96	1	vF; S; R; lbM	1
1580	3104			7 46 20.9	-0.423	1			...	8 of 10 neb, in line with h. 469, 470.	0
1581	{ 469, a }		R. 8 novæ	7 47 ±		...	32 57 ±		...		0
1582						...			...		
1583						...			...		
1584						...			...		
1585						...			...		
1586						...			...		
1587						...			...		
1588						...			...		
1589	472	IV. 22		7 46 40.1	+2.488	1	116 2 1.1	9.07	1	pB; vL; R; er; *8 M.....	3
1590	470	III. 836		7 46 42.0	4.905	2	32 57 5.9	9.13	2	F; vS; R; *9 sf; f of 2	3
1591	471	III. 830		7 47 11.2	4.660	1	36 46 9.8	9.16	1	F; pS; E?; bM vS*? L* nf	2
1592	471, a		R. nova	7 47		...	36 46		...	Makes D neb with h. 871 ...	0
1593	3103		Δ. 535	7 47 18.9	2.133	2	128 11 8.3	9.11	2	l; Cl; B; Ri; L; IC; st 12 ...	3
1594			M. 47	7 48 20.5	2.751	W.	105 3 19.3	9.21	W.	Place from Wollaston's Cat.	0*
1595		VII. 58		7 48 39.8	2.700	1	107 21 4.6	9.22	1	Cl; pL; pRi; pC; st S.....	1
1596	473	II. 302		7 48 50.3	3.602	1	65 52 5.8	9.26	1	F; S; IE; bM; er	3
1597	473, a		R. nova	7 48 ±		...	65 52 ±		...	vF; E; * inv near N	0
1598	{ 474 = 3106 }	VII. 10		7 49 0.6	2.544	2	113 56 10.5	9.25	2	Cl; L; cRi; vIC	5
1599	3105			7 49 1.8	2.453	1	117 29 59.5	9.25	1	Cl; L; IC.....	1
1600	475	III. 837		7 50 26.3	4.881	1	33 3 54.6	9.42	1	vF; vS; R; glbM	2
1601	{ 479 = 3107 }	VII. 23	Δ. 626	7 50 36.9	2.396	3	119 41 58.1	9.37	3	Cl; pL; cRi; pC; st 11...13	4
1602	477			7 50 53.9	3.682	1	62 35 59.9	9.43	1	vF; S; R; bM.....	1
1603	477, a		R. nova	7 50		...	62 35		...	F; S.....	0
1604	476	III. 750		7 50 55.0	4.067	(1)	49 47 50.2	9.44	(1)	cB; S; R; sbM	2
1605	476, a		R. nova	7 50 +		...	49 47 ±		...	Follows III. 750 (h. 476) ...	0
1606		III. 838		7 51 11.3	4.903	1	32 42 17.4	9.48	1	eF; vS.....	1
1607	478	III. 709		7 51 25.4	+4.530	1	38 52 4.7	+9.49	1	F; L; R; vgbM; r; am st ...	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and m.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
1608	3108			7 52 3.2	+2.778	2	103 59 2.7	+ 9.49	2	cF; S; vIE90°; glbM; am st.	2
1609	3109			7 52 12.0	1.565	1	141 55 10.1	9.47	1	pF; S; R; vgpmbM	1
1610		III. 839		7 53 15.3	4.672	1	36 10 28.9	9.63	1	eF; vS	1
1611	480	VI. 37		7 53 15.8	2.858	1	100 14 21.4	9.58	1	Cl; pL; vRi; C; st 11...20...	2*
1612	481	II. 544		7 53 39.5	3.409	1	73 54 23.9	9.63	1	pB; pL; iR; vgbM; er; *225°.5, 60".	3
1613		VIII. 1		7 54 31.8	2.675	3	108 41 12.4	9.68	3	Cl; B; pRi; iC; stS	4
1614	482	III. 605		7 54 45.3	3.586	1	66 13 22.6	9.72	1	vF; S; iR	2
1615	483, a		R. nova	7 54 32.6	3.273	...	80 7 43.3	9.71	...	γ in Lord Rosse's diagram	0*
1616			D'Arrest, 51	7 54 40.8	3.27	[2]	80 12 48	9.70	[2]	eF; III. 512 f10°.5; n50"	0
1617	483	III. 512		7 54 47.2	3.273	2	80 11 58.3	9.71	2	F; S; R; psmbM; r	3
1618	484	III. 7		7 54 48.7	3.255	1	81 4 16.3	9.71	1	F; vS; vIE; 2 st p	3
5066				7 55 12.5		...	65 25 19.4		...	See No. 5066.	
1619	3111			7 56 1.4	1.004	6	150 29 10.5	9.75	6	Cl; vB; vL; pRi; st 7...13...	6
1620	3110			7 56 17.4	2.825	1	101 54 45.3	9.81	1	F; vS; R; bet 3 st 13, 14	1
1621	3112			7 56 48.9	2.457	1	117 47 40.5	9.85	1	Cl; B; pRi; pC	1
1622	485			7 57 22.1	4.938	1	31 49 50.5	9.95	1	pF; pL; R; psbM; *9, np 3.	1
1623	486	III. 877		7 59 0.5	2.845	1	101 2 6.9	10.03	1	eF; pL; R; vglbM; am st...	2
1624	488	VIII. 30		7 59 28.2	2.461	(2)	117 46 23.5	10.05	1	Cl; vL; pRi; iC; st 10...15...	3
1625	487	III. 752		7 59 54.2	3.451	1	71 46 29.0	10.10	1	eF; iE; vS*n	2
1626	489	II. 726		8 1 15.2	3.857	2	55 38 18.6	10.22	2	pB; pL; R; vglbM; r; 2 st nf	4
1627	3113			8 1 22.2	2.419	2	119 29 53.7	10.19	2	Cl; pL; Ri; C; st 9, 13...14.	2
1628	490	III. 840		8 1 50.7	4.775	1	33 55 30.7	10.29	1	pF; pL; R; psbM; *8, 164°-3	2
1629	491	IV. 55		8 3 18.1	4.271	1	43 35 28.4	10.38	1	⊕; pB; pL; R; rrr st 20...	3
1630	3114	VII. 11		8 4 7.8	2.819	2	102 24 59.0	10.40	2	Cl; vL; Ri; iC; st 11...13	3
1631	492	III. 710		8 4 26.2	4.404	1	40 30 20.4	10.48	1	F; L; E; vgbM	2
1632	3115		B.A.C. 3073	8 4 42.2	2.817	1	102 30 54.5	10.45	1	Nebulous *6.7	1
1633	493	II. 719		8 4 51.0	3.919	1	53 19 26.7	10.49	1	F; pL; iR; vgbM; * nr	3*
1634	494	II. 627		8 6 2.9	3.524	1	68 13 36.1	10.57	1	F; S; iE 45°; *8, np 4'	4
1635	3116		Δ. 563	8 6 26.4	2.215	3	126 58 24.1	10.57	3	Cl; B; L; iC; iE; st 9...12...	3
1636	3117		Δ. 411	8 6 31.6	1.769	2	138 51 3.1	10.57	2	Cl; B; L; iC; st 7...16	2
1637	496	VI. 22	C. H.	8 6 50.1	2.964	2	95 22 30.6	10.62	3	Cl; vL; pRi; pmC; st 9...13.	7
1638	495			8 7 37.5	4.892	1	31 46 48.9	10.73	1	pB; S; mE 0°; psmbM	1
1639		III. 711		8 8 46.0	4.438	1	39 32 33.9	10.73	1	eF; cL; iE 45°±	1
1640	497	II. 303		8 10 33.7	3.568	1	66 6 1.0	10.90	1	F; S; R; mbM; r	4
1641	498	III. 256		8 10 43.9	3.094	1	88 48 46.3	10.91	1	vF; cS; iF; 3Sst inv?	3
1642	499	III. 606		8 11 6.8	3.500	1	69 3 9.5	10.95	1	vF; S; R; sbM; stellar	3
1643	3118			8 11 21.1	2.499	1	117 2 10.9	10.93	1	F; pL; gmbM; am 60 st	1
1644			D'Arrest, 52	8 11 42	3.51	[3]	68 34 36	10.98	[3]	F; pL	0
1645	500	III. 607		8 12 13.6	3.512	1	68 26 14.6	11.02	1	vF; cS; R	2
1646	501	II. 634		8 12 26.5	3.510	1	68 30 37.2	11.04	1	eF; S; R; bM	2
1647	3119			8 12 27.9	2.633	1	111 22 33.6	11.02	1	vF; S; R; gbmM; am 60± st.	1
1648		III. 288		8 12 48.8	2.549	1	115 1 53.2	11.04	1	vF; cL; er	1
1649	{ 503 = 3120 }	VII. 64		8 12 58.1	2.421	3	120 12 19.8	11.06	3	{ Cl; pL; pRi; iC; iR; } st 11...14.	4
1650			D'Arrest, 53	8 13 10	3.50	[2]	68 41 48	10.98	[2]	vF; cE; 3vSst f	0
1651	502	VI. 39		8 13 15.1	+2.444	1	119 18 33.1	11.07	1	Cl; vL; cRi; iC; st 9, ...	3
1652	3176			8 13 25.4	-140.624	1	179 41 7.5	7.74	1	F; S; R; glbM; Polariss. Austr.	1*
1653		II. 259		8 14 16.8	+3.543	1	66 57 59.4	11.18	1	F; S; iF; r	1
1654	3121	III. 902		8 14 49.0	2.818	2	102 52 35.0	11.20	2	F; vIE; gbmM; r; am 50 st	3
1655	3122			8 15 35.1	2.272	4	125 46 36.2	11.24	4	* = h. 4083 in pS neb; am 70 st.	4
1656	3123			8 15 44.3	2.435	1	119 52 8.5	11.25	1	Cl; cL; pRi; pC; R; st 12...	1
1657	504	III. 753		8 17 3.0	3.488	2	69 13 21.1	11.37	3	vF; pS; R; glbM; * p 75"	4
1658	3124			8 17 39.0	2.461	1	119 2 32.8	11.46	1	Cl; pmCM; iF; st 9, 10...13.	1
1659	3125			8 17 39.2	2.369	1	122 31 10.1	11.47	1	Cl; F; S; R; gbmM; st 15	1
1660	505	II. 315		8 18 41.2	3.615	1	63 35 7.7	11.49	1	pF; S; R; vsbM*	2
1661	506	III. 599		8 19 34.0	3.511	1	68 3 46.8	11.56	1	vF; pL; iF; r; * sp 2'	2
1662	507	III. 234		8 23 58.8	+3.529	1	66 58 8.1	+11.87	1	vF; S; stellar	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
1663	3126			8 24 27.7	+0.438	1	157 39 23.2	+11.84	1	F; pS; R; gbM	1
1664	508			8 24 32.8	4.510	2	36 41 28.9	11.93	2	eF; S; R; *95°; p of 2	2
1665	509	III. 292		8 24 48.6	3.691	1	59 59 13.9	11.93	2	vF; pL; R; lbM; r; *nr ...	4
1666	510			8 25 8.2	4.500	2	36 44 52.5	12.05	2	eF; S; R; f of 2; *310°	2
1667	510, <i>a</i>		R. nova	8 25 8.2	4.500	::	36 40 52.5	12.05	::	Place from 510 h. by MS. ...	0*
1668	510, <i>b</i>		R. nova	8 25 ±		...	36 44 ±		...	No description or place	0
1669	511			8 25 27.2	3.628	1	62 32 51.4	11.98	1	eF	1
1670	512	II. 318		8 26 43.8	3.661	2	61 3 35.1	12.07	2	F; vIE; mbM; r	3
1671	3130			8 26 48.7	1.170	2	150 38 29.3	12.01	2	Cl; pS; lRi; lC	2
1672	{ 513 = 3127 }	IV. 35		8 26 57.1	2.771	2	105 39 51.8	12.06	2	F; S; att to *13; *7 nf, 10° ...	3
1673	3128			8 27 10.4	2.830	1	102 41 44.4	12.08	1	B; S; E; psbM; bet 2 st.	1
1674	3129	II. 266		8 27 14.9	2.628	3	112 29 47.4	12.08	3	eB; L; vME 110° 3	5
1675	515	III. 257		8 28 48.7	3.093	1	88 48 50.0	12.20	1	eF; pL; iF	2
1676	514	II. 319		8 29 0.5	3.662	3	60 49 18.6	12.22	3	F; pS; R; bM; r	4
1677	3131			8 30 27.5	2.178	3	130 11 3.7	12.29	3	*9 inv in pB, pL, R, neb ...	3†
1678	{ 516 = 3132 }	VII. 63		8 31 28.3	2.476	2	119 28 1.1	12.37	2	Cl; cL; pRi; pC; st 11...13..	4
1679		III. 982	H. O. N.	8 31 32.3	6.536	1	16 45 45.1	12.47	1	vF; S; stellar	1
1680		III. 235		8 31 51.1	3.539	1	65 57 15.6	12.42	1	eF; S	1
1681	517		M. 44	8 32 9.0	3.462	1	69 32 36.2	12.44	1	Præsepe Cancri	3
1682		III. 983	H. O. N.	8 32 33.3	6.501	1	16 51 47.2	12.54	1	vF; S; stellar	1
1683	3133			8 32 58.2	2.357	1	124 16 9.1	12.47	1	Cl; pmC; irr Δ; st 13	1
1684	518	I. 204		8 33 37.9	4.342	4	39 17 45.8	12.56	4	cB; S; E 130° ±; psmbM*?..	5
1685	3134			8 33 41.5	1.596	2	144 37 47.0	12.50	2	pB; S; R; 3 or 4 vS st p nr ...	2
1686	519			8 34 20.5	3.005	1	93 37 53.4	12.50	1	vF; pL; gbM; r; 2 pB st s, sl	1
1687	3136			8 34 20.5	2.000	1	135 43 55.5	12.55	1	Cl; S; st L & S	1
1688	{ 521 = 3135 }	III. 49		8 34 51.4	3.345	3	75 13 2.6	12.62	3	F; S; vIE 135° ±; psbM	5
1689	522	II. 727		8 35 17.8	3.802	2	54 47 16.5	12.65	3	F; L; R; r	4
1690		II. 908		8 35 58.9	6.024	1	19 11 53.8	12.76	1	pB; pL; iF; er	1
1691	520	I. 288		8 37 4.6	8.188	1::	11 15 26.7	12.89	1	vB; cL; lE 90° ±; g, svmbM.	2
1692	523			8 37 28.1	4.503	1	35 36 46.6	12.82	1	eF; psbM	1
1693	4017		Δ. 609	8 37 45.6	2.423	1	122 9 4.7	12.79	1	Cl; pS; lRi; lC; vIF; st 12, 13	1
1694	3137			8 37 50.2	2.060	2	134 27 15.7	12.79	2	Cl; L; Ri; pmE; st 11...14..	2
1695	3138			8 38 1.3	1.977	2	136 42 16.0	12.80	2	Cl; pS; mC; iR; gbM; st 13...15.	2
1696		III. 50		8 38 43.9	3.309	1	76 54 43.4	12.88	1	eF; cL; R; lbM	1*
1697	3139			8 38 59.6	2.800	1	104 47 19.7	12.89	1	vF; vS; R; bM; *15m nr ...	1
1698	524			8 39 29.7	3.309	1	76 53 9.2	12.94	1	Cl; st 9...10	1
1699			D'Arrest, 54	8 40 20	3.43	[1]	70 27 48	12.99	[1]	eF	0
1700	525			8 40 20.3	4.183	1	42 25 39.3	13.01	1	Cl; lC	1
1701	3140			8 40 50.8	1.694	1::	143 27 29.7	12.99	1::	Cl; L; P; lC; st 10...13	1
1702	3142			8 41 3.4	1.928	1	138 16 31.0	13.00	1	Cl; pL; P; lC; st 13	1
1703	3141		Δ. 489? 490?	8 41 7.1	2.174	1	131 22 30.3	13.01	1	Cl; pRi; lCM; st 12...13 ...	1
1704	526	II. 80		8 41 23.5	3.432	3	70 24 45.8	13.06	3	B; pL; lE 10° or biN; mbM*	5
1705	526 <i>a</i>		R. nova	8 41 23 ±		...	70 24 ±		...	Nearly in contact with h. 526 (see description of h. 526).	?
1706			D'Arrest, 55	8 41 52	4.46	[1]	35 58 48	13.11	[1]	vF; R; *15 p 12°, 270°	0
1707	527	II. 48		8 42 1.3	3.430	1	70 28 19.0	13.10	1	eeF; pL; lbM; r	2*
1708	528	VIII. 10		8 42 32.9	3.283	3	78 8 45.2	13.14	4	Cl; vIC; P	5
1709	529	III. 294		8 42 59.4	3.685	1	58 36 38.1	13.17	1	pF; vS; R; bM	2
1710	529, <i>a</i>		R. nova	8 43 ±		...	58 36 ±		...	Makes a close D neb with h. 529.	0
1711	530	I. 242		8 43 26.1	4.348	2	38 9 35.3	13.21	2	vB; L; vg, vsmbM*10	3
1712	531		M. 67	8 43 34.3	3.291	4	77 40 36.0	13.20	5	l; Cl; vB; vL; eRi; lC; st 10...15.	8*
1713	532	I. 200		8 43 58.0	+3.746	3	56 3 38.9	+13.23	3	vB; vL; vME 40° 9; gmbM..	5

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1714	h. 533	H. III. 712		h m s 8 45 2.1	s +4.247	2	40° 18' 38".6	+13.32	2	F; pL; R; gbM; 4 S st nr ...	3
1715	533, a		R. 3 novæ	8 45 ±		...	40 18 ±		...	4 (incl h. 533) nearly in a line	0
1716											
1717											
1718		II. 658		8 45 42.6	3.909	1	49 55 12.8	13.36	1	pF; vS; mbM	1
1719	534	III. 831		8 46 48.6	4.366	1	37 23 56.2	13.44	1	vF; S; R; psbM	2
1720	535	II. 823		8 46 53.6	4.333	1	38 7 1.2	13.44	1	pB; mE 0° ±; psmbM	2*
1721	536	II. 280		8 47 20.4	3.027	1	92 32 2.2	13.44	1	pF; cS; E 90° ±; bet 2 st ...	2†
1722	536, a		R. 3 novæ	8 47 ±		...	92 32 ±	13.48	...	No description	0
1723											
1724											
1725	536, b		R. nova	8 48 0.0	3.025	...	92 28 2.2		...	MS. No description	0
1726	538			8 48 33.4	3.025	1	92 39 34.6	13.52	1	vF; pS; R; r; *9 p	1
1727			D'Arrest, 56	8 48 43	3.025	[2]	92 34 48	13.53	[2]	vF; S; R; *15 p, 44" n; h. 538 nr.	0
1728	537	IV. 66		8 48 46.5	4.438	1	35 41 54.1	13.57	1	pB; fan-shaped; *11 att	2†
1729		III. 625		8 48 54.1	3.892	1	50 7 22.8	13.56	1	vF; vS	1
1730		II. 281		8 48 58.2	3.022	2	92 50 24.5	13.55	1	vF; pS, R	2
1731		III. 841		8 49 15.0	4.536	1	33 46 55.0	13.60	1	vF; S	1
1732	540			8 50 3.8	4.065	1	44 33 26.5	13.65	1	pB; L; E; vgbM *18	1
1733	3143			8 50 5.7	1.451	1	148 41 21.4	13.58	1	eF; S; R; psbM	1
1734	3144			8 50 52.5	2.632	2	114 8 0.1	13.67	2	pF; S; R; vgpmbM	2
1735	542			8 51 19.6	3.189	??	83 7 51.0	13.70	1::	F; pL; R	1*
1736		II. 557		8 51 19.6	3.189	1	83 9 32.0	13.70	1	F; pL; mE	1*
1737	541	III. 540		8 51 25.7	3.785	1	53 44 3.6	13.72	1	vF; S; E 110° ±; 2 vF st inv.	4
1738	539			8 51 28.6	8.388	1:	10 14 54.9	13.83	1::	pB; S; E 45° ±; *nf	1
1739	543	II. 529		8 51 59.2	2.997	2	94 21 30.2	13.74	2	cF; pL; R; vgbM	3
1740		III. 264		8 52 29.4	3.017	2	93 11 6.1	13.77	2	vF; vS; stellar	2
1741	544			8 53 6.7	3.781	1	53 42 41.9	13.83	1	eF; S; stellar	1
1742	545	II. 834		8 53 51.4	4.769	1	29 30 44.0	13.90	1	cF; pS; iR; er	2*
1743	546			8 54 2.0	3.023	1	92 50 15.1	13.87	1	vF; L; R; bM	1*
1744	547			8 55 0.5	3.018	1::	93 10 51.9	13.93	1	eF; R	1
1745	3145			8 55 30.8	2.097	1	135 20 55.2	13.94	1	!; eeF; vL; vvmE 19°	1†
1746			D'Arrest, 57	8 55 58	3.47	[2]	67 28 48	14.00	[2]	D neb; pB; S, not R; comes s 4'.	0
1747			D'Arrest, 58	8 55 59	3.46	[2]	67 32 48	14.00	[2]	vF; vS	0
1748	549			8 56 5.6	4.305	1	37 41 23.9	14.03	1	4 S st in neb	1
1749	549, a		R. nova	8 56		...	37 41		...	Makes D neb with h. 549 ...	0
1750	550	I. 249		8 56 26.3	4.788	1	28 58 21.8	14.06	1	cB; cL; E 90° ±; er	3
1751		III. 608		8 56 34.0	3.527	1	64 26 53.5	14.05	1	eF; S; R; vlbM	1
1752	551	III. 60		8 56 43.6	3.399	2	70 59 30.5	14.05	2	vF; S; R; r; *nr	3
1753	548			8 57 9.1	7.170	1:	12 56 4.8	14.16	1::	pB; pL; E; vglbM	1
1754	552	III. 825		8 57 15.0	3.760	4	54 3 51.7	14.09	4	eF; S; R; vglbM; *12. 345°, 50".	5
1755			D'Arrest, 59	8 57 28	3.39	[4]	71 7 5	14.09	[4]	pF; S; R; bMN=*15	0
1756		III. 291		8 57 34.4	3.514	3	64 0 27.3	14.11	3	vF; cL; R; bMN; 2 est p ...	3*
1756			D'Arrest, 60	8 57 35	3.53	[2]	64 0 24	14.10	[2]	*15.16 inv in pB; pL neb 40" diam.	0*
1757			D'Arrest, 61	8 57 55	3.53	[1]	64 6 12	14.12	[1]	vF; vS	0
1758		III. 626		8 58 51.7	3.928	2	47 44 29.7	14.19	2	vF; S; iF; lbM; r	2
1759	553	II. 828		8 58 52.0	4.387	1	35 35 23.3	14.21	1	pB; pS; E; vgbM	2
1760	554	III. 647		8 59 32.7	3.811	1	51 48 43.9	14.23	1	vF; cS; R	3
1761	560	III. 275		9 0 13.6	2.820	1	104 56 21.5	14.25	1	vF; pS; bM; S *30" n	2
1762	557	III. 236		9 0 16.2	3.450	1	67 59 30.1	14.27	1	cF; vS; R; er; bet 2 pB st ...	3
1763	558	II. 520		9 0 16.9	3.136	1	86 2 37.8	14.26	1	vF; pL; E; gbM; er	3
1764	556			9 0 29.6	4.225	1	39 2 40.0	14.30	1	eF; sbM *15; 1st of 3	1
1765	555	I. 250		9 0 41.1	4.728	1	29 23 40.6	14.32	1	cB; cL; iE; psmbMLBN ...	2
1766	559			9 0 51.3	4.224	2	39 0 48.6	14.32	2	pF; S; E; psbM; 2nd of 3 ...	2
1767	559, a		R. nova	9 0 ±		...	39 ±		...	One of 4 (h. 556, 559, 561); one vF; one E.	0
1768	562	II. 490		9 0 56.7	+3.694	3	56 18 26.3	+14.31	3	F; L; mE 150°; r; 2 st n	4

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. II.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s		° ' "	"			
1769	561			9 1 0.0	+4.226	2	39 4 21.1	+14.27	2	vF; vS; IE; 3rd of 3	2
1770	3146			9 1 27.6	2.671	1	113 5 19.6	14.32	1	eF; IE; lbM?	1
1771	564	I. 2		9 2 52.2	3.195	2	82 23 38.9	14.43	2	cB; cL; R; vg; vsmbM; r? ..	7
1772	563			9 2 55.9	4.014	1	44 28 15.5	14.45	1	pB; L; R; vgbM; r	1
1773	565	III. 61		9 3 1.4	3.377	1	71 44 55.2	14.44	1	vF; S; R; am 5St; (PD 70°)	2*
1774	566	II. 564		9 3 44.1	3.732	4	54 24 3.7	14.49	4	pB; S; R; psmbM	6
1775	566, a		R. nova	9 3		...	54 24		...	eF; companion of h. 566, 567.	0
1776	567	III. 826		9 4 3.5	3.728	1	54 29 46.3	14.51	1	vF; S; R; S* 7.5 p	2
1777	{ 569 = 3147 }	I. 66		9 4 50.4	2.837	2	104 14 58.2	14.54	2	B; S; pmE 90°±; psmbM...	3
1778	568	I. 167		9 5 16.1	3.858	(1)	49 21 24.4	14.58	(1)	cB; R; mbMBN	2
1779		III. 295		9 5 21.2	3.613	1	59 23 25.4	14.58	1	vF; vS; R; 2pB st sp	1
1780	{ 571 = 3148 }	I. 59		9 6 5.8	2.670	3	113 36 36.3	14.61	3	B; L; mE63°7; gmbM	5
1781	570	I. 216		9 6 25.9	5.535	1	20 12 11.7	14.69	1	B; pL; IE90°±; mbM; r; vS*sf inv.	4
1782	3150			9 6 36.0	0.866	1:	157 21 50.0	14.60	1	vF; vS; mE105°	1
1783	3149			9 7 10.6	2.253	3	131 51 37.1	14.67	3	!; O; pB=*9; vS; R; am st.	4
1784	572			9 8 11.6	3.705	2	54 59 4.8	14.76	2	vF; S; R; * p1*, n5'	2
1785	573	III. 296		9 8 17.0	3.624	1	58 31 50.8	14.76	1	eF; S; R; lbM	2
1786	575	III. 62		9 8 21.5	3.392	(1)	70 27 55.8	14.76	1	vF; S; R; r	2
1787	575	III. 63		9 8 21.8	3.392	(1):	70 28 25.8	14.76	1	vF; S; R; r	2
1788		II. 708		9 8 31.6	3.899	1	47 27 36.4	14.78	1	pB; S; stellar	1*
1789	574	III. 832		9 8 39.8	4.271	1	36 54 12.0	14.80	1	vF; S; IE; *att; *inv	2
1790		III. 878		9 8 45.5	4.971	2	25 18 38.6	14.82	2	vF; L; R; mbM	2
1791	577			9 8 54.8	3.413	1	69 13 19.7	14.79	1	vF; S; R; np of 2	1*
1792			D'Arrest, 62	9 9 5	3.34	[3]	69 22 9	14.80	[3]	vF; vS; h. 578 f7*5; Δ.P.D. 118".	0*
1793	3152		Δ. 265	9 9 9.9	1.185	4	154 17 18.8	14.76	4	!; ⊕; vL; eRi; vgeCM; 45°d; st 13...15.	4
1794	578			9 9 12.3	3.410	1:	69 21 43.3	14.81	1	vF; S; R; sf of 2	1
1795		III. 749		9 9 35.0	5.953	1	17 34 40.4	14.88	1	F; cS; bM	2
1796	{ 580 = 3151 }	II. 505		9 9 37.1	2.817	2	105 43 14.6	14.82	2	pB; pS; E 45°±; psmbM ...	3
1797		II. 868		9 9 39.0	4.975	(1)	25 9 42.1	14.87	(1)	F; S; iF; 1st of 2	1
1798	576	II. 869		9 9 41.2	4.974	1	25 10 11.1	14.87	1	F; S; E; 2nd of 2	2
1799	3153	III. 242		9 10 3.6	2.688	1	113 2 16.5	14.85	1	F; S; IE; gbm	3
1800	579			9 10 17.0	4.684	1	28 57 59.7	14.89	1	F; pmE	1
1801	3154		Δ. 564	9 10 21.5	2.417	2	126 1 53.8	14.86	2	!; OpB; pL; R; vglbm; in L, C, Cl.	2†
1802	3155			9 10 39.7	2.628	1	116 14 46.4	14.88	1	eF; *11 att	1
1803	3156			9 10 41.4	0.713	1:	159 3 42.5	14.85	1	pF; vS; R; glbm	1
1804	581, a		R. nova	9 10 43.8	3.680	::	55 49 34.9	14.88	::	R.MS. No description	0*
1805	581, b		R. nova	9 10 43.8	3.680	::	55 19 34.9	14.88	::	R.MS. No description	0*
1806	581			9 10 50.9	3.680	1	55 40 39.3	14.91	2	vF; E; I. 113 f	2*
1807			D'Arrest, 63	9 10 52	3.68	[2]	55 47 48	14.91	[2]	vF; vS; R; h. 581, 6' n	0*
1808	581, c		R. nova	9 10 53.4	3.680	::	55 29 4.9	14.92	::	R.MS. No description	0*
1809	581, d		R. nova	9 10 58.4	3.680	::	55 47 34.9	14.93	::	R.MS. No description	0*
1810	582, a		R. nova	9 11 3.1	3.680	::	55 28 34.9	14.93	::	R.MS. No description	0*
1811	582	I. 113		9 11 12.9	3.680	2	55 39 34.9	14.93	2	cB; cL; IE; mbf; 3 sts	4*
1812	582, b		R. nova	9 11 14.5	3.680	::	55 40 20.3	14.92	::	β in Lord R.'s diag. } v nr I.	0*
1813	582, c		R. nova	9 11 16.0	3.680	::	55 27 34.9	14.92	::	α in Lord R.'s diag. } 113.	0*
1814	582, d		R. nova	9 11 17.6	3.680	::	55 27 34.9	14.92	::	R.MS. No description	0*
1815	582, e		R. nova	9 11 32.6	3.680	::	55 42 29.3	14.92	::	ε of Lord R.'s diagram	0*
1816	3157			9 11 41.7	+0.760	1	158 45 49.0	+14.90	1	F; pS; R; glbm	1

No. of Cata- logue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s		° ' "	"			
1817	582, f		R. 2 novæ	9 11 ±		...	55 40 ±		...	2 of 15 seen.....	0*
1818				9 11 45.3	+2.817	1	105 53 11.2	+14.94	1	eF; R; bM; *f8*.5.....	1
1819	585			9 11 50.3	3.813	1	50 7 15.1	14.97	1	vF; vS; R.....	3
1820	583	III. 627		9 12 6.0	3.680	::	55 46 35.3	14.98	::	ξ of Lord R.'s diagram	0*
1821	582, g		R. nova	9 12 13.5	3.714	1	54 2 26.7	14.99	1	cF; S; R; *10 np 2'	3
1822	586	III. 827		9 12 19.6	4.187	1	38 25 33.3	15.01	1	vB; L; vme 150°.8; vmbM =*10.	3
1823	584	I. 205		9 12 29.8	1.357	1	152 28 51.8	14.96	1	F; vS; bet 2 st.....	1
1824	3158			9 12 41.2	3.385	1	70 28 52.3	15.01	1	S* and neb.....	1
1825		III. 64		9 12 53.2	3.832	1	49 15 51.9	15.03	1	cF; cS.....	1
1826		III. 628		9 13 2.9	2.392	1	127 25 37.3	15.01	1	vF; S; R; *12 att sf	1
1827	3159			9 13 30		...	105 55		...	np 587 h.; close	0
1828	587, a		R. nova	9 13 30.4	2.819	1	105 55 14.5	15.05	1	vF; cL; E 45° ±; glbM; *11 sf 9°.	2
1829	587	III. 488		9 13 52.9	2.330	1	129 57 21.8	15.06	1	eF; cL; R; vglbM rr	1
1830	3160			9 14 20.7	3.826	1	49 16 29.6	15.12	1	vF; cS; R; *10 p 2'; 1st of 2	2
1831	588	III. 629		9 14 24.2	3.827	1	49 14 39.6	15.12	1	vF; S; vglbM; 2nd of 2	2*
1832	590	III. 630		9 14 30.9	4.108	2	40 11 39.9	15.13	2	cF; cS; vIE; pglbM; 1st of 2	3
1833	589	III. 714		9 14 ±		...	40 11 ±		...	Seen with h. 589, 591	0
1834	589, a		R. nova	9 14 41.5	2.895	1	101 18 57.6	15.12	1	pB; pL; R; gmbMN	4
1835	592	I. 132		9 14 44.2	4.108	2	40 9 20.2	15.14	2	cF; cS; IE; bM; 2nd of 2	3
1836	591	III. 713		9 15 45.1	3.684	3	54 53 13.7	15.19	3	vB; pL; R; smbM	4
1837	593	I. 137		9 16 48.2	2.920	1	99 50 3.2	15.24	1	cF; S; E; bet 2 st 12, 16	2
1838	594	III. 520		9 17 8.3	3.259	1	77 46 1.8	15.26	1	F; vS; p of 2	1
1839		II. 57		9 17 11.2	3.258	1	77 46 16.1	15.27	1	pF; S; f of 2	1
1840		II. 58		9 17 11.3	2.710	2	112 34 33.8	15.26	2	B; S; R; gbM.....	2
1841	3161			9 17 21.9	2.016	1	140 30 21.5	15.25	1	Cl; IC	1
1842	3162			9 17 28.1	1.694	7	147 42 57.5	15.25	7	!!; O = *8; vS; R; *15, 59°.13, 13".	8
1843	3163			9 17 37.9	4.454	1	32 1 44.8	15.26	1	cF; S; E; vglbM	2
1844	595	III. 846		9 18 8.2	3.255	5	77 57 53.6	15.32	6	pF; pS; R; bM; p of 2, 109°	7
1845	597	II. 546		9 18		...	77 58		...	Forms Δ with 2 E neb	0
1846	597, a		R. nova	9 18 13.1	3.255	6	77 58 34.9	15.33	6	vF; pL; R; bM; f of 2	7
1847	598	II. 547		9 18 33.5	4.758	2	26 54 27.1	15.37	2	B; cS; R; mbM; am st	3
1848	596	I. 260		9 19 30.4	2.502	1	123 29 36.4	15.38	1	vF; S; vglbM; rrr; st 11m... ..	1
1849	3164			9 19 42.4	3.445	1::	66.22 59.3	15.41	1	eF; vS; E 90° ±	1
1850	599			9 20 8.5	2.740	1	111 8 37.6	15.42	1	eeF; pL	1
1851	3165			9 20 11.4	1.370	2	153 12 29.0	15.40	3	F; S; R; pmbM; B*nr	3
1852	3168			9 20 12.5	2.627	2	117 25 24.9	15.43	2	cF; S; R; gmbM	2
1853	3166			9 20 24.7	2.904	1	101 2 12.2	15.44	1	pF; pS; vIE; vglbM; r	2
1854	600	II. 555		9 20 41.0	2.688	1	114 11 43.5	15.45	1	F; S; R; bM	1
1855	3167			9 21 59.9	3.562	2	59 49 57.2	15.54	2	vF; S; R; vsbM*12	3
1856	602	III. 297		9 22 2.3	3.195	1	81 39 42.2	15.54	1	vF; E; er; 2 or 3 st inv	4
1857	603	III. 8		9 22 12.0	4.421	1	31 54 24.1	15.57	1	vF; vS; R; vglbM; *7' s	1
1858	601			9 22 42.2	1.839	1	145 30 15.5	15.55	1	F; pL; R; gmbM; am 80 st ..	2
1859	3169			9 24 5.4	2.861	1	104 6 31.2	15.64	1	vF; vS; stellar.....	1
1860		III. 276		9 24 14.6	3.409	2	67 53 46.8	15.66	3	cB; vL; E; gmbM; r; sp of 2 ..	4†
1861	604.1	I. 56		9 24 15.4	2.592	1	119 46 47.5	15.65	1	F; S; IE; psbM	1
1862	3170			9 24 16.3	3.410	2:	67 52 50.1	15.67	2:	vF; cL; R; psbM; r; nf of 2 ..	4†
1863	604.2	I. 57		9 24 37.6	3.203	2	80 57 11.4	15.68	2	F; pS; IE; gbM	3
1864	606	II. 495		9 25 2.9	2.830	1	106 7 27.0	15.70	1	pF; S; IE 90...180; mbsf ..	2
1865	607	II. 506		9 25 15.9	7.880	1	9 37 30.0	15.80	1	eF; vS	1
1866		III. 977		9 25 26.7	4.972	1	23 26 15.8	15.76	1	eF; S; psbM	1
1867	605			9 25 37.0	1.992	3	142 17 26.0	15.70	3	Cl; cL; pRi; pC; st 10...14.	3*
1868	3171			9 26 15.3	3.228	1	79 13 53.1	15.77	1	F; pL; R; gbM; p of 2	4
1869	608	II. 40		9 26 33.1	+3.227	1	79 16 23.4	15.78	1	vF; S; R; bMN, f of 2	2
1870	609	III. 513		9 26 54.2	-0.275	1	166 0 46.2	15.74	1	pF; pL; R; gbM	1
1871	3174			9 26 57.2	+3.408	2	67 40 32.3	15.81	2	F; S; vIE	3
1872	610	II. 260		9 27 24.1	3.589	1	57 40 57.2	15.84	1	vF; cS; R; sbMN	3
1873	611	III. 298		9 27 49.1	+2.769	1::	110 13 55.2	+15.84	1::	eF; S; R; p of 2	1
1874	3172										

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
1875	3173	III. 597		9 28 2.3	+2.768	2	110 17 59.8	+15.86	2	vF; pS; IE; vglbM; f of 2...	3
1876	3175			9 28 32.8	2.840	1	105 46 48.4	15.88	1	pB; S; R	1
1877	3177			9 28 57.4	1.993	1	142 49 22.7	15.89	1	Cl; pRi; pC; * taken.....	1
1878			D'Arrest, 64	9 29 32	3.43	[1]	66 10 24	15.94	[1]	eF; vS; IE; vlbM; 1st of 3...	0
1879			D'Arrest, 65	9 29 34	3.43	[1]::	66 12 12	15.94	[1]	eF; S; 2nd of 3	0
1880			D'Arrest, 66	9 29 42	3.43	[1]	66 8 18	15.95	[1]	eF; vS; 3rd of 3	0
1881	3179			9 30 6.6	2.221	1::	136 18 58.8	15.96	1::	Cl; eL; vRi; st L & S	1
1882	3178	II. 556		9 30 13.5	2.768	2	110 30 23.1	15.97	2	pB; pS; vIE; gmbM	5
1883	612	III. 963		9 30 28.4	6.574	1	12 47 47.8	16.06	1	eF; S; iFig; * f 3'	2
1884	614	III. 4		9 30 39.3	3.215	2	79 51 27.0	16.00	3	vF; S; vIE; bM; Δ st nf....	5
1885	613			9 30 42.6	3.626	2	55 21 52.3	16.01	2	pL; vIE0°; vglbM.....	2
1886	3180			9 31 15.5	2.756	2	111 24 53.9	16.03	2	F; S; R; glbM; 2 or 3 S st nr	2
1887	615	III. 519		9 31 34.7	3.178	(1)	82 24 1.5	16.05	1::	vF; pL; vgbM	2
1888	616	IV. 68		9 32 17.7	4.410	2	30 31 15.3	16.11	2	eF; vS; R; vgvmbMN	3
1889	3182			9 32 38.7	3.291	1::	74 32 3.3	16.11	1::	eeF; susp	1
1890	3181			9 32 44.3	3.292	1	74 26 8.3	16.11	1	vF; S; R; n of 2	1
1891	620	III. 541		9 32 45.9	3.659	1	53 29 0.6	16.12	1	eF; pS; iR; glbM; r	4
1892	617			9 33 7.8	5.746	1	16 22 12.4	16.18	1	eF; *13 nr	1
1893	618			9 33 12.5	5.149	1	20 45 24.1	16.17	1	F; pL; R; vglbM; * n	1
1894	621			9 33 24.7	3.131	1	85 46 2.7	16.19	1	vF; R; gbM	1
1895	619	III. 315		9 33 39.4	5.734	1	16 23 43.3	16.21	1	vF; vS; R; bM	2
1896	622	I. 114		9 34 34.3	3.571	4	57 31 54.6	16.22	3	B; vL; IE; vgbM; p of 2	5
1897	623	III. 751		9 34 46.9	3.660	1	53 6 33.9	16.23	1	eF; vS; R; bM; r	3
1898	626	II. 275		9 34 49.3	3.084	1	89 1 56.6	16.22	2	pF; pL; R; vglbM	4
1899	624	II. 491		9 34 51.5	3.572	4	57 25 34.9	16.23	4	pB; pL; IE; vglbM; f of 2...	5
1900	628	III. 527		9 34 59.1	2.961	1	97 57 26.9	16.23	1	vF; pS; iR; vglbM.....	3
1901	627			9 35 7.4	3.573	1	57 21 1.2	16.24	1	F; nf of 3	1
1902	3183		Δ. 397	9 35 16.9	2.143	1	139 41 32.9	16.23	1	Cl; S; lRi; pC; st 13	1
1903	4018			9 35 22.9	2.628	1	119 24 42.2	16.24	1	eF; pS; B*8m f	1
1904	630	I. 61		9 35 27.7	3.029	1	93 4 16.5	16.25	1	B; eS; iR; bM; *9 sp 3'	3
1905	625	I. 285		9 35 35.6	5.048	1	21 26 21.7	16.29	1	B; vL; mE 152°4; st inv	2
1906		I. 282		9 35 47.6	6.111	1	14 15 5.9	16.33	1	cb; pL; iF	1
1907	631	III. 521		9 36 17.2	2.937	1	99 44 57.7	16.29	1	pF; pS; vIE; psbM.....	2
1908	632	III. 528		9 36 18.4	2.948	1	98 58 2.7	16.29	1	vF; pS; IE 0°±; vglbM.....	3
1909	629	I. 78		9 36 41.2	5.576	1	17 4 29.8	16.36	1	vB; eL; R; psmbM; * inv f.	3
1910	3184			9 36 52.2	2.330	1	133 33 40.3	16.31	1	Cl; P; E; st 10...11	1
1911	3185	III. 289		9 37 7.5	2.791	1	109 49 54.9	16.33	1	F; pS; R; bM; r; stellar	4*
1912	633	III. 34		9 37 23.0	3.231	(1)	78 20 10.5	16.35	2::	eF; vS; R; bM (? P.D. 15')..	3
1913		II. 311		9 38 10.3	2.780	3	110 37 58.7	16.39	3	pB; pS; iR; mbM	3
1914		II. 624		9 38 39.5	3.156	1	83 41 19.3	16.41	1	F; pS; IE 90°±	1
1915	3186			9 38 52.0	2.825	1	107 44 26.6	16.42	1	F; R; gbM; * f	1
1916	634, a		R. nova	9 38		...	67 20		...	Makes a D neb with h. 634; which follows it.	0
1917	634			9 38 55.9	3.391	1	67 20 5.9	16.43	1	F; vS; bM; sp of 2	1
1918	635	III. 277		9 38 58.1	2.884	1	103 41 8.9	16.43	1	eF; S; R; bM; stellar; p of 2	2
1919	637	III. 278		9 39 4.6	2.884	1	103 43 23.9	16.43	1	eF; S; R; bM; stellar; f of 2	2
1920	636			9 39 18.1	3.391	1	67 16 32.5	16.45	1	F; S; R; bM; nf of 2.....	1
1921	3189			9 39 21.6	2.008	1	144 8 1.9	16.43	1	Cl; P; IC; st mm.	1
1922	3187			9 39 28.0	2.778	1	110 56 47.5	16.45	1	vF; S; * 20 f 1'	1
1923	3188	V. 50		9 39 32.5	2.619	2	120 32 50.5	16.45	2	i; vF; vL; vg, vsbMN 4"; 19°5 d.	3
1924	638	II. 717		9 40 2.0	3.823	1	45 15 27.0	16.50	1	pF; pL; iR; bM; r.....	2
1925	638, a		R. nova	9 40 4.8	3.823	...	45 16 27.0	16.48	...	RMS	0
1926	638, b		R. nova	9 40 4.8	3.823	...	45 16 57.0	16.48	...	Suspected; MS	0
1927	3191		Δ. 397	9 40 5.0	2.167	1?	139 47 14.1	16.47	1?	Cl; S; lRi; iF; st 12...15	1
1928	638, c		R. nova	9 40 7.6	2.823	...	45 12 27.0	16.48	...	MS	0
1929	3190			9 40 10.4	2.634	1	119 48 2.4	16.48	1	F; S; R; *12 att 320°	1
1930	638, d		R. nova	9 40 13.3	3.823	...	45 21 27.0	16.49	...	MS	0
1931	639	V. 26		9 40 15.0	3.584	1	55 56 14.0	16.50	1	i; cb; L; vimE 90°.....	3+
1932	638, e		R. nova	9 40 30.2	3.823	...	45 13 27.0	16.52	...	MS	0
1933	640			9 41 1.2	+3.825	1	45 1 50.5	+16.55	1	pF; R; bM; r; p of 2.....	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
1934	h. 638, f	H.	R. nova	h m s 9 41 4.0	s +3.823	...	45° 15' 27.0"	+16.55	...	MS	0
1935	641			9 41 7.2	3.825	1	45 0 16.5	16.55	1	F; psbM; rr; f of 2.....	1
1936	} 641, a		R. novæ	9 41 ±		...	45 0 ±		...	Several near	0
1937				9 41 ±		...	45 0 ±		...		0
1938			D'Arrest, 67	9 41 32	3.49	[1]	54 38 7	16.56	[1]	vF; pL; R; cometary	0
1939	642			9 41 32.1	3.251	1	76 32 0.8	16.56	1	F; pL; R; glbM.....	1
1940	642, a		R. nova	9 41		...	76 32		...	3 "novæ," with 642 (Vide h. 646, 648).	0
1941	644			9 42 3.2	3.011	1	94 33 38.4	16.58	1	eF; L; p of 2	1
1942	646	III. 51		9 42 31.0	3.250	3	76 31 58.3	16.61	3	eF; pS; IE 0°±; r	4
1943	647			9 42 38.9	3.012	1	94 31 9.3	16.61	1	F; R; vglbM; f of 2	1
1944	645	I. 115		9 42 39.6	3.579	2	55 47 35.6	16.62	2	pB; pS; vIE; mbM; *10 sf 100°.	4
1945	648	III. 52		9 42 54.1	3.249	1	76 36 18.9	16.63	1	eF; pL; E; r	2
1946	3192			9 42 58.1	2.782	1	111 5 19.6	16.62	1	eF; vS; R; *9s	1
1947	643	V. 23		9 43 9.9	5.466	1	17 8 23.4	16.68	1	vF; vL; IE; r	3
1948	3193			9 43 17.7	2.821	1	108 31 38.2	16.64	1	F; S; R; lbM	1
1949	649		M. 81	9 43 48.9	5.066	1	20 16 10.0	16.70	1	l; eB; eL; E 156°0; g, svmbMBrN.	4
1950		{ IV. 79 = 4H. ON }	M. 82	9 43 52.3	5.142	1	19 34 16.3	16.71	1	vB; vL; vmE "a beautiful ray."	2*
1951	650			9 44 0.3	3.497	2	60 7 7.4	16.68	2	F; S; sbM *12; bet 2B st ...	2
1952	3194		B. 2686	9 44 1.8	1.975	2	145 45 42.8	16.66	2	Cl; pL; pRi; iF; st 11...12...	2
1953		W. H. nova?	M. 81??	9 44 38.0	5.064	1	20 12 18.9	16.73	1	vB; eL; mE; 5 or 6 st (?) inv	1*
1954	3197			9 45 5.8	1.674	1	152 2 12.3	16.71	1	Cl; eL; iC	1
1955	3195			9 45 6.2	2.705	1	116 22 6.6	16.72	1	F; pS; R; lbM	1
1956	3196	II. 98		9 45 25.6	3.300	1	72 39 56.5	16.75	1	⊕; F; L; R; vglbM; rr; 2B stsp.	4
1957	651	II. 835		9 46 19.3	4.327	2	30 2 40.3	16.81	2	eF; pS; IE; vgbM; *10 n 7'	4
1958	652	III. 254		9 46 26.5	3.097	1	87 55 58.0	16.80	1	vF; vL; vmE 111°5	3
1959	3198			9 46 38.8	2.834	1	107 59 3.0	16.80	1	vF; pS; R; lbM	1*
1960	3199			9 47 1.9	2.704	1	116 40 25.6	16.82	1	pF; R	1*
1961	3201			9 47 41.3	2.707	1	116 37 56.5	16.85	1	pF; S; R; gbM	1*
1962	3202	III. 272		9 47 50.8	2.836	1	107 58 44.8	16.86	1	F; pL; R; glbM	2*
1963	3200	III. 600		9 47 53.6	3.293	1	72 54 4.1	16.87	1	vF; S; vIE; gbM	2
1964	656	VI. 4		9 47 59.1	3.133	2	85 4 19.1	16.87	2	F; pL; vIE; vgbM; rr; *7f90°.	4
1965	3203			9 48 15.8	2.692	2	117 38 37.4	16.88	2	pB; S; R; vgmbM; *11 att 203°8.	2
1966		III. 978		9 48 24.1	7.497	1	9 3 41.8	16.96	1	eF; pL; vlbM; 2 S st s	1
1967	3205			9 48 31.2	0.647	1	163 16 15.5	16.85	1	F; L; iR; glbM; S * inv ...	1
1968	3204	III. 601		9 48 39.8	3.297	1	72 30 4.3	16.91	1	vF; eS; vIE; er	2
1969	654	II. 333		9 48 42.7	5.376	1	17 9 25.2	16.94	1	pF; vS; R; bM; *11 nr.....	3
1970	653	II. 903		9 48 44.4	6.102	1	13 10 9.5	16.95	1	vF; pL; r	2
1971	655	II. 334		9 48 57.4	5.366	1	17 12 16.5	16.95	1	vF; vS; vglbM	3
1972		II. 909	H. ON 5	9 49 50.5	5.382	1	17 12 24.7	16.99	1	F; pL; R; 3rd of 3.....	1
1973	657	II. 492		9 50 7.9	3.533	3	56 57 50.4	16.98	3	pB; pL; E 90±; gbM; *9 nf	4
1974		III. 293		9 50 32.8	3.474	1	60 21 51.7	16.99	1	eeF; eS; stellar (?).....	1*
1975	659	II. 59		9 50 39.1	3.209	1	78 58 42.7	16.99	1	pB; pS; R; gmbMN; 3 st nr	2
1976	3206	III. 273		9 50 47.0	2.831	2	108 40 49.7	16.99	2	vF; pS; IE; glbM	3
1977		III. 853		9 51 10.7	4.126	1	33 42 23.9	17.03	1	vF; S; vglbM	1
1978	660	III. 542		9 51 18.7	3.585	1	53 55 47.9	17.03	1	vF; pL; iR; vglbM	4
1979	3207			9 51 19.5	3.261	1	74 53 56.9	17.03	1	vvF; *14 att; *11 f	1
1980	3208			9 51 32.2	2.848	1	107 30 25.9	17.03	1	eF; S; R	1
1981	3209	II. 268		9 52 4.6	2.731	1	116 15 22.5	17.05	1	pB; S; R; mbM	2
1982	658	I. 286		9 52 7.4	4.925	(2)	20 35 17.7	17.09	1	cB; eL; mbM; R with ray...	3
1983		V. 47		9 52 24.3	4.119	1	33 38 27.7	17.09	1	vB; L; mE 135°±	1
1984		III. 934		9 52 34.6	3.241	1	76 19 59.7	17.09	1	vF; S; R	1
1985		III. 596		9 52 37.0	2.786	2	112 7 58.4	17.08	2	vF; eS; lbM; ΔS st np	2
1986	3210			9 52 40.5	2.671	1	119 41 50.4	17.08	1	vF; S; R; * att	1
1987	3211			9 52 45.4	+2.722	1	116 28 7.7	+17.09	1	vF; S; R; *13 att sf	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
1988	3212			9 52 50.7	+2.832	1	109° 50' 56.7	+17.09	1	vF; S; R.....	1
1989	3213			9 53 4.5	2.607	1	123 33 46.0	17.10	1	pB; S; R; pmbM; bet 2 st...	1
1990	661	III. 24		9 53 17.7	3.367	1	66 55 52.6	17.12	1	vF; S.....	2
1991	3214			9 53 19.7	2.705	1	117 38 50.3	17.11	1	pF; pS; R; vS at inv.....	1
1992	3215	II. 293		9 53 38.9	2.831	1	108 57 52.9	17.13	1	pB; pS; iR; bM; p of 2.....	2
1993	3216			9 53 53.3	2.655	1	120 52 53.2	17.14	1	F; L; E; vglbM.....	1
1994	3217			9 54 2.5	2.832	1	108 57 58.5	17.15	1	eF; R; lbM; f of 2.....	1
1995	663			9 54 22.3	3.398	2	64 37 15.1	17.17	2	pB; S; mE 90°±; psbMN...	2
1996	664	III. 478		9 54 25.4	3.525	1	56 37 10.4	17.18	1	eF; S.....	3
1997	3218			9 54 27.6	2.655	1	120 59 37.8	17.16	1	pB; pS; R; gpmbM.....	1
1998	662	III. 916		9 54 36.7	4.298	1	29 13 26.7	17.19	1	vF; vS; R; bM; *11, 142° 2	2
1999	665	IV. 48		9 55 19.4	3.675	1:	48 35 27.3	17.21	1:	eF; pL; E 45°±; vF * inv...	2
2000	3219			9 55 46.4	2.120	2	144 6 16.3	17.21	2	Cl; C; lE; st 13...16.....	2
2001	666	II. 320		9 55 58.4	3.504	1	58 8 19.2	17.24	1	F; S; R; sbM.....	2
2002	3220			9 56 17.6	2.660	2	121 0 33.2	17.24	2	F; S; R; glbM.....	2
2003	3221			9 56 40.1	2.746	2	115 29 25.8	17.26	2	cF; vL; vmE 82° 3; lbM...	2+
2004		II. 898		9 56 54.3	3.242	1	75 49 0.4	17.28	1	F; L red * n 3'.....	1
2005	667			9 57 14.8	3.818	3	42 3 28.0	17.30	3	pB; S; R; SmbM *12.....	3
2006	3222			9 58 3.2	2.716	1	117 46 13.6	17.32	1	eF; L; Δ 2 st 8m.....	1
2007	3224		Δ. 297	9 58 11.2	1.934	3	149 26 46.6	17.32	3	Cl; eL; lC; B; st 9...14.....	3
2008	{ 668 = 3223 }	I. 163		9 58 14.3	2.988	2	97 2 32.9	17.33	2	vB; L; vmE 45°; vg, vsmbMEN.	3
2009	3225			9 59 14.8	2.627	1	123 32 36.4	17.38	1	F; pS; R; gbM.....	1
2010			Auw. N. 26	9 59 18.4	3.247	...	74 55 35.7	17.39	...	F; (Lassell, Mar. 31, 1848)...	0
2011		II. 305		9 59 26.0	3.003	1	95 51 17.9	17.43	1	F; S; lE; er.....	2
5067				9 59 51.1		...	89 15 7.3		...	See No. 5067.	...
2012	3226			10 0 2.2	2.848	1	108 33 36.3	17.41	1	F; pL; R; lbM; * s.....	1
2013	3227			10 0 14.8	2.700	2	119 15 12.6	17.42	2	cF; S; R; gbM.....	2
2014	669	III. 65		10 0 40.4	3.299	(1)	70 53 43.5	17.45	1:	eF; cS; vLE; r.....	2*
2015	670			10 0 45.2	3.195	1	79 20 43.5	17.45	1	eF; S; psbM; 31 Leon sf 100°	1
2016	671			10 0 54.9	3.296	1	71 4 49.8	17.46	1	pB; pS; pmE; gbM.....	1
2017	3228			10 1 8.2	2.523	4	129 45 5.5	17.45	4	ll; O; vB; vL; lE; *9M; 4° 0 d.	4+
2018	3229			10 1 46.2	1.540	1?	156 41 44.1	17.47	1?	B; R; bM.....	1
2019	672			10 2 9.8	3.759	1	43 21 12.6	17.52	1	F; S; R; gbM.....	1*
2020	3231			10 2 9.9	1.545	2	156 41 30.4	17.48	2	pB; pS; R; gbM; * 13 n...	2
2021	3230			10 2 30.4	2.719	1	118 22 22.3	17.51	1	vF; S; lE.....	1
2022	3232			10 3 9.9	2.982	1	97 47 52.5	17.55	1	F; R.....	1
2023	673	III. 518		10 3 17.7	2.936	1	101 44 15.8	17.56	1	F; pL; R; vg, slbM; f of 2...	4
2024	674	I. 79		10 4 47.5	5.285	1	15 54 35.5	17.65	1	vB; L; R; vg, vsmbM.....	2
2025	675			10 4 48.3	3.861	1	38 49 21.9	17.63	1	* 7 m in photosphere 2' or 3' d	1
2026	677	III. 53		10 5 8.4	+3.222	1	76 38 47.9	17.63	1	eF; pL; vLE; r; st inv.....	3
2027	3234			10 5 10.7	-0.511	1	169 43 54.4	17.58	1	F; S; lE; vlbM; * 15 inv...	1
2028	680	III. 255		10 5 25.3	+3.113	1	86 10 27.2	17.64	1	F; cS; R; psbM; Δ B st f...	3
2029	3233			10 5 27.7	2.700	1	120 16 4.2	17.64	1	vF; pS; E; * 8.9 sp.....	1
2030	678	II. 639		10 5 31.3	3.592	1:	50 33 28.5	17.65	1:	eB; cS; R; psbM; r.....	2
2031	} 678, a		R. 2 novæ	10 5 ±		...	50 33 ±		...	3 seen; one (? which) = h. 678	0
2032				10 5 42.7	5.424	1?	14 53 50.7	17.69	1:	vF; S; R.....	1
2033	676			10 5 43.5	3.345	2	66 34 30.8	17.66	3	pF; cL; R; vglbM; r; S * inv	5
2034	682	II. 43		10 5 48.2	3.589	1:	50 40 11.1	17.67	1	F; S; R; gbM.....	2
2035	681	II. 640		10 5 49.2	4.054	1	32 38 47.1	17.67	1	eF; S; R; vglbM.....	1
2036	679			10 6 15.4	3.116	::	85 56 15.7	17.69	::	vvF; mE 0°±.....	0
2037	684, a		R. nova	10 6 29.6	3.116	3	85 52 53.7	17.69	4	B; pS; R; psmbM; p of 2...	8
2038	684	I. 3		10 6 36	3.34	[1]	59 42 7	17.69	[1]	F; S; ?? Cl of vS st.....	0
2039			D'Arrest, 68	10 6 45.3	4.194	1	29 4 36.3	17.71	1	F; psbM; stellar; * 7.8 np 5'	1
2040	683			10 6 58.5	3.116	3	85 50 20.3	17.71	4	B; pL; vLE; pgmbM; * 11, 78° 2, 80°.	8
2041	685	I. 4		10 7 32.0	3.745	1	42 42 39.2	17.74	1	F; S; R.....	1
2042	686			10 8 7.9	+87.502	1::	0 6 46.2	+19.47	1::	vF; R; gbM; * 11, 2' s; Po-	1*
2043	250									larissima Borealis.	

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
2044	3235			10 8 9.4	+2.754	1	116 59 43.5	+17.75	1	eF; S; R; 2 B st f	1
2045		III. 964		10 8 16.1	5.415	1	14 38 38.0	17.80	1	cF; S; stellar; S * f nr	1
2046	3236			10 8 18.9	2.738	2	118 11 3.8	17.76	2	cB; L; mE 50°.5; vglbM ...	2
2047	687	III. 25		10 8 51.5	3.318	3	68 10 42.7	17.79	3	cF; S; R; psBM.....	4
2048	3237			10 9 23.5	2.905	1	105 5 56.0	17.80	1	pB; pL; gpmbM.....	1
2049	688, a		R. nova	10 9 44.5	3.623	...	47 53 13.9	17.83	...	MS; no description.....	0
2050	688, b		R. nova	10 9 47.2	3.623	...	47 54 1.9	17.83	...	MS; no description.....	0
2051		I. 265		10 9 47.5	4.082	1	31 5 44.2	17.84	1	cB; cL; iR; vgbM	1
2052	688	I. 168		10 9 49.9	3.623	1	47 53 1.9	17.83	1	pB; vL; R; vgbM	4
2053	689			10 9 53.3	3.627	1	47 41 6.2	17.84	1	pF; vL; R; vgbM; 12°.5 d; * 11 n 2'.	1
2054	692, a		D'Arrest, 69	10 9 57	3.32	[2]	67 35 48	17.83	[2]	pF; pL; gmbM (δ in Lord R.'s diagram).	0
2055	692, b		R. nova	10 10 4.7	3.324	...	67 25 19.5	17.85	...	Marked γ in Lord R.'s diagr.	0
2056	690	III. 910		10 10 17.0	4.047	1	31 53 7.8	17.86	1	vF; pL; r	2
2057	692, c		R. nova	10 10		...	67 28		...	mE, parallel to h. 692, with which it forms D neb.	0
2058	692	II. 44		10 10 23.3	3.324	3	67 28 10.5	17.85	3	B; pS; E; psbMN; sp of 2	4+
2059	691			10 10 25.8	3.726	1	42 51 47.8	17.86	1	F; S; R; bM	1
2060		III. 704		10 10 27.6	3.728	1	42 43 44.8	17.86	1	eF; vS; (?)	1
2061	693	II. 45		10 10 41.6	3.324	3	67 24 24.8	17.86	3	B; S; vLE; psbM; r; * 9, 352°.0, 75" nf of 2.	4
2062		III. 695		10 10 46.8	+5.391	1	14 29 43.0	17.90	1	vF; vS.....	1
2063	3241			10 10 51.5	-0.506	2	170 10 11.6	17.82	2	!; O; pB; S; lE; 13°.0 d; 3S st nr.	2+
2064	694	III. 348		10 10 53.6	+3.398	1:	61 38 27.1	17.87	1:	eeF; pS; lE.....	2
2065		III. 966		10 10 57.9	6.114	1	11 4 53.6	17.92	1	vF; vS.....	1
2066	695	I. 199		10 11 14.4	3.702	1	43 44 15.7	17.89	1	pB; vL; mE 45°±; vgbM...	3
2067	3239			10 11 46.1	2.128	4	147 15 39.7	17.89	4	!; vB; vL; falcate; * N.....	4+
2068	3238		Δ . 445	10 11 52.8	2.452	2	135 42 6.0	17.90	2	$\oplus$; vL; iR; lCM; gbM; st 13...16.	2
2069	696	II. 720		10 12 5.3	3.643	1::	46 18 42.6	17.92	1:	cF; S; R; vgbM; 1st of 3...	2
2070	3240			10 12 5.5	2.777	2	116 0 17.3	17.91	2	pB; S; cE; gbM.....	2
2071	698			10 12 12.9	3.396	1	61 28 54.6	17.92	1	eF; pL; gbM	1
2072	699	II. 721		10 12 21.7	3.641	1	46 20 50.2	17.94	1	cF; S; R; vgbM; 2nd of 3...	2
2073	697	I. 266		10 12 30.9	4.011	1	32 22 5.2	17.94	1	pB; cL; E; vglbM	2
2074	700	II. 722		10 12 33.3	3.641	1	46 18 50.2	17.94	1	cF; S; R; stellar; 3rd of 3...	2
2075	701			10 12 47.0	3.365	1	63 47 46.5	17.95	1	F; S; R; has a *	1
2076	3242			10 13 17.6	1.946	2	151 58 35.5	17.95	2	O; =* 10 m; R; am 150 st	2
2077		III. 979	H. ON	10 13 34.0	6.559	1	9 27 5.6	18.02	1	Stellar; 1st of 3	1
2078		III. 980	H. ON	10 13 34.2	6.565	1	9 26 5.6	18.02	1	vF; S; 2nd of 3	1
2079		III. 981	H. ON	10 13 34.3	6.571	1	9 25 5.9	18.03	1	vF; S; 3rd of 3	1
2080	702	III. 330		10 13 55.0	3.342	1	65 21 53.7	17.99	1	vF; pS; R; bM	2
2081		I. 283		10 14 23.0	5.297	1	14 37 52.9	18.03	1	cB; cL; cR	1
2082		III. 911		10 14 26.4	3.998	1	32 15 55.6	18.02	1	vF; cL; iF	1
2083			D'Arrest, 70	10 14 39	3.31	[2]	67 42 12	18.01	[2]	eF; mE; a ray	0
2084			Auw. N. 27	10 14 55.0	3.288	...	69 24 42.6	18.02	...	F; lbMr (Winnecke, June 1855).	0
2085	3243			10 15 20.5	2.683	1	123 32 59.9	18.03	1	pB; vL; vLE; psbMN	1
2086	3244			10 15 27.7	2.677	1	123 59 28.2	18.04	1	vF; pS; R; vgbM	1
2087	703	II. 882		10 15 43.0	4.028	1	31 9 19.1	18.07	1	cF; pL; lE; vgbM	2
2088		II. 28		10 15 55.1	3.289	1	69 23 27.8	18.06	1	vF; cL; R } D neb; 45°, 2' {	1*
2089		II. 29		10 15 58.5	3.289	1	69 22 28.1	18.07	1	vF; cL; R }	1*
2090	3245		Δ . 386	10 16 10.6	2.352	1::	141 1 1.8	18.06	1::	Cl; 9 L & a few S st	1
5068				10 16 14.1		...	89 13 46.0		...	See No. 5068.	
2091	705			10 16 15.7	3.207	3	76 44 21.4	18.08	3	!; * or ** in neb	3
2092	704			10 16 25.6	4.456	1	22 29 6.0	18.10	1	Cl; cL; P; lC; st 10...12 ...	1
2093			D'Arrest, 71	10 16 33	3.39	[1]	61 16 42	18.09	[1]	eeF; * 11 p, ls, 150" p of 2...	0
2094	706			10 17 5.1	3.374	1	62 16 5.0	18.10	1	pB; pS; R; psbM	1*
2095			D'Arrest, 72	10 17 7	3.38	[2]	61 16 12	18.11	[2]	F; S; f of 2	0
2096	707			10 17 16.6	+4.142	1	28 1 16.9	+18.13	1	eF; vS; psbM; 2 st 11, 12, f	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
2097	h. 709	H. III. 631		h m s 10 17 27.7	s +3.556	1:	49° 39' 49.9"	+18.13	1:	vF; vS; R; pgbM	3
2098	708	III. 883		10 17 28.5	3.980	1	32 4 25.9	18.13	1	F; S; R; psbM	2
2099	$\left\{ \begin{array}{l} 710 \\ = \\ 3246 \end{array} \right\}$	IV. 10		10 17 31.0	3.255	3	72 8 18.9	18.13	3	vF; * 9 inv nr M	5
2100	3247			10 17 53.2	2.851	2	111 4 56.9	18.13	2	eF; S; R; * nr	2
2101	3249			10 17 56.8	2.717	1	121 45 28.9	18.13	1	F; pmE; glbM; * 11 np ...	1
2102	3248	IV. 27	Lal. 20204	10 18 2.2	2.886	4	107 55 50.2	18.14	4	!; O; vB; IE, 135°; 32'' d $\pm$; blue.	7+
2103	4019			10 19 20.9	2.612	1	129 6 23.4	18.18	1	vF; * 11 m 90'' n	1
2104	711	I. 86		10 19 24.9	3.385	4	60 47 1.7	18.19	5	vB; pL; E; smbMEN	6
2105	712			10 19 26.0	3.116	1	85 26 24.7	18.19	2	eF; S; R; 2 st Δ ; * 6, 300°, 8'	3
2106	3250			10 19 37.9	2.196	1::	147 10 35.7	18.19	1	st inv in neb	3
2107	713	II. 347		10 20 2.0	3.314	1	66 26 28.6	18.22	3	pB; S; R; psbM	4
2108	3251			10 20 6.9	2.690	1	124 14 38.3	18.21	1	eF; pL; R; vglbM	1
2109	3252			10 20 25.6	2.625	4	129 13 45.9	18.23	4	pB; pL; R; vg, psbM; * 13, 45°.	5
2110			D'Arrest, 73	10 20 29	3.35	[1]	63 11 42	18.23	[1]	vF; pL; 3 B st sp	0
2111		III. 316		10 20 51.1	5.054	1	15 27 11.9	18.27	1	eF; pS; mE; r	1*
2112	714	I. 72		10 21 27.2	3.392	2	59 47 27.1	18.27	3	cB; L; E 45° $\pm$; psmbMN...	5
2113	3253			10 21 32.2	2.111	1	149 57 49.8	18.26	1	Cl; pS; vC; st 15	1
2114	3254			10 21 53.0	2.553	1	133 11 7.1	18.27	1	cB; S; R; gmbM	2
2115	3255			10 22 29.8	2.689	3	124 56 49.0	18.30	2	vF; vS; R; psbM; 1st of 4...	3
2116	3256			10 22 36.9	2.689	4	124 53 9.0	18.30	4	eF; S; R; psbM; 2nd of 4...	4
2117	715	II. 870		10 22 42.9	4.272	1	24 14 50.9	18.33	1	F; S; R; gbM	2
2118	3257			10 22 52.7	2.689	4	124 52 39.0	18.32	4	vvF; vS; R; psbM; 3rd of 4	4
2119	3258			10 23 3.0	2.546	1	133 56 12.6	18.32	1	F; S; R; am st	1
2120	3260			10 23 8.2	2.554	1	133 29 7.6	18.32	1	eF; S; R	1
2121	3259			10 23 11.4	2.692	2	124 53 48.9	18.33	2	vF; vS; R; psbM; 4th of 4	2
2122	3261			10 23 13.8	2.556	1	133 24 32.9	18.33	1	F; S; mE 280° $\pm$; psbM ...	1
2123	718	III. 349		10 23 15.9	3.377	1	60 29 34.9	18.33	1	pF; S; R; psbM; * sf nr ...	5
2124	716			10 23 18.3	3.895	1	33 11 37.5	18.35	1	eF; bet 2 S st	1
2125	717	II. 871		10 23 29.5	4.246	1	24 32 16.8	18.36	1	vF; vS; R; psmbM *	2
2126	3262			10 23 33.9	2.696	1::	124 38 10.2	18.34	1::	eF; vS; R; 1st of 4	1
2127	3263			10 23 37.9	2.697	2?	124 39 10.2	18.34	2?	F; S; R; 2nd of 4	2
2128	3264			10 23 41.1	2.699	1	124 30 10.5	18.35	2	F; S; R; bM; 3rd of 4	3
2129	719	III. 331		10 23 44.1	3.329	1	64 24 28.5	18.35	1	eF; vS; R; glbM	2
2130	3265			10 23 45.9	2.697	1	124 39 3.5	18.35	1	pF; S; E; pmbM; 4th of 4...	2
2131	720	II. 358		10 24 27.7	3.360	4	61 36 59.4	18.38	4	F; pL; R; glbM; * f	5
2132	3266			10 24 34.0	2.681	2	126 1 22.1	18.37	2	F; L; vLE; psbM	3
2133	3267			10 24 57.7	2.635	1	129 13 51.7	18.39	1	F; S; * 8 p	1
2134	721	II. 359		10 25 4.1	3.369	3	60 46 20.7	18.39	3	cB; cS; R; pgmbM	4
2135	3268			10 25 26.1	2.637	3	129 13 52.0	18.40	3	F; S; R; * nf	2
2136	3269			10 25 34.3	2.711	1	124 8 4.3	18.41	1	eF; pL; E; glbM	1
2137	3271			10 26 51.1	2.536	1	135 22 3.5	18.45	1	pF; S; R; gbM	1
2138		III. 912		10 26 58.6	3.944	1	30 44 23.1	18.47	1	eF; vS	1
2139	3270			10 27 1.3	2.808	4	116 44 11.8	18.46	4	pB; S; IE; gbM; 1st of 9 ...	4
2140	722	III. 917		10 27 9.3	3.945	1	30 40 11.4	18.45	1	vF; pS; R; psbM	2
2141			D'Arrest, 74	10 27 11	3.28	[2]	67 37 18	18.47	[2]	F; pL; * p 24°, 225'' s	0
2142	723	III. 918		10 27 12.7	3.942	1	30 43 31.4	18.48	1	eF; cS; R; vglbM	2
2143	3272			10 27 47.4	2.713	1	124 34 31.4	18.48	1	eF; vS; R	1
2144	3276			10 28 7.7	2.258	1::	147 28 14.7	18.49	1::	Cl; B; Ri; pL	1*
2145	724	I. 164		10 28 12.8	3.474	1	51 57 17.3	18.51	1	cB; L; mE 135° $\pm$; glbM ...	4
2146	725	III. 767		10 28 44.6	3.698	1	39 9 51.9	18.53	1	vF; pS; IE	2
2147	726	III. 54		10 28 57.8	3.193	1	76 34 38.9	18.53	1	eF; cL; R; vgbM; r	2
2148	$\left\{ \begin{array}{l} 727 \\ = \\ 3273 \end{array} \right\}$	III. 55		10 29 10.6	3.206	3	75 6 29.2	18.54	3	cF; cS; R; pmbM; r; am B st	5
2149		II. 46		10 29 13.4	3.285	1	67 15 27.2	18.54	1	pF; S; r; Δ pB st n	1
2150	728	II. 46??		10 29 18.1	3.283	2	67 23 45.2	18.54	2	cB; S; IE; psbM; r	3
2151	3274			10 29 20.3	+2.757	1	121 37 4.2	+18.54	1	eF; S; R	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	a						
2152	3275			10 29 22.6	+2.756	1	121° 38' 41".2	+18.54	1	vF; S; R	1
2153	730	III. 66		10 29 27.1	3.245	2	71 8 48.2	18.54	2	vF; vS; vIE; glbM; r	3
2154	729	III. 615		10 29 33.9	3.470	2	51 49 25.5	18.55	2	vF; cS; psbM; er	4
2155	3277			10 29 38.5	2.818	1	116 26 17.5	18.55	1	vF; S; R; 2nd of 9	1
2156	3278			10 29 41.3	2.813	1	116 52 38.5	18.55	1	eeF; 3rd of 9	1
2157	3279			10 29 48.4	2.816	4	116 42 50.5	18.55	4	F; S; R; 4th of 9	4
2158	731	IV. 60		10 29 59.7	3.770	1	35 45 49.1	18.55	1	○? cB; pL; R; vg, vsmbMN 15"	3†
2159	3280			10 30 1.1	2.815	2	116 48 5.8	18.56	2	B; L; R; p of D neb; 5th of 9	3
2160	3281			10 30 9.5	2.815	1::	116 49 17.1	18.57	1::	B; L; R; f of D neb; 6th of 9	1
2161	3282			10 30 28.7	2.815	2	116 53 42.1	18.57	2	cF; E; gbM; 7th of 9	2
2162	3283			10 30 39.9	2.815	1	116 56 48.4	18.58	1	8th of 9	1
2163	3284			10 30 59.8	2.817	3	116 52 45.7	18.59	3	F; S; R; bM; 9th of 9	3
2164	3285			10 31 6.1	2.636	2	130 54 24.7	18.59	2	cF; pL; pmE; lbM	2
2165		III. 700		10 31 7.4	3.526	1	47 36 30.3	18.61	1	cF; L; iE; mb, s of M	1
2166	732	II. 745		10 31 12.8	3.626	2	41 51 29.3	18.61	2	F; pS; mE 0...90°; * 10 nf.	4
2167	3286		Δ. 322?	10 32 2.9	2.277	1	147 53 48.6	18.62	1	pB; vL; iF; * inv	1
2168	734	II. 348		10 32 17.5	3.299	2	65 11 2.2	18.64	2	vF; S; R; gbM; vS * att	3
2169	733			10 32 34.7	5.264	3	12 26 32.4	18.68	3	pB; S; iE; psmbM	3
2170		I. 272		10 32 34.7	3.158	1::	79 59 19.5	18.65	1::	B; S; iR; mbMBN	2
2171	3287		Δ. 355	10 33 1.6	2.409	1	143 24 4.5	18.65	1	Cl; P; st 9	1
2172	3288			10 33 26.3	2.729	1	125 19 17.0	18.70	1	cF; vS; mE; * 15 att	1
2173	735	II. 641		10 33 28.5	3.451	1	51 58 16.4	18.68	1	eF; vS; R; bM	4
2174	3289			10 33 43.6	2.823	1	117 1 41.4	18.68	1	vF; pL; iE; glbM	1
2175	737	II. 77		10 34 42.7	3.195	2	75 31 44.6	18.72	2	F; cL; E; vgbM; r; * 7p10	4
2176	736	III. 317		10 35 17.6	4.724	1	15 54 50.5	18.75	1	pF; S; R; gbM	2
2177		III. 5		10 35 25.5	3.157	1	79 49 40.2	18.74	1	cF; eS	1
2178	739	I. 81		10 35 49.5	3.298	1	64 20 20.5	18.75	1	eB; L; gbM; * inv; 2st f	3
2179	740	I. 26		10 36 8.6	3.178	1	77 16 41.8	18.76	1	cB; pL; E; mbM	2
2180	3290	V. 7		10 36 10.9	3.203	1	74 23 11.8	18.76	1	cF; vL; R; vglbM; er	2
2181	3291			10 36 23.8	2.732	4	125 37 46.8	18.76	3	pF; S; mE 0°±; vsmbM; 1st of 3.	4
2182	738	I. 80		10 36 29.0	4.648	1	16 25 22.7	18.79	1	B; S; iE; psbM; * 11, 281° 8, 20° 0.	2
2183	742			10 36 33.7	3.358	1	58 32 21.1	18.77	1	eF; vS; 2st 9-10, s	1
2184	743		M. 95	10 36 36.7	3.175	2	77 34 22.1	18.77	4	B; L; R; pgmbMrN	8
2185	741	III. 842		10 36 39.2	3.732	1	33 18 30.6	18.78	1	F; cS; R; psbM; * 90"	2
2186	3292			10 36 41.4	2.734	4	125 38 53.1	18.77	3	F; S; vIE; psbM; 2nd of 3...	4
2187	744	III. 107		10 36 53.2	3.133	2	82 30 48.4	18.78	2	vF; pS; R; bM; * 9, 150"	4
2188	3293			10 37 11.7	2.735	...	125 38 59.7	18.79	...	cF; vS; vIE; vS * att; 3rd of 3	2
2189	745	V. 52		10 37 21.0	4.018	1	26 2 25.3	18.81	1	pB; L; E 0°; glbM	2*
2190	746	III. 318		10 37 57.1	4.583	1	16 49 37.9	18.83	1	vF; L; R; vgbM; r; * sf	2
2191	747			10 38 59.5	3.091	1	87 28 32.5	18.85	1	eF; L; eE; vglbM; a ray...	1
2192	3294			10 39 4.1	2.644	1?	132 58 43.5	18.85	1	F; E; gbM; * 6, 7 v nr	1*
2193	748	II. 78		10 39 10.5	3.189	2	75 31 1.5	18.85	2	pB; cL; iR; vglbM; r; 1st of 3	5
2194	749		M. 96	10 39 20.4	3.173	4	77 26 55.8	18.86	4	vB; vL; iE; vsvmbM; r	8
2195	750	II. 81		10 39 34.8	3.219	2	71 59 40.1	18.87	2	cB; pL; vIE; gbM; r	3
2196	751			10 39 37.3	3.188	1	75 28 25.1	18.87	1	F; R; 2nd of 3	1
2197	3295		Δ. 309	10 39 36.8	2.313	3	148 56 44.8	18.86	3	η Argūs. The great neb	Mon.†
2198	753			10 39 49.3	3.187	1	75 35 15.1	18.87	1	F; R; 3rd of 3	1
2199	752	III. 701		10 39 49.4	3.506	1	46 4 18.4	18.88	1	vF; cS; iR	2
2200			D'Arrest, 75	10 40 12	3.12	[1]	83 12 24	18.88	[1]	vF; S	0
2201	754	II. 99		10 40 16.7	3.189	2	75 16 24.7	18.89	2	vB; cL; R; svmbMBN	5*
2202	3296			10 40 22.4	2.702	3	129 17 2.7	18.89	3	cF; S; R; glbM	4
2203	757	I. 17	Mechain.	10 40 25.7	3.177	3	76 41 10.7	18.89	3	vB; cL; R; psbM; r	8
2204	755	II. 360		10 40 27.4	3.323	5	60 39 47.7	18.89	5	pB; pS; R; sbM	6
2205	756	II. 565		10 40 33.2	3.390	2	54 33 18.7	18.89	2	pF; cL; iR; vglbM; 1st of 3	3
2206	3297			10 40 39.5	2.874	2	113 41 36.7	18.89	2	F; pL; iR; glbM	2
2207	758	I. 18		10 40 52.2	3.177	3	76 38 27.0	18.90	3	vB; L; R; psmbM; 2nd of 3	8
2208	759			10 40 54.9	3.116	1	84 15 58.0	18.90	1	vF; R	1
2209	760			10 40 55.7	+3.116	2	84 20 22.0	+18.90	2	F; S; iE; bM	2

No. of Cata- logue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	"			
2210	762			10 41 2.5	+3.116	1	84 18 1.3	+18.91	1	Suspected; *nr	1
2211	761	II. 41		10 41 3.8	3.176	2	76 43 38.3	18.91	2	F; L; E 90°±; vglbM; 3rd of 3.	9
2212	3298			10 41 32.5	2.807	1	120 48 44.6	18.92	1	F; S; pmE 0°	1
2213	763	III. 881		10 41 36.0	4.082	1	23 28 40.9	18.93	1	vF; S; psbM; st nr	2
2214	3299			10 41 41.8	2.870	3	114 25 38.6	18.92	3	F; S; R; psbM; 2st 10f	3
2215	764	II. 872		10 41 59.8	4.075	1	23 30 51.2	18.94	1	cF; S; IE; vgbM	2
2216	765	I. 116		10 42 2.4	3.364	2	56 16 38.2	18.94	2	cB; pS; iE; 1st of 2	4†
2217	766	I. 117		10 42 7.5	3.364	2	56 15 59.2	18.94	2	pB; pS; iE; 2nd of 2	4†
2218		I. 284		10 42 34.7	5.103	1	11 57 49.1	18.97	1	cB; vS; iF	1
2219		III. 792		10 42 57.4	3.717	1	33 49 54.1	18.97	1	vF; S; E; er	1
2220	768	II. 361		10 43 2.3	3.314	3	60 47 45.1	18.97	3	pF; S; R; bM	4
2221	767	II. 335		10 43 15.4	4.609	1	15 34 35.7	18.97	1	pF; L; iE; vgbM	2
2222	771			10 43 17.6	3.619	1	38 13 52.4	18.98	1	pB; R; pgbM	1
2223	769	III. 919		10 43 20.3	3.884	1	27 53 7.4	18.98	1	vF; vS; R; vS* nr	2
2224	770	III. 913		10 43 21.3	3.792	1	30 49 46.4	18.98	1	vF; cS; R; 2pB st s	2
2225	772	II. 718		10 43 33.3	3.496	1	45 32 51.4	18.98	1	pB; S; vIE; stellar; 3S st nr.	3
2226	772, a		R. nova	10 43 +		...	45 +		...	3' dist. from h. 772	0
2227	773	II. 362		10 43 33.5	3.307	4	61 17 10.4	18.98	4	B; pL; R; mbM	5
2228	776	III. 522		10 43 28.5	2.982	(3)	102 6 30.0	19.00	1:	F; S; R; lbM	4
2229	774	I. 27		10 43 29.8	3.180	2	75 50 51.4	18.98	2	B; S; IE 135°±; smbMN...	5
2230	775	II. 363		10 43 41.7	3.308	2	61 9 14.7	18.99	4	cF; S; R; bM	5
2231		IV. 6		10 43 49.0	3.117	4	83 26 44.4	18.99	4	B; vL; R; bM; r	5*
2232		II. 131		10 43 52.8	3.355	1	56 29 54.7	18.99	1	F; S	1
2233		I. 118		10 43 54.7	3.327	1	57 16 54.7	18.99	1	cB; cL; iR; mbM (758° P.D.).	1*
2234	777	III. 88		10 43 57.2	3.121	3	83 25 0.7	18.99	3	F; vL; R; vgbM; rr	5*
2235	778	II. 494		10 44 1.8	3.356	1	56 21 52.0	19.00	1	pF; pL; IE; sp of 3	3
2236	779	I. 118?		10 44 27.8	3.355	1	56 18 10.3	19.01	1	pB; L; iE; gbM; 2nd of 3...	2*
2237		III. 108		10 44 33.7	3.141	1	80 44 57.3	19.01	1	eF; eS; R	1
2238	780	I. 172		10 44 35.5	3.395	2	52 38 29.3	19.01	2	pB; pL; vmE 42°5; *inv?	3*
2239	782			10 44 40.7	3.356	1:	56 10 30.3	19.01	1:	pB; nf of 3 in a line	1
2240	783	III. 20		10 44 40.7	3.156	1	79 6 47.3	19.01	1	vF; vL; R; vgbM	2
2241	784	III. 497		10 44 43.3	3.105	1	85 28 16.3	19.01	1	F; pS; R; vglbM	3
2242	781	II. 887		10 44 54.9	3.865	1	27 58 21.6	19.02	1	cF; pS; IE; vgbM	2
2243	786	II. 47		10 44 59.2	3.257	2	66 19 33.6	19.02	3	pB; pL; IE 120°; gbM	5
2244	785	III. 914		10 45 10.9	3.740	1	32 8 27.9	19.03	1	vF; S; IE	2
2245	787	I. 267		10 45 58.4	3.730	1	32 16 18.5	19.05	1	cB; pL; iR; vglbM; *10nf2'	2
2246	3301			10 45 58.5	2.660	2	134 24 21.2	19.04	2	Cl; pL; P; IC; iF; st 9...13	2
2247	3300			10 45 59.6	3.203	1	72 29 30.5	19.05	1	eF; vL; vglvbm; B *sp ...	1
2248	788	I. 233		10 46 5.2	3.666	1	34 56 48.5	19.05	1	B; pL; mE 67°0; gbM	3
2249	3302			10 46 20.2	2.809	1	122 10 58.5	19.05	1	F; S; R; *6.7 sf	1
2250	3303			10 46 25.7	2.919	1	110 6 14.5	19.05	1	vF; L; R; vglbM; r	1
2251	789	II. 364		10 46 40.2	3.291	4	62 1 7.1	19.07	4	F; pL; vIE; vlbM	5
2252	3304			10 46 52.3	2.912	1	111 2 13.1	19.07	1	F; S; R; bM	1
2253	790			10 47 1.0	3.206	1	71 54 33.4	19.08	1	pF; iE; np of 2	1
2254	791	II. 82		10 47 3.0	3.205	1	71 58 7.4	19.08	2	pF; S; E; gbM; r; sf of 2...	3
2255	792	IV. 29		10 47 11.5	2.958	1	105 16 57.4	19.08	1	eF; att to *12f	2
2256	793			10 47 21.7	3.207	1	71 38 54.7	19.09	1	2 or 3 Sst & neb	1
2257		I. 268		10 47 23.7	3.721	1	32 8 1.7	19.09	1	vB; vS; R; stellar	1
2258	794	II. 16		10 48 2.0	3.132	1	81 33 42.0	19.10	1	vF; vS; vIE; psbM	5
2259	3305			10 48 29.5	2.879	1	115 23 58.3	19.11	1	F; S; R; glbM	1
2260	795			10 48 57.1	4.652	2	14 3 47.2	19.14	2	eF; pL; R; vglbM; *nf ...	2
2261	796			10 48 57.6	3.146	1	79 30 48.9	19.13	1	vF; *9, 90°; p of 2	1
2262	798			10 49 26.2	3.145	1	79 29 49.2	19.14	1	vF; R; vsmbM*12; f of 2...	1
2263	797	III. 632		10 49 33.9	3.425	2	48 18 8.2	19.14	2	F; eS; R; bM	4
2264	3306			10 49 58.9	2.976	1	103 33 21.5	19.15	1	eeF; S	1
2265	799	II. 888		10 50 5.6	3.759	1	29 44 47.8	19.16	1	vF; S; R; vgbM	2
2266		III. 972		10 50 15.1	3.819	1	27 39 6.1	19.17	1	vF; S; R; bM	1
2267		III. 67		10 50 35.3	+3.197	1	72 9 7.1	+19.17	1	vF; E; bet 2 st	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. II.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	"			
2268	800	III. 332		10 50 49.8	+3.253	1	65 1 18.4	+19.18	1	vF; R; gbM; *13 {H.1'n h. 2's}	2
2269	801	III. 705		10 51 18.6	3.485	1	43 8 17.7	19.19	1	eF; S; R	2
2270	3308			10 52 16.3	2.675	1	135 49 22.3	19.21	1	eF; S; R; gbM	1
2271	3307			10 52 17.7	2.869	1	117 43 48.3	19.21	1	pF; S; R; bM; am st	1
2272	802			10 52 29.0	4.670	1	13 25 47.9	19.23	1	Very doubtful object	1
2273	{ 804 = 3309 }	II. 100		10 52 37.6	3.178	4	74 24 56.6	19.22	4	F; L; R; glbM; r	5
2274	805	I. 87		10 52 44.4	3.288	10	60 16 20.6	19.22	10	cB; cL; R; gmbM	13
2275	803	I. 269		10 52 50.3	3.686	1	31 34 53.9	19.23	1	{H. eF; } h. cB; } v1E; pS; *13 s att	2
2276	806	II. 101		10 52 56.1	3.170	2	75 20 58.9	19.23	2	vB; pL; 1E 80°+; smbMN.	4*
2277	807	III. 21		10 53 14.5	3.158	1	77 5 16.2	19.24	1	eF; cS; R; bMN	3
2278	808			10 53 49.7	3.274	2	61 32 11.5	19.25	2	vF; R; bM; *sp	2
2279	809	III. 498		10 54 0.3	3.100	3	85 37 30.5	19.25	3	vF; pL; mE	4
2280	3310			10 54 11.7	2.432	1	149 34 46.5	19.25	1	Cl; pL; pRi; lC; st 13	1
2281		III. 824		10 54 29.4	2.945	1	108 43 14.1	19.27	1	vF; vS; iR; glbM	1
2282		III. 75		10 54 30.4	3.171	1	74 52 13.1	19.27	1	eF; pL	1
2283		III. 793		10 54 45.6	3.636	1	33 2 13.4	19.28	1	vF; vS; stellar	1
2284	{ 967 968 }	III. { 967 968 }		10 55 9.9	4.582	1	13 27 12.0	19.30	1	{vF eF}; D neb; v near	1
2285											
2286	3311			10 55 32.8	2.457	1	149 5 48.7	19.29	1	3S st 10 m in vF neb	1
2287	810	I. 88		10 55 33.8	3.271	5	61 16 22.7	19.29	6	B; L; E; mbMN; rr; p of 2	8
2288	3312			10 55 50.2	2.976	1	104 44 26.0	19.30	1	pF; S; R; glbM; *14 nr	1
2289	811	III. 22		10 55 53.7	3.147	3	78 10 23.0	19.30	3	vF; cS; R; vglbM	4
2290	812	IV. 7		10 55 59.3	3.195	2	71 6 42.0	19.30	2	eF; pL; R; sbMS*; *9 att 25°.	4
2291	814	II. 507		10 56 3.9	2.970	1	105 32 3.0	19.30	1	F; {H. S h. vL}; bM; *nfinv...	2
2292		III. 598		10 56 5.4	3.106	1	84 28 16.3	19.31	1	eF; S; 1E; ?	1
2293	813	II. 365		10 56 6.3	3.277	2	60 21 55.3	19.31	2	F; L; cE; *7, 310° 8'	5
2294		V. 39		10 56 27.4	2.923	2	112 21 17.3	19.31	2	vF; vL; mE	2
2295	815	II. 366		10 56 28.0	3.269	1	61 11 48.3	19.31	1	F; pS; R; pgbM; f of 2	2
2296		V. 40		10 56 43.8	2.923	2	112 29 17.6	19.32	2	vF; vL; mE	2
2297	3313			10 56 50.2	2.955	1	108 1 41.6	19.32	1	vF; pL; R; vglbM	1
2298	816	II. 336		10 57 1.2	4.248	1	16 40 22.2	19.34	1	pB; vS; iR; psmbM*	2
2299	817	II. 884		10 57 13.1	3.622	1	32 43 27.2	19.34	1	eF; S; R; vglbM	2
2300	3314			10 58 19.3	2.446	1	150 36 38.5	19.35	1	Cl; pRi; pC	1
2301	818	I. 13		10 58 38.7	3.076	1	89 16 57.8	19.36	1	cB; cL; mE 140°±; vsmbMN	4
2302		II. 904		10 59 4.5	4.481	1	13 33 18.7	19.39	1	F; pL; lbM	1
2303	819	III. 23		10 59 13.5	3.145	2	77 51 22.4	19.38	3	F; S; 1E; psbM; 2st np in line.	5
2304	820	III. 350		10 59 41.3	3.263	3	60 43 33.7	19.39	3	eF; S; *10 p 60''	5
2305	3316			11 0 17.4	2.956	1	108 43 4.0	19.40	1	F; S; R; psbM; p of 2	1
2306	3317			11 0 19.9	2.956	1	108 47 19.0	19.40	1	eF; S; R; vlbM; f of 2	1
2307	821	III. 915		11 0 20.6	3.609	1	32 1 29.0	19.40	1	vF; S; R; pgbM	2
2308	3315		Δ. 323	11 0 33.8	2.530	2	147 55 1.0	19.40	2	II; Cl; eL; R; lC; st 8...12...	4
2309	3318			11 0 45.4	2.823	1	126 25 0.3	19.41	1	eeF; vSatt	1
2310	823	III. 111		11 1 17.1	3.104	1	84 24 45.6	19.42	1	eF; vS; R; bM; r	2*
2311	822			11 1 18.9	3.260	1	60 47 14.0	19.40	1	F; S; R; bM	1
2312	825			11 1 31.1	3.258	1	60 34 29.9	19.43	1	eF	1
2313	824			11 1 32.2	3.319	1	53 12 44.9	19.43	1	vF; R; psbM; *7 p 7'	1
2314	826	III. 920		11 2 29.3	3.683	1	27 54 1.5	19.45	1	eF; vS; E 0°±; r	2
2315	828	II. 42		11 2 38.9	3.137	1	78 30 58.5	19.45	1	F; S; 1E; vlbM	2*
2316	827			11 2 40.8	3.314	1	53 12 35.5	19.45	1	eF; S; *8, p	1
2317		I. 220		11 2 44.9	3.518	2	35 51 45.8	19.46	2	cB; cL; cE 160°	2
2318	831	V. 46		11 3 3.0	3.554	1	33 34 44.1	19.47	1	cB; vL; vmE 79°0; pbM; r.	3
2319	829	III. 351		11 3 3.7	3.254	5	60 28 50.8	19.46	6	!; F(? var); S; R; bM; *9 f1'; 1st of 4.	8*
2320	832	III. 352		11 3 9.6	+3.253	3	60 33 4.1	+19.47	3	eF; vS; 2nd of 4	5

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
2321	833			11 3 12.0	+3.253	1	60 36 21.1	+19.47	2	vF; pS; R; bM; 3rd of 4 ...	2
2322	3319			11 3 19.8	2.830	2	126 46 48.1	19.47	2	B; S; R; pgmbM; 1st of 3...	2
2323		III. 79		11 3 25.7	3.144	1	77 19 26.1	19.47	1	eF; pS; lE; r	1
2324	834			11 3 26.1	3.139	1	78 3 34.1	19.47	1	F; S; R; gbM	1
2325	830	II. 337		11 3 30.4	4.167	1	16 22 0.4	19.48	1	pF; pS; lE; gbM; *15, 22° 1, 70 ⁿ .	2
2326	835			11 3 36.1	3.252	3	60 32 50.1	19.47	3	vF; pL; 4th of 4	3
2327	3320			11 3 56.5	2.832	2	126 46 43.4	19.48	2	pF; S; R; bM; 2nd of 3.....	2
2328	836	III. 89		11 4 2.4	3.108	2	83 25 3.4	19.48	2	eF; R; sbM; r	3
2329	3321			11 4 9.7	2.835	1	126 42 10.4	19.48	2	vF; pL; R; * inv; 3B st nr...	2
2330		II. 819		11 4 32.1	2.972	1	107 31 27.7	19.49	1	pF; pL; iF; bM	1
2331	3323			11 4 32.5	2.534	1	149 28 52.7	19.49	1	Cl; pRi; lC	1
2332	3322			11 4 39.9	2.842	1	126 5 31.7	19.49	1	eF; S; R; glbM; 3 st 11 f ...	1
2333	3324			11 5 52.5	2.522	1	150 36 57.3	19.51	1	F; lE; 1st of 6	1+
2334		III. 723		11 5 53.4	3.424	1	40 53 27.6	19.52	1	eF; vS; p of 2	1
2335	837			11 5 54.4	2.988	1	105 11 45.6	19.52	1	Neb (?)	1
2336	3325			11 5 56.2	2.526	2	150 28 3.6	19.52	2	F; lE; sbM; 2nd of 6.....	2+
2337	3326			11 6 3.2	2.526	2	150 32 58.6	19.52	2	*12 with fan-shaped neb att; 3rd of 6.	2+
2338	3327			11 6 14.4	2.528	2	150 30 34.6	19.52	2	B; bM*; 4th of 6	2+
2339		II. 728		11 6 18.3	3.422	2	40 51 28.9	19.53	2	pB; pL; R; vgmbM	2
2340	3329			11 6 21.4	2.541	2	150 26 42.6	19.52	2	F; L; E 0°; bM; 5th of 6 ...	2+
2341	3328	II. 269		11 6 28.1	2.923	2	115 59 51.9	19.53	2	B; pL; E; vsmbMN; 2 Bst Δ	4
2342	3330			11 6 31.0	2.540	2	150 35 29.9	19.53	2	cF; S; E 160° ±; 6th of 6 ...	2+
2343	838		M. 97	11 6 34.8	3.514	1	34 13 38.2	19.54	1	!!; O; vB; vL; R; vvg, vsbM O; 19° 0 d.	4+
2344	839	III. 921		11 6 51.9	3.621	1	28 32 29.2	19.54	1	vF; L; E; vgbM; in Δ of L st	2
2345	3332			11 6 59.1	2.546	1	150 2 19.2	19.54	1	Cl; pRi; C; E.....	1
2346	3331	III. 529		11 7 4.5	3.000	1	103 19 48.2	19.54	1	vF; S; iR; lbM	3
2347	840	I. 29		11 7 16.1	3.144	3	76 25 29.5	19.55	3	B; cL; E 90° ±; psmbM.....	4
2348		III. 770		11 7 26.0	3.520	1	33 29 29.5	19.55	1	vF; vS; stellar	1
2349		III. 706		11 7 30.1	3.404	2	41 45 30.5	19.55	2	vF; vS; v lE; stellar; cB* n...	2
2350	841	II. 102		11 7 43.8	3.154	1	74 26 53.8	19.56	2	pF; L; R; glbM	2
2351	3333			11 7 49.1	2.946	1	112 57 42.8	19.56	1	vF; pS; R; bM	1
2352	843	II. 49		11 8 4.2	3.172	1	71 7 42.8	19.56	1	B; pS; R; pgmbM	2
2353	842	II. 709		11 8 6.6	3.339	1	47 38 28.8	19.56	1	pF; S; lE 0° ±; vgbM	4
2354	3334			11 9 7.0	2.558	1	150 29 39.4	19.58	1	⊕ and neb; st 15...18	2
2355		II. 626		11 9 7.9	3.098	1	84 42 34.4	19.58	1	pB; S; lE; mbM.....	1
2356	844	III. 27		11 9 21.8	3.169	2	71 14 8.7	19.59	2	F; S; R; sp of 3	3
2357	3335			11 9 29.9	2.885	1	123 3 54.7	19.59	1	eF; S; R; gbM	1
2358	845	II. 50		11 9 30.6	3.169	2	71 11 38.7	19.59	2	vB; L; R; vmbM; 2nd of 3..	3
2359	846	II. 51		11 9 35.3	3.169	2	71 5 49.7	19.59	2	B; pL; R; psbM; 3rd of 3...	3
2360	847	I. 270		11 10 11.5	3.548	3	30 27 29.0	19.60	3	vB; pS; lE 90° ±; vsymbMSN.	5
2361	849	II. 521		11 10 16.1	3.098	2	84 40 50.0	19.60	2	pF; cS; iR; psmbM; *10, 330°, 3'.	5
2362	848	I. 271		11 10 20.9	3.533	1	31 14 40.3	19.61	1	vB; cL; mE 305° 0; smbMN	2
2363	850	II. 729		11 10 29.9	3.366	1	43 29 5.3	19.61	1	F; pL; lE 90° ±; glbM; r ...	3
2364	851	III. 333		11 10 39.5	3.197	1	65 49 46.3	19.61	1	cF; vS; smbM; stellar; p of 2	2
2365	851, α		R. nova	11 10 +		...	65 49 +		...	F; S; bM; place from MS ...	0
2366		III. 76		11 10 53.0	3.149	1	74 29 36.6	19.62	2	eF; pL.....	1
2367	3336			11 10 59.5	2.939	1	115 21 55.6	19.62	1	F; S; R; gbM.....	1
2368		III. 334		11 11 0.0	3.197	1	65 42 34.6	19.62	1	vF; S; f of 2	1
2369	852	I. 244		11 11 9.6	3.520	1	31 30 10.6	19.62	1	cB; cL; R; vgmbM	3
2370	3338			11 11 21.4	1.992	1	165 27 17.6	19.62	1	F; pS; pmE; gbM	1
2371	3337	I. 241	Δ. 617	11 11 29.6	2.899	1	122 2 45.9	19.63	1	cB; vL; E 160° ±; am 4 st...	2
2372	853	II. 879		11 11 32.8	3.745	1	21 59 31.9	19.63	1	pB; S; R; gbM	2
2373	854		M. 65	11 11 36.7	3.139	3	76 8 46.9	19.63	3	B; vL; mE 165° ±; gbMBN.	8*+
2374	855			11 11 40.5	3.111	1	81 42 52.9	19.63	1	eF	1
2375		II. 885		11 12 17.2	3.512	1	31 22 36.2	19.64	1	F; S; lE; 135° ±	1
2376	856	II. 52		11 12 43.1	+3.164	1	70 52 55.5	+19.65	1	B; S; v lE; sbM	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and m.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s		° ' "	"			
2377	857 = 875?		M. 66	11 12 48.3	+3.137	2	76 15 17.5	+19.65	3	B; vL; mE150°; mbM; 2st up	9*†
2378	859	V. 8		11 12 57.4	3.140	4	75 38 28.5	19.65	4	pB; vL; vmE102°0	8†
2379	858	I. 226		11 13 1.7	3.435	1	36 4 6.5	19.65	1	pB; L; R; svmbMrN	2†
2380	860	II. 338		11 13 3.3	3.211	7	62 16 12.5	19.65	7	cF; L; R; vglbM	9
2381	861			11 13 5.2	3.089	2	86 16 11.5	19.65	2	pB; S; R; smbMN	3
2382		II. 30		11 13 6.1	3.163	1	71 4 37.5	19.65	1	pB; *inv	1*
2383	862	II. 550		11 13 21.8	3.027	2	99 30 44.8	19.66	2	F; vS; R; lbM; *7 f; p of 2	4
2384	863	II. 551		11 13 36.0	3.028	2	99 28 56.8	19.66	2	F; vS; R; psbM; *7p; f of 2	4
2385	856, a		R. nova	11 13 46.1	3.164	::	70 52 55.5	19.66	::	pF; S; R; vlbM; foll h. 856, 15'.	0
2386	864	II. 33		11 13 53.9	3.090	1	85 59 53.1	19.67	1	B; pL; R; psbM	3
2387	865	I. 245		11 14 10.0	3.515	4	30 9 43.1	19.67	4	pB; pL; R; vgbM	7
2388	867 = 861?	II. 32		11 14 20.6	3.088	1?	86 16 50.1	19.67	1?	pB; S; E; bM	1*
2389	866	III. 15		11 14 20.9	3.170	2	69 4 20.4	19.68	2	cF; cL; IE; gbM; sp of 2	3
2390	868			11 14 51.7	3.290	1	49 22 36.7	19.69	1	pB; S; pinE; bMN=close?	1
2391	869	III. 16		11 14 52.5	3.169	2	69 1 43.7	19.69	2	vF; pS; R; gbM; nf of 2	3
2392	870	III. 335		11 15 1.4	3.190	1	64 56 3.7	19.69	1	cF; vS; R; bM; np of 2	2
2393	872	III. 336		11 15 6.1	3.190	1	64 57 48.7	19.69	1	vF; vS; sf of 2	2
2394	871	II. 775		11 15 6.6	3.274	1	51 27 49.7	19.69	1	pF; cL; IE; vgbM	2
2395		II. 880		11 15 24.0	3.766	2	19 47 39.0	19.70	2	F; S; IE15°+	2
2396	873	I. 5		11 15 33.5	3.150	3	72 38 30.0	19.70	3	pB; pS; iR; bM; r	5
2397	874	II. 782		11 15 40.7	3.423	1	35 23 23.0	19.70	1	pB; S; R; vgbM; *12 p	2
2398	876	III. 768		11 15 58.6	3.409	1	36 18 44.3	19.71	1	cF; vS; R; stellar	3
2399	878	II. 53		11 16 22.1	3.154	1	71 25 28.3	19.71	1	cF; S; IE; r	3
2400	877	IV. 59		11 16 23.3	3.273	1:	50 42 2.3	19.71	1:	{ H. cB; S; R; svmbMN } h. F; R; *17 M.	3
2401		II. 635		11 16 27.0	3.037	1	97 52 41.3	19.71	1	F; pL; iR; vgbM	1
2402	3339	III. 530		11 16 37.0	3.014	1	103 3 31.3	19.71	1	F; S; R; stellar; p of 2	2
2403	879	IV. 4		11 16 39.9	3.070	1	90 19 53.3	19.71	1	vF; S; att to *13 m	3
2404	881	I. 219		11 17 10.2	3.271	2	50 28 40.6	19.72	2	cB; cL; iR; pgmbM	3
2405	882	I. 20		11 17 11.3	3.123	3	77 53 21.6	19.72	3	F; E90°+; B*f34°	5*
2406	3340	III. 531		11 17 11.8	3.015	2	103 4 29.6	19.72	1	pF; pL; iR; vlbM	4
2407	880	II. 845		11 17 14.0	3.569	1	25 47 2.9	19.73	1	F; pS; iR; gbM; *9 np	4
2408	883	II. 829		11 17 24.4	3.462	1	31 30 53.9	19.73	1	vF; pL; pmE135°±; er	3
2409	884	III. 337		11 17 25.2	3.182	1	65 16 34.9	19.73	1	vF; vS; R	2
2410	885	III. 922		11 17 54.2	3.503	1	28 45 10.2	19.74	1	vF; vS; 2 vS st inv	2
2411	886	I. 131		11 17 58.3	3.034	1	99 1 48.9	19.73	2	pB; L; E0°±; gbM	3*
2412	3341			11 18 20.4	2.956	1	115 58 29.2	19.74	1	F; vL; gvlbM; *7 s 6'	1
2413	887	I. 194		11 18 26.4	3.301	2	45 38 32.2	19.74	2	vB; cL; vmE0°±; vsmbMN; st p.	4
2414	888			11 18 34.5	3.328	2	42 14 27.2	19.74	2	cF; S; R; vsbM*; 2st 11 nf.	2
2415		II. 886		11 18 36.9	3.442	1	32 6 42.5	19.75	1	pF; iF	1
2416	889			11 18 47.7	3.199	1	61 21 44.5	19.75	1	vF; S; R; psbM; *12 nf	1
2417		III. 112		11 18 51.5	3.051	2?	95 4 52.5	19.75	2?	cF; cL; R; r (v near vB*)	3*
2418	3342		Δ. 481	11 18 56.6	2.858	2	132 27 50.5	19.75	2	Cl; cL; pRi; IC; st 10...14...	2
2419	891 = 3343	II. 159		11 19 10.3	3.145	4	72 22 2.5	19.75	4	B; pS; R; bM	5
2420	890	I. 262		11 19 19.3	3.618	1	22 38 33.8	19.76	1	cB; S; iR; spmbMN	2
2421	892	I. 246		11 19 32.8	3.431	1	32 20 57.8	19.76	1	cB; pL; E	3
2422	893			11 19 52.5	3.144	2	72 12 19.8	19.76	3	pB; pL; E; vgbM	3
2423	894	II. 160 = III. 28		11 20 23.8	3.144	2	72 0 27.1	19.77	2	pB; L; vIE; vgbM; r	5
2424	895	II. 770		11 20 32.3	3.202	5	59 42 52.1	19.77	5	pB; pS; R; lbM; r	6
2425	896	I. 247		11 20 44.8	+3.441	1	30 40 47.4	+19.78	2	pB; pS; vIE 80°±; pgbM; Sst sf nr.	4

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
2426	897	II. 339		11 20 45.7	+3.182	1	63 34 36.4	+19.78	1	pB; pL; lE; bM	2
2427	898	II. 54		11 20 49.6	3.142	2	72 18 40.4	19.78	2	F; pS; lE; r	4
2428		II. 152		11 20 59.6	3.111	1	79 41 44.4	19.78	1	F; mE; r	1
2429	3334	III. 532		11 21 9.0	3.023	1	102 24 55.4	19.78	1	cF; S; E; gbM	2
2430	899			11 21 22.8	3.231	1	53 48 40.7	19.79	1	cF; S; R; sbM*?	1
2431	900			11 21 30.0	3.158	1	68 25 51.7	19.79	1	eF; vS; E90°	1
2432	3345			11 21 33.5	2.706	1	149 11 3.7	19.79	1	B; pL; lR; pgpmbM	1
2433	901	II. 349		11 22 5.3	3.171	1	65 7 52.0	19.80	1	pF; pL; lE	2
2434	902	II. 13		11 22 53.7	3.109	1	79 57 4.3	19.81	1	pF; pL; R; vsmbM; r	4
2435	3346			11 22 53.9	2.920	3	125 37 34.3	19.81	3	pB; cS; R; psmbM	3
2436	903			11 23 0.3	3.109	1:	79 52 19.3	19.81	1	pB; cS; E90°	1
2437	904	II. 350		11 23 43.7	3.161	1	66 27 39.6	19.82	1	F; S; *7.8 nf 5'	2
2438	905			11 24 15.2	3.185	1	60 44 5.6	19.82	1	F; vS; R; smbM	2
2439	906	II. 367		11 24 17.1	3.183	3	61 4 23.6	19.82	3	F; cS; R; sbMN	4
2440	907	III. 353		11 24 28.7	3.183	3	60 52 0.9	19.83	3	F; S; R; psbM	5*
2441	3347	II. 562		11 24 29.0	3.024	2	103 27 38.9	19.83	2	pF; S; R; vglbM	5
2442	3348			11 24 36.6	2.958	2	119 29 4.9	19.83	2	pB; S; mE; *13 att	2
2443	908	I. 221		11 24 46.4	3.342	2	36 9 21.9	19.83	2	pB; vL; R; vglbM	4
2444	909	II. 836		11 25 43.2	3.442	2	27 21 32.2	19.84	2	cF; S; R; gvlbM; r	3
2445	910	II. 730		11 25 44.5	3.284	2	42 10 57.2	19.84	2	pB; vL; lE0°; vsmbM*15; *11 n.	3+
2446	912	II. 351		11 26 2.1	3.162	1	64 46 58.5	19.85	1	F; S; R; bM	2
2447	911	I. 222		11 26 2.3	3.333	1	36 5 47.5	19.85	1	pB; pL; lE0°±; gbM; *12 nr	3
2448		III. 80		11 27 5.9	3.115	1	76 43 50.8	19.86	1	vF; vS; R	1
2449	913	H. 552		11 27 8.0	3.042	2	99 3 46.8	19.86	2	F; S; R; psbM; *14 sp 225°	3
2450		III. 771		11 27 10.4	3.340	1	34 23 50.8	19.86	1	eF; S; lR; L * in field	1
2451	3349	III. 935		11 27 37.1	3.038	1	103 19 8.1	19.87	1	eF; S; R; gbM	2
2452	914	I. 287		11 27 44.4	3.602	1	18 41 33.1	19.87	1	pB; L; mE130°4; mbM ..	2
2453		III. 772		11 27 44.9	3.340	1	34 15 51.1	19.87	1	vF; stellar	1
2454		II. 783		11 28 6.8	3.331	1	34 41 52.1	19.87	1	pB; pL; bM	1
2455	915	III. 847		11 28 23.8	3.387	1	29 15 14.4	19.88	1	vF; vS; R; vgbM	2
2456	916			11 28 33.7	3.255	1	43 56 23.4	19.88	1	vF; S; R; vgbM	1
2457	3350			11 28 37.1	2.938	2	127 10 35.4	19.88	2	pF; pL; vLE; glbM	2
2458		III. 969		11 28 56.6	3.748	1	14 14 50.7	19.89	1	eF; S	1
2459	3351			11 28 59.4	2.939	2	127 12 45.4	19.88	2	F; cS; lE; gvlbM	2
2460	917	II. 905		11 29 3.2	3.784	1	13 56 57.4	19.88	1	pB; pL	2
2461	918	H. 784		11 29 4.5	3.319	1	34 55 43.7	19.89	1	pF; L; lE	2*
2462	920			11 29 4.7	3.203	1	52 49 3.7	19.89	1	eF; pL; pmE; gbM	1
2463	919	III. 843		11 29 8.5	3.360	1	30 49 18.7	19.89	1	vF; R; stellar; vS*1 d sf ...	1
2464			D'Arrest, 78	11 29 16	3.14	[1]	67 24 12	19.88	[1]	B; pS; mbMN=*13; *11 p 4 ^a , s 175 ⁿ .	0
2465	921	II. 837		11 29 29.6	3.398	1	27 29 1.7	19.89	1	F; vLE; gbM	2
2466			D'Arrest, 79	11 29 39	3.13	[2]	71 20 42	19.89	[2]	F; S; R	0
2467	922			11 29 41.9	3.150	1	65 7 26.7	19.89	1	vF; S; R	1
2468	3352		Δ. 289	11 29 42.0	2.764	2	150 49 29.7	19.89	2	Cl; pL; pRi; pC; st 8...13 ..	2
2469	923	III. 29		11 29 54.8	3.128	1	71 23 48.7	19.89	1	vF; eS; stellar	2
2470	924			11 29 55.8	3.125	1	72 20 37.7	19.89	1	vF; S; bM	1
2471	925	II. 731		11 30 10.0	3.262	2	41 19 6.0	19.90	2	pB; S; pmE	4
2472	926	II. 838		11 30 13.1	3.364	2	29 36 37.0	19.90	2	pF; S; R; gbM; r	3
2473	927	II. 352		11 30 27.9	3.143	2	66 32 34.0	19.90	2	vF; S; E; r	4
2474	928	III. 81		11 30 58.4	3.109	1	77 6 36.3	19.91	1	cF; cS; R; psbM	4
2475	3353			11 31 34.0	2.881	1	139 55 51.3	19.91	1	eF; S; R; am 50 Sst	1
2476	929	I. 227		11 31 41.7	3.315	2	32 57 27.3	19.91	2	pF; L; vLE; vgbM; r	4
2477	930	II. 732		11 32 1.9	3.241	1	42 45 34.6	19.92	1	F; S; att to *15; another * cont.	2
2478	3354			11 32 6.8	2.954	1	126 57 46.6	19.92	2	cB; R; sbMN*; *9 sf	2
2479	931			11 32 18.7	3.171	3	57 18 46.6	19.92	3	pB; pL; pmE; gbM; p of 2...	3
2480	932			11 32 21.7	3.171	3	57 17 20.6	19.92	3	cB; pL; pmE0°; pgbM	3
2481	933	III. 109		11 32 29.4	3.123	2	71 31 5.6	19.92	2	cF; vS; pmE; sbM; 2 Sst f; 1st of 3.	3
2482	935	III. 609		11 32 35.2	+3.048	1	98 35 14.6	+19.92	1	vF; vS; R; gbM; *8 s 6' ...	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
2483	h.	II. 773		h m s 11 32 45.1	+ 3.305	1	32 59 54.6	+ 19.92	1	cF; pS; vS*v nr	1
2484		III. 844		11 32 47.4	3.327	1	30 36 24.9	19.93	1	vF; S; mE	1
2485	937	II. 839		11 32 47.9	3.345	2	28 56 14.9	19.93	2	F; cS; R; mbM	3
2486	{ 934 = 3355 }			11 32 54.0	3.116	2	73 53 41.9	19.93	3	cF; R; p of 2	3†
2487	938	II. 340		11 32 54.5	3.144	1	64 31 35.9	19.93	1	F; cS; IE; stellar; r	3
2488	{ 936 = 3356 }	II. 103		11 32 57.8	3.116	4	73 52 36.9	19.93	6	F; pS; E; pglbM; r; f of 2...	8†
2489	936, a		R. nova	11 33 ±		...	73 52 ±		...	Smaller than h. 936	0
2490	939	II. 161		11 33 0.4	3.122	3	71 29 46.9	19.93	3	pF; pL; R; bM; r; 2nd of 3	4
2491	939, a	} R. 3 novæ		11 33 ±		...	71 29 ±		...	No description (for 939, c, see No. 5069).	0
2492	939, b										
5069											
2493	940	III. 30		11 33 5.0	3.122	3	71 27 36.9	19.93	2	vF; pS; r; 2 vB st p; 3rd of 3	4
2494		II. 830		11 33 23.7	3.300	1	32 58 24.9	19.93	1	pB; E	1
2495	941	III. 375		11 33 26.1	3.129	1	68 53 9.9	19.93	1	cB; cS; R; bM; r	3
2496		D'Arrest, 80		11 33 29	3.12	[1]	71 25 42	19.93	[1]	F; pL; *9.10, s 5'	0
2497				11 33 35.4	3.135	1	66 46 41.9	19.93	1	vF; vS	1
2498	942			11 33 45.6	3.237	2	41 30 46.9	19.93	2	F; S; vIE; glbM	4
2499	943	I. 21		11 33 45.7	3.104	3	77 45 10.9	19.93	4	B; L; vIE	8
2500	944	III. 320		11 33 48.3	3.142	1	64 24 10.9	19.93	1	cF; vS; R; p of 2; * 6 sf 3'	3
2501	945	I. 94		11 33 53.4	3.183	3	52 40 24.2	19.94	3	cB; pL; pmE 90°±; bM	5*
2502	946	III. 339		11 34 20.2	3.140	1	64 25 31.2	19.94	1	cF; S; f of 2	2
2503	947			11 34 40.4	3.099	1	78 55 43.2	19.94	1	F; 1st of 4	1
2504	948	III. 284		11 34 47.6	3.058	1	95 23 14.2	19.94	1	F; pS; R; psbM	3
2505	950			11 34 51.9	3.099	1	78 54 43.2	19.94	1	vF; 2nd of 4	1
2506	949	III. 376		11 34 52.5	3.126	2	68 54 26.5	19.95	2	vF; cS; R; bM; bet 2st.	5
2507	951	II. 153		11 34 58.4	3.099	1	78 57 13.5	19.95	1	pF; pS; 3rd of 4	2
2508	3357			11 35 9.4	3.039	1	103 4 35.5	19.95	1	F; cS; IE; psbM	1
2509	952	III. 774		11 35 10.2	3.258	1	36 26 51.5	19.95	1	vF; cS; pmE	3
2510	953	II. 154		11 35 11.4	3.099	1	78 58 3.5	19.95	1	pF; pS; 4th of 4	2
2511	954	II. 341		11 35 14.4	3.143	3	62 44 1.5	19.95	3	pF; S; R; psbM; stellar	4
2512		D'Arrest, 81		11 35 21	3.12	[1]	70 22 42	19.94	[1]	F; S; lbM	0
2513	955			11 35 51.3	3.253	1	36 29 51.5	19.95	1	vF; vS	2
2514	956			11 35 56.5	3.141	1	62 43 32.5	19.95	1	eF	1
2515	957			11 36 11.6	3.043	1	102 5 22.8	19.96	1	F; vS; R; bM	1
2516		III. 340		11 36 15.0	3.130	1	66 24 56.8	19.96	1	vF; pL; 2 suspected neb nr	1
2517		III. 102		11 36 15.9	3.097	1	79 6 56.8	19.96	1	eF; pS	1
2518		D'Arrest, 82		11 36 21	3.12	[1]	70 7 42	19.95	[1]	vF; vS; slbMN* 13 m	0
2519	958			11 36 24.4	3.306	2	29 6 47.8	19.96	2	pB; E; gbM; * 8 nf 5'	1
2520	959			11 36 29.2	3.286	2	31 16 21.8	19.96	2	pB; cS; E; psbM* 12	3
2521	960			11 36 32.6	3.121	2	69 15 52.8	19.96	2	cF; S; R; 1st of 5	1
2522	} 960, a		R. 4 novæ	11 36 ±		...	69 15 ±		...	8 "knots" (vide h. 960, 1, 2, 3)	0
2523											
2524											
2525	961			11 36 39.1	3.120	2	69 20 12.8	19.96	2	cF; S; R; 2nd of 5	1
2526	962	III. 377		11 36 44.2	3.120	3	69 17 13.8	19.96	3	F; S; R; vglbM; 3rd of 5	5
2528	963			11 36 46.7	3.120	1	69 14 9.8	19.96	1	vF; pS; 4th of 5	1
2529	964			11 36 53.3	3.264	1	33 34 40.8	19.96	1	F; pL; R; vglbM	1
2530	965			11 36 53.6	3.159	1	55 42 18.8	19.96	1	F; S; R; psbM	1
2531		III. 35		11 36 57.9	3.098	1	78 55 57.8	19.96	1	cF; vS	1
2532		III. 776		11 37 2.1	3.265	1	33 19 56.8	19.96	1	eF; pL; IE	1
2533	966	III. 378		11 37 6.3	3.120	1	69 15 33.8	19.96	1	eF; vS; R; 5th of 5	3
2534		III. 36		11 37 11.9	3.098	1	78 55 57.8	19.96	1	eF; vS	1
2535		III. 386		11 37 37.2	3.118	1	69 27 57.1	19.97	1	vF; vS; r	1
2536	970			11 37 45.7	3.118	1	69 14 20.1	19.97	1	F; S; R; bM(?)	1
2537		III. 385		11 37 53.2	3.117	1	69 35 57.1	19.97	1	vF; vS; r	1
2538	971			11 38 30.2	+ 3.096	2	78 23 56.4	+ 19.98	2	F; S; iR; psbM	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
2539	972	III. 833		11 38 34.3	+3.218	3	39 1 19.4	+19.98	3	cF; cS; R; psbM	4
2540	967			11 38 34.6	3.152	1?	56 6 59.4	19.98	1?	eF; R; gbM; 1st of 4	1*
2541	973	II. 104		11 38 35.6	3.102	3	75 27 32.4	19.98	3	B; S; R; smbM*	5
2542		III. 104		11 38 36.0	3.091	1	80 40 29.4	19.98	1	vF; vS; suspected	1
2543		III. 387		11 38 37.2	3.116	1	69 27 58.4	19.98	1	vF; vS; r	1
2544		III. 103		11 38 37.6	3.093	1	80 1 58.4	19.98	1	vF; r	1
2545		I. 201		11 38 55.9	3.202	2	41 43 58.4	19.98	2	B; L; mE 25° ±	2
2546	974			11 38 56.4	3.150	1	56 1 20.4	19.98	1	vF; R; 2nd of 4	1
2547		II. 881		11 39 0.8	3.391	1	19 49 57.4	19.98	1	F; pL; mE 105° ±	1
2548	{ 968 = 975 969			11 39 1.4	3.150	2	56 3 20.4	19.98	2	vF; R; gbM; 3rd of 4	2
2549	{ 976 = 3358			11 39 14.9	3.149	2	56 7 5.4	19.98	2	vF; R; gbM; 4th of 4	2
2550				11 39 16.6	2.903	2	145 36 20.4	19.98	2	vF; lE; 2 st inv	1
2551		III. 372		11 39 21.8	3.112	1	68 35 58.4	19.98	1	vF; cL	1
2552	977	III. 388		11 39 30.9	3.116	1	68 49 39.4	19.98	1	cF; S; iR; gbM; r; *7 sp 6'	4
2553	3359	III. 828		11 39 45.1	3.014	2	117 8 33.7	19.99	2	cF; vS; vLE; bM; vF*sf ...	2
2554	{ 979 = 3360	I. 120		11 39 56.8	3.040	2	106 4 52.7	19.99	2	pB; L; iR; vgpmbM	3
2555	978	II. 785		11 40 6.3	3.238	2	33 15 24.7	19.99	2	pB; S; lE; pgbM	3
2556	980	II. 723		11 40 45.5	3.136	3	58 51 30.7	19.99	3	pB; S; bM	4
2557	{ 981 = 3361	II. 553		11 40 53.3	3.053	2	100 10 27.7	19.99	2	pB; pL; R; gbM; r	4
2558		III. 940		11 40 54.2	3.462	1	14 52 59.0	20.00	1	vF; S; R; bM	1
2559	982	II. 738		11 41 12.9	3.192	1	40 30 26.0	20.00	1	B; pL; R; mbM	2+
2560	983	I. 248		11 41 18.7	3.250	2	29 48 43.0	20.00	2	B; pL; iR; pgmbM; p of 2 ..	4
2561	984	II. 832		11 41 29.7	3.248	1	29 47 51.0	20.00	1	pF; pL; vLE; gbM; f of 2 ...	3
2562		II. 739		11 41 36.1	3.189	1	40 30 59.0	20.00	1	F; vS	1
2563	986	II. 408		11 41 41.5	3.144	3	54 11 35.0	20.00	3	F; S; R; bM	5
2564	985	I. 228		11 41 45.6	3.224	4	33 8 26.0	20.00	4	B; pL; lE; svmbM	6
2565	987			11 41 51.5	3.123	1	62 46 31.0	20.00	1	pB; R; smbM	1
2566	988	I. 82		11 41 53.3	3.124	5	62 11 43.0	20.00	5	B; pL; vLE 0° ±; bMN	8
2567		III. 970		11 41 58.8	3.564	1	11 7 57.0	20.00	1	pF; pL; r	1
2568	989	III. 321		11 42 1.2	3.122	2	63 5 50.0	20.00	2	F; pS; lE; vglbM	3
2569	3362			11 42 1.4	2.998	2	126 44 21.0	20.00	2	pB; cS; vLE; lbM	2
2570	3363	II. 864		11 42 9.2	3.019	1	118 32 27.0	20.00	1	pB; S; R; mbM	2
2571		III. 715		11 42 11.9	3.185	1	40 48 0.0	20.00	1	eF; pL	1
2572	990			11 42 19.8	3.065	1	90 19 7.0	20.00	1	eF; S; psbM	1
2573	3364			11 42 34.3	2.967	1	137 29 23.0	20.00	1	Cl; vL; lC; st 9...14	2
2574	991	III. 341		11 42 42.2	3.117	(1)	64 17 48.0	20.00	1:	vF; S; p of 2	2
2575	992	II. 342		11 42 49.0	3.120	3	62 44 0.3	20.01	3	F; pL; R; pgbM	4
2576		II. 786		11 43 0.7	3.209	1	33 53 0.3	20.01	1	F; E	1
2577		III. 113		11 43 4.8	3.066	1	94 21 0.3	20.01	1	eF; eS; bet 2 st	1*
2578	993	II. 787		11 43 20.3	3.206	1	34 4 32.3	20.01	1	eF; R; gbM	2
2579	994			11 43 20.4	3.190	1	37 23 43.3	20.01	1	F; L; vmE; vgbM	1
2580	995	III. 90		11 43 20.5	3.084	2	82 38 55.3	20.01	3	F; vS; R; lbM; *13 np 80'' ..	4
2581	3365			11 43 23.9	2.934	4	146 24 8.3	20.01	5	O; !; S; R; blue; =*7m; 1*5=d.	6
2582	996			11 43 33.5	3.115	1:	64 17 48.3	20.01	1	Neb; f of 2	1
2583		II. 824		11 43 36.9	3.193	1	36 21 30.3	20.01	1	pB; L; mE	1
2584	997	II. 788		11 43 37.1	3.203	1	34 8 32.3	20.01	1	pF; S; R; pspmbM	2
2585		III. 716		11 43 47.7	3.180	1	39 1 0.3	20.01	1	vF; vS	1
2586	3366	I. 259		11 43 57.8	3.025	2	118 3 5.3	20.01	2	B; pL; lE; gmbM; r; vS*sp inv.	3
2587		II. 825		11 43 59.4	3.189	1	39 8 0.3	20.01	1	pB; S; iF; bM	1
2588			D'Arrest, 83	11 44 6	3.11	[1]	67 19 48	20.01	[1]	vF; vS	0
2589	998	III. 379		11 44 11.2	+3.108	1	67 12 20.3	+20.01	1	eF; eS; vLE; er; st nr.	4

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s						
2590	999	II. 740		11 44 24.7	+3.170	1	40 31 56.6	+20.02	1.	pF; S; R; pspmbM	2
2591	1000	III. 616		11 44 29.7	3.139	2	51 13 20.6	20.02	2.	eF; cL; iF; glbM; *6n5'; *7f	4*
2592		III. 769		11 44 39.8	3.182	1	37 16 0.6	20.02	1.	eF; S	1
2593			D'Arrest, 84	11 44 49	3.17	[1]	40 35 48	20.01	[1]	vF; v diffie; H. II. 740 np ...	0
2594	1001			11 45 8.9	3.124	3	56 48 50.6	20.02	3.	pF; S; IE; psbM	3
2595	3367			11 45 19.8	3.033	1	116 7 21.6	20.02	1.	vF; cL; vmE 59°3	1
2596	1003	III. 389		11 45 28.9	3.103	1	68 35 6.6	20.02	1.	vF; cS; R	2
2597	1002	I. 203		11 45 30.5	3.150	2	45 5 35.6	20.02	2.	B; vL; R; bMpBN; er	3†*
2598		III. 971		11 45 31.1	3.378	1	14 7 0.6	20.02	1.	eF; vS; R	1
2599	1004	III. 380		11 45 34.2	3.103	1	68 14 0.6	20.02	1.	vF; cS; R	3
2600	1005	I. 173		11 45 38.3	3.132	3	52 13 50.6	20.02	3.	vB; pL; R; smbM*9	4
2601	1007	III. 322		11 45 50.3	3.111	3	63 0 39.6	20.02	3.	pF; pS; R; psbM	4
2602	1006	I. 251		11 45 50.5	3.211	2	28 33 14.6	20.02	2.	B; pL; R; gmbM; r; *f	3
2603	1008	II. 403		11 46 7.9	3.101	1	68 28 0.6	20.02	2.	F; pS; iE; lbM; *p	5
2604	1009	I. 202		11 46 21.7	3.155	2	41 21 59.9	20.03	2.	cB; pL; pmE; vgbM	4*
2605	1010	III. 342		11 46 22.4	3.105	1	65 49 28.6	20.02	1.	vF; cS; vIE	2
2606	1011	V. 45		11 46 27.4	3.169	1	36 53 3.9	20.03	1.	cB; L; E 0°+; vsbMLrN ...	3†
2607	1012	III. 612		11 46 28.1	3.068	2	93 13 23.9	20.03	2.	cF; cS; IE 90°±; bM; r ...	4
2608	1013	III. 381		11 46 30.0	3.101	1	68 20 44.9	20.03	1.	eF; R	2*
2609		II. 623		11 46 53.7	3.043	2	112 24 0.9	20.03	2.	cF; S; E 170°±; lbs	2
2610	3368	III. 290		11 46 56.8	3.047	1	109 47 29.9	20.03	1.	eF; pL; pmE 56°8	2
2611		II. 294		11 46 58.4	3.048	1	108 47 0.9	20.03	1.	F; S; E; r	1
2612	1014	II. 833		11 47 1.8	3.188	2	30 51 20.9	20.03	2.	pF; pS; pmE; vgbM	3
2613	1015	IV. 67		11 47 27.9	3.184	2	30 43 51.9	20.03	2.	pF; cL; R; vg, sbM	3
2614	3369		Δ. 349	11 47 29.3	2.976	1::	144 56 21.9	20.03	1.	Cl; pL; pRi; gplmbM; st 13	3
2615		III. 905		11 47 31.4	3.254	1	19 54 0.9	20.03	1.	eF; vS	1
2616	3370	I. 67		11 47 32.4	3.056	2	103 11 30.9	20.03	2.	cB; pL; iR; gmbM; Δ 2st...	6
2617	1016			11 47 39.1	3.108	2	60 57 8.9	20.03	2.	vF; S; E; *10nf att	3
2618		II. 789		11 48 4.2	3.166	1	33 54 0.9	20.03	1.	pB; E	1
2619		II. 790		11 48 4.2	3.166	1	33 54 0.9	20.03	1.	F; S	1
2620	1017	IV. 62		11 48 6.5	3.165	1	34 6 0.9	20.03	1.	B; pL; R; g, sbM disc	2
2621	1018	II. 162		11 48 16.5	3.086	2	77 15 16.9	20.03	2.	pB; L; iR; bM; *10, 25°, 5'	4
2622	1018, a		R. nova	11 48 +		...	77 15 -		...	nf h. 1018	0
2623	1020			11 48 18.9	3.060	1	101 15 54.9	20.03	1.	F; S; R; psbM; p of 2	1
2624	1019	II. 724		11 48 20.0	3.109	1	59 13 39.9	20.03	1.	pF; vS; R; bM	2
2625	1021			11 48 32.4	3.060	1	101 12 44.9	20.03	1.	vF; S; R; bM; f of 2	1
2626	1022	II. 132		11 48 45.9	3.079	3	82 28 4.2	20.04	3.	B; pL; cE 30°; vsmbMN ...	4
2627	1023	II. 840		11 48 54.9	3.178	2	28 42 30.2	20.04	2.	cF; S; IE; bM; *8, 90°, 6'...	3
2628		III. 274		11 49 3.7	3.052	1	109 7 1.2	20.04	1.	vF; pL; iF	1
2629	1024	III. 343		11 49 10.1	3.098	1	65 20 55.2	20.04	1.	cF; cS; R; psbM	2
2630	1026			11 49 23.3	3.104	1	60 13 26.2	20.04	1.	eF; S; R; bM	1
2631	1025	III. 707		11 49 23.7	3.136	1	40 52 57.2	20.04	1.	vF; cS; another suspected ...	3
2632	1027			11 49 30.3	3.107	3	57 11 59.2	20.04	3.	pF; S; pmE 90°±; *11 nr...	1
2633	1028			11 50 9.6	3.100	1	61 20 43.2	20.04	1.	vF; S; R; bM*; p of 2	2
2634	1029	II. 791		11 50 14.3	3.147	1	33 46 31.2	20.04	1.	pF; S; IE; psbM	2
2635	1030	IV. 61		11 50 15.6	3.142	1	35 50 34.2	20.04	1.	cB; vL; pmE; sbMBrN	3
2636	1032			11 50 34.2	3.085	3	74 55 29.2	20.04	2.	vF; pL; R; 2st f	3
2637	1031	I. 229		11 50 34.3	3.145	1	33 46 21.2	20.04	1.	cB; pS; R; vg, smbM	2
2638	1033, a		R. 3 novæ	11 50 ±		...	63 57 ±		...	one S, R; the other two E ...	0
2639						...					
2640						...					
2641	1033	III. 323		11 50 36.1	3.095	1	63 57 34.2	20.04	1.	pF; vS; E 25° bet 2 st	2
2642	1040, a		R. nova	11 50 44.7	3.121	::	41 54 13.2	20.04	::	S; R; 7' np h. 1040	0
2643	1034	III. 344		11 50 45.6	3.093	1	66 0 39.2	20.04	1.	vF; vS; R; n of 2	2
2644	1035	III. 345		11 50 45.6	3.093	1	66 4 59.2	20.04	1.	vF; vS; R; s of 2	2
2645	1036	III. 354?		11 50 52.9	3.097	2	61 20 22.2	20.04	2.	F; S; R; *12 near	3
2646		III. 324		11 50 53.4	3.095	1	63 53 51.2	20.04	1.	eF; suspected	1
2647	1037			11 50 56.3	3.070	1	91 21 8.2	20.04	1.	F; S; R; bM; *11 nf	1
2648		III. 325		11 50 57.6	3.087	1	66 6 2.2	20.04	1.	eF; vS	1
2649	1038	II. 368		11 51 3.0	3.097	4	61 1 33.2	20.04	3.	pB; pS; R; psbM; r	5
2650	1039			11 51 4.6	+3.097	1	61 10 ±?	+20.04	0	vB; mE; mbM (?P.D.)	1*

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
2651	1040			11 51 14.4	+3.121	1	41° 59' 10.2	+20.04	1	F; pL; mE; vglbM	1
2652	1041	II. 733		11 51 19.2	3.116	2	45 17 9.2	20.04	2	B; cL; mE 62° 3; vsvmbM * 10	3*†
2653	1042			11 51 24.9	3.086	2	73 2 32.2	20.04	2	pB; pS; R; psbM	2*
2654	1043, u		R. nova	11 51 33	3.095	:	61 39 50	20.04	:	vF	0
2655	1043	II. 369		11 51 33.0	3.095	(1)	61 44 50.2	20.04	2	F; L; E; gbfM	4
2656	1044			11 51 42.8	3.083	1	75 0 46.2	20.04	1	eF; * 9 sf 5'	1
2657	1045	II. 725		11 51 43.9	3.097	1	58 47 57.5	20.05	1	pB; pL; E 19° 5; biN	2
2658		II. 295		11 51 56.2	3.059	1	107 35 2.5	20.05	1	F; vS; iF; bM	1
2659	1046	III. 617		11 51 57.7	3.104	1	51 24 39.5	20.05	1	eF; pL; R	3
2660	1047	I. 223		11 52 6.8	3.122	1	38 15 44.5	20.05	1	vB; cL; mE 160° ±; vsvmbMBN.	3
2661	3371	II. 296		11 52 23.1	3.059	1	108 29 25.5	20.05	1	⊕; pF; pL; R; rr; st 16 ...	2
2662		III. 3		11 52 45.0	3.083	2	73 0 1.5	20.05	2	vF; vS; vLE; r	2
2663	1048	I. 121		11 53 12.3	3.072	2	90 19 7.5	20.05	2	cB; L; vLE; psmbM; B st nr	4
2664	1049	II. 404		11 53 21.0	3.084	6	69 8 34.5	20.05	6	pF; pL; R; gbM; * 12 nf...	7
2665		II. 508		11 53 21.4	3.062	1	107 3 2.5	20.05	1	pB; S; iE; bM	1
2666		III. 903		11 53 45.4	3.155	1	19 52 1.5	20.05	1	eF; S; iF; gvlbM	1
2667	3372	III. 279		11 53 50.5	3.064	1	105 10 13.5	20.05	1	eF; pL; * 945° ±	2
2668	1050	I. 253		11 54 12.4	3.125	1	27 19 59.5	20.05	1	{ H. vb; vL; E } { h. pB; 25"; R }	2*
2669	1051	III. 77		11 54 13.1	3.079	2	75 49 13.5	20.05	2	eF; pL; R; r	3
2670	1052	IV. 28.1		11 54 43.7	3.064	1	108 5 11.5	20.05	1	pB; cL; R; vgbM	2†
2671	1053	IV. 28.2		11 54 43.7	3.064	1	108 7 11.5	20.05	1	pF; pL	2†
2672	1054	I. 252		11 54 58.6	3.117	2	27 5 10.5	20.05	2	B; cL; R; g; psvmbMrN ...	3
2673	1055			11 55 10.9	3.074	1	84 52 38.5	20.05	2	pF; S; R; psbM; * f 30° ...	2
2674	1056	III. 491		11 55 18.9	3.072	1	89 25 53.5	20.05	1	cF; cS; R; bM	3
2675	1057	II. 276		11 55 32.9	3.072	1	87 14 48.5	20.05	1	pF; L; R; sbM; * sf	4
2676	1058	II. 741		11 55 36.5	3.095	1	40 35 7.5	20.05	1	pB; pS; R	2
2677	1059			11 55 38.9	3.079	2	71 12 12.5	20.05	2	vF; vS; R; psbM	2
2678	1060	III. 390		11 55 41.8	3.079	1	70 28 21.5	20.05	3	eF; pS; R; glbM	4
2679		II. 509		11 55 47.7	3.066	1	105 36 2.5	20.05	1	F; cL; iR; lbM	1
2680	1061	IV. 56		11 55 58.3	3.089	4	44 41 3.5	20.05	4	B; vL; E; vg; vsmbM * 11...	5†
2681	3373			11 55 59.6	3.039	1	152 24 13.5	20.05	1	Cl; pRi; iC	1
2682		III. 794		11 56 8.6	3.099	1	31 18 2.5	20.05	1	eF; S	1
2683	1062			11 56 32.1	3.078	1	68 10 1.5	20.05	1†	pB. P.D. very doubtful ...	1*
2684	1063			11 56 36.6	3.078	1::	68 8 1.5	20.05	1†	pB. P.D. very doubtful ...	1*
2685	1064			11 56 40.6	3.078	1::	67 55 1.5	20.05	1†	pB. P.D. very doubtful ...	1*
2686	1065			11 56 51.4	3.077	3	68 59 39.5	20.05	3	vF; S; R; D neb pos 70° ...	3*
2687	1066	I. 174		11 56 53.2	3.080	4	57 19 32.5	20.05	4	pB; vL; mE 97°; vgbM	5
2688			D'Arrest, 85	11 56 57	3.07	[1]	70 46 42	20.06	[1]	B; E; gbm * 17 p, 82" dist.	0
2689	1067			11 56 57.5	3.077	2	68 59 58.5	20.05	2	pF; R	2*
2690	1068			11 56 59.7	3.076	4	68 51 54.5	20.05	4	pB	4*
2691	1069	III. 37		11 57 1.0	3.074	5	78 21 53.5	20.05	5	F; pS; R; gbM	6
2692		II. 781		11 57 1.6	3.087	2	36 40 2.5	20.05	2	pF; S; stellar	2
2693	1070	III. 392		11 57 1.7	3.076	1	68 53 38.5	20.05	1	vF; vS	2*
2694	1071	III. 391		11 57 2.0	3.076	2	68 49 13.5	20.05	2	F; vS	2*
2695	3374			11 57 4.9	3.046	1	156 31 32.5	20.05	1	vF; vS; R; bM*; am st ...	1
2696	1072	II. 277		11 57 17.2	3.072	4	87 19 26.5	20.05	4	F; pS; R; pgbM; np of 2 ...	7
2697	1073	III. 393		11 57 20.8	3.076	1	68 54 1.5	20.05	1	eF; vS	2*
2698	1074			11 57 20.9	3.072	1	87 9 8.5	20.05	1	F; S; R	1
2699	1075	III. 394		11 57 24.7	3.076	1	69 3 31.8	20.06	1	vF; vS	2*
2700	1076	III. 258		11 57 29.5	3.072	1	87 26 4.8	20.06	1	cF; cS; vLE; bM; sf of 2 ...	3
2701		III. 395		11 57 32.3	3.075	2†	69 7 1.8	20.06	2†	vF; vS	1*
2702		III. 396		11 57 32.3	3.075	2†	69 7 1.8	20.06	2†	vF; vS	1*
2703	1077			11 57 38.1	3.072	1	91 36 8.8	20.06	1	F; L; R; * 10 n 60"	1
2704	1078	III. 355		11 57 42.0	3.076	4	62 13 27.8	20.06	4	cF; pS; E; gbM	5
2705			D'Arrest, 86	11 58 13	3.08	[1]	38 54 7	20.06	[1]	F; iE; I. 206 nr	0
2706	3375	III. 754		11 58 24.4	3.070	1	115 44 43.8	20.06	1	pB; S; R; bM	2
2707		I. 224		11 58 40.3	3.074	2	38 56 2.8	20.06	2	B; pL; pmE; vsbM	2
2708		I. 206		11 58 42.2	3.074	3	38 42 22.8	20.06	3	B; cL; pmE 135° ±; lbM ...	3
2709	3376			11 58 44.2	+3.072	1	103 45 28.8	+20.06	1	eF; L; pmE; vgbM; 2 st 11 nr	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. II.'s Catalogues of Nebulæ.	Sir W. II.'s Classes and Nos.	Other Authorities.								
2710	h. 1079	II. III. 382		11 58 45.3	+3.072	1	68 38 1.8	+20.06	2	vF; vS.....	4
2711	1081	I. 207		11 58 51.9	3.072	1	41 44 34.8	20.06	1	pB; vL; mE 32° 0	4
2712	1080	III. 400		11 58 53.1	3.072	2	52 21 4.8	20.06	2	eF; vS; R; stellar; *10 sp 2'	6
2713	1082	III. 383		11 58 54.0	3.072	2	68 36 5.8	20.06	3	eF; eS; R; bM	5
2714		III. 384		11 58 57.0	3.072	1	68 35 2.8	20.06	1	eF; eS	2
2715	1084	III. 717		11 58 59.8	3.072	1	39 38 47.8	20.06	1	pB; vL; vmE 166° 5; vgvbM	3
2716	1083	III. 326		11 59 0.1	3.072	2	63 39 44.8	20.06	2	eF; vS; R; vgbM	3
2717	1085	I. 225		11 59 15.4	3.070	1	36 30 34.8	20.06	1	B; pS; R; bMBN; *12 sp, v, nr	3
2718	3377		Δ. 291	11 59 28.0	3.077	3	150 27 56.8	20.06	3	Cl; pL; pC; iR; st 10...14...	3
2719	1086	II. 370		11 59 28.8	3.070	6	61 2 37.8	20.06	6	pB; pS; IE; bM	7
2720	3378	II. 865		11 59 29.4	3.074	2	119 0 26.8	20.06	2	pF; pS; R; psbM; r; p of 2	3
2721	3379	II. 866		11 59 34.4	3.074	2	119 0 41.8	20.06	2	pF; pS; R; pgbM; f of 2	3
2722	1087			11 59 39.8	3.062	1	22 3 42.8	20.06	1	B; S; R; gbM	1
2723	1088	I. 195		11 59 52.9	3.067	2	46 9 26.8	20.06	2	vB; pS; mE 151° 0	4
2724	3380			11 59 55.8	3.077	2	129 25 18.8	20.06	2	F; S; vLE; glbM; 3Bst nr	2
2725	1089			12 0 0.5	3.068	1	55 13 46.8	20.06	1	eF	1
2726	3381	III. 533		12 0 0.9	3.074	1	102 24 23.8	20.06	1	eF; S; iR; gbM	2
2727	1090			12 0 1.7	3.070	1	74 49 13.8	20.06	1	eF; suspected	1
2728	1092, a		R. nova	12 0 19.8	3.072	...	86 30 33.5	20.06	...	Hook-shaped; h. 1092 is nf 45°; 14' dist.	0
2729	1091	III. 708		12 0 26.9	3.064	1::	46 12 1.8	20.06	1::	vF; vS.....	2
2730		II. 14		12 0 42.1	3.070	1	79 41 2.5	20.05	1	IE	1*
2731		III. 904		12 0 56.3	3.040	1	19 38 2.5	20.05	1	eF; vS; E	1
2732	1093			12 0 59.2	3.064	1	56 13 7.5	20.05	1	eF; vS; R; mbM	1
2733	1092	V. 4		12 0 59.4	3.072	2	86 20 39.5	20.05	2	eF; vL; E 90° ±; bM *16...	4+
2734	1094	{ I. 33 = II. 60 }		12 1 1.2	3.070	2	78 50 36.5	20.05	3	pB; pL; mE 120°; bM; r	5
2735			Auw. N. 28	12 1 3.1	3.045	...	24 2 50.6	20.05	...	pB; pL; cE; mbMN (Hind, Jan. 5, 1850).	0
2736	1095	III. 68		12 1 28.2	3.067	1	73 5 11.5	20.05	1	vF; S; R; psbM; bet 2 vS st	2
2737	1096	I. 279		12 1 29.2	3.005	2	12 25 10.5	20.05	2	F; pL; vLE; glbM	4
2738		I. 263		12 1 33.2	3.032	1	20 25 2.5	20.05	1	cB; IE; bM	1
2739	{ 1097 = 3382 }	II. 548		12 1 42.3	3.074	3	98 15 19.5	20.05	3	F; pL; pmE 95° ±; vglbM	4
2740	1098	III. 356		12 1 47.9	3.063	2	59 56 7.5	20.05	2	eF; S; R; 1st of 3	3
2741	1099	III. 357		12 1 52.9	3.063	3	59 58 57.5	20.05	3	eF; S; iR; 2nd of 3	4
2742	1100	I. 278		12 1 59.4	3.003	2	14 19 20.5	20.05	2	pB; cL; R; gmbM	3
2743	1101	II. 371		12 2 1.4	3.062	2	60 3 7.5	20.05	2	pF; pL; IE; 3rd of 3	4
2744	1108	II. 321		12 2 6.7	3.061	1	59 18 5.5	20.05	1	F; vL; vgbM	3
2745		I. 196		12 2 23.2	3.053	2	45 32 32.5	20.05	2	B; pL; IE; vgbM; *np	2
2746	1102	III. 795		12 2 29.8	3.037	1	30 22 0.5	20.05	1	vF; pS; IE; gbM; r	3
2747	1103	III. 814		12 2 31.4	3.044	1	36 6 6.5	20.05	1	vF; S; iF; vglbM; er	2*
2748	1104	IV. 54		12 2 43.5	3.051	1::	46 44 3.5	20.05	1	cB; R; vg, vsbMN	2
2749	1107	II. 747		12 2 54.1	3.047	2	42 46 19.5	20.05	2	pF; cL; vmE 109° 0; vgbM	3
2750	1105	I. 169		12 2 54.5	3.053	1	49 20 37.5	20.05	1	B; vL; vglbM	2
2751	1106, a		R. nova	12 2 +		...	70 40 +		...	S; prec h. 1106	0
2752	1106	I. 19		12 2 55.5	3.064	3	70 40 17.5	20.05	3	⊕; vB; pL; R; gbM; rrr	4
2753		III. 327		12 2 56.8	3.060	1	62 48 2.5	20.05	1	vF; pS	1
2754	1109	II. 802		12 3 22.6	3.030	1	30 56 13.5	20.05	1	F; S; E	2
2755	1110	I. 73		12 3 26.7	3.057	1	58 49 5.5	20.05	1	B; S; R; pgmbM	3
2756	1111	I. 165		12 3 27.0	3.050	2	49 49 31.5	20.05	2	vB; S; R; vsmbMBN; p of 2	4+
2757	1112	II. 83		12 3 28.7	3.064	3	73 11 28.5	20.05	3	pB; pL; R; pgmbM; r	4
2758		I. 11		12 3 36.9	3.063	1	70 52 2.5	20.05	1	B; pL; E; bM	1
2759		III. 845		12 3 42.6	3.026	1	30 53 2.5	20.05	1	vF; S; E 90° +	1
2760	1113	II. 642		12 3 46.5	3.049	2:	49 45 45.5	20.05	2:	pF; S; E; vgbM; f of 2	4+
2761	1114	I. 208		12 4 0.9	3.036	1	38 43 54.5	20.05	1	pF; cL; vmE 60° +	4
2762	1115	II. 405		12 4 4.2	3.061	1	69 2 58.5	20.05	1	F; pS; IE; bM; pB * nf	3
2763	1116	III. 941		12 4 4.7	2.945	1	13 5 48.5	20.05	1	eF; pS; R; Δ 2 st	2
2764		II. 803		12 4 39.4	+3.018	2	31 27 2.5	+20.05	2	F; S; R	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. II.'s Catalogues of Nebulae.	Sir W. II.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s		° ' "	+ "			
2765	1117	II. 353		12 4 46.9	+3.057	1	65 5 49.5	+20.05	1	B; L; iE; bM.....	2
2766		III. 399		12 4 56.4	3.046	2	53 2 2.5	20.05	2	vF; pL; vE; er.....	2
2767	1118			12 5 4.5	3.045	1	52 43 3.5	20.05	1	F; pL; R; vgbM; * sp 10'...	1
2768	1119	II. 105		12 5 8.6	3.063	1	76 1 5.5	20.05	1	pB; pL; iF; psbM; r; * inv	3
2769	1120	III. 358		12 5 14.5	3.051	2	60 3 17.5	20.05	2	F; S; 1st of 4.....	4
2770	1123	II. 792		12 5 16.6	3.016	1	33 2 58.5	20.05	1	F; S; iE; gbM.....	2
2771	1121	II. 372		12 5 17.2	3.051	2	60 0 1.5	20.05	1.:	F; S; 2nd of 4.....	4*
2772	1122	III. 359		12 5 18.7	3.051	2	60 5 31.5	20.05	2.	F; S; 3rd of 4.....	4
2773	1124	III. 360		12 5 27.2	3.050	2	60 3 31.5	20.05	2	F; cS; 4th of 4.....	4*
2774	3383	III. 534		12 5 30.9	3.081	1	103 14 38.5	20.05	1.	vF; pL; R; vgbM.....	2
2775	1125			12 5 35.2	2.993	1	78 22 7.5	20.05	2	vF; vL; E 45° ±; *7 f.....	2
2776	1126	I. 9		12 5 42.9	3.070	1	87 55 28.5	20.05	1	pB; pS; pmE 135° ±; bMN	5
2777	1127	II. 133		12 5 53.0	3.067	3	82 11 3.5	20.05	3	pF; S; iE 0° ±; r.....	5
2778		III. 777		12 6 4.8	3.016	1	36 20 2.5	20.05	1	eF; S; stellar.....	1
2779	1128	III. 697		12 6 13.1	3.031	2	45 32 36.5	20.05	2	vF; cL; mE 170° ±.....	5
2780	3384			12 6 14.4	3.152	1	151 56 12.2	20.04	1	Cl; mC; st eS.....	1
2781	1129	II. 373		12 6 15.1	3.048	3	60 42 57.2	20.04	3	eF; L; R; gbM.....	5
2782		II. 813		12 6 28.8	3.018	1	38 30 2.2	20.04	1	pB; S; iE.....	1
2783	1131	II. 106		12 6 38.7	3.061	1	75 48 11.2	20.04	1	F; L; iE; vglbM; r.....	3
2784	1133	II. 409		12 6 39.4	3.038	1	52 35 28.2	20.04	1	eF; pS; R; vglbM; r.....	3
2785	1130			12 6 39.6	3.066	4	82 1 3.2	20.04	4	cF; R; bM; near S*.....	4
2786	1132		M. 98	12 6 40.0	3.060	3	74 19 1.2	20.04	4	B; vL; vmE 152° 1; vsvmbM	7
2787	1134	II. 163		12 6 44.6	3.061	1	76 3 5.2	20.04	1	vF; pL; E; vgbM.....	2
2788	1135	II. 867		12 7 6.6	3.004	1	34 40 46.2	20.04	1	pB; vS; vsbM *12.....	2
2789		III. 796		12 7 8.8	2.988	1	29 34 2.2	20.04	1	eF.....	1
2790	1136	II. 374		12 7 26.4	3.045	4	60 48 15.2	20.04	4	pB; S; R; vsmbM *.....	5
2791	1137	II. 134		12 7 28.5	3.066	1	83 24 54.2	20.04	1	pF; pmE; vgbM.....	2
2792	1139	II. 793		12 7 29.7	2.997	1	33 12 18.2	20.04	1	pF; pS; iE; gbM.....	3
2793		III. 797		12 7 32.0	2.983	2	29 16 2.2	20.04	2	vF; S.....	2
2794	1138	II. 164		12 7 32.7	3.061	2	77 3 13.2	20.04	2	cF; { H. vmE h. R, 2 obs. } lbM ...	3
2795		II. 165		12 7 44.7	3.060	1	76 4 2.2	20.04	1.	F; vmE.....	1
2796	1140	I. 175		12 8 1.5	3.037	2	56 1 26.2	20.04	2	vB; S; R; psmbM.....	3
2797	1141	III. 397		12 8 8.0	3.051	1	68 33 36.2	20.04	2	vF; cL; iR; vgbM.....	4
2798			D'Arrest, 87	12 8 9	2.96	[1]	25 25 42	20.04	[1]	pB; pS; R; *12 f; ln.....	0
2799	1142	II. 107		12 8 19.4	3.058	1	75 19 25.2	20.04	1	vF; pL; R; gbM.....	2
2800		II. 375		12 8 21.7	3.041	1	60 43 22.2	20.04	1	F; pS.....	1
2801	1143	III. 850		12 8 23.4	2.945	1	23 14 29.2	20.04	1	pF; pS; R; vgbM.....	2
2802	1144	II. 108		12 8 31.3	3.057	2	75 19 1.2	20.04	2	B; L; E 90° ±; g, sbM; r...	3
2803	1145	II. 354		12 8 31.6	3.047	1	65 13 59.2	20.04	1	cF; vS; R.....	2
2804	1146	I. 95		12 8 36.0	3.030	1	52 54 3.2	20.04	1	cB; cL; iE; biN.....	3+
2805	1147	II. 135		12 8 44.8	3.065	3	82 49 9.2	20.04	3	B; pS; E; sbM *11.....	4
2806	1148	I. 35		12 8 46.7	3.058	2	76 4 2.2	20.04	3	vB; vL; vmE 17° ±; sbMN	5+
2807	1149	II. 748		12 8 49.1	3.009	4	42 8 55.2	20.04	5	pF; L; mE 45° 0; * n, p of 2	6+
2808		III. 718		12 8 55.0	3.006	1	41 5 2.2	20.04	1	vF; vS.....	1
2809	3385			12 9 7.3	3.126	4	132 32 44.2	20.04	4	pF; pL; pmE; vglbM.....	4
2810	1150			12 9 9.1	2.931	1	22 59 24.2	20.04	1	pB; S; R; psbM.....	1
2811	1151	I. 209		12 9 12.9	3.005	1	41 20 34.2	20.04	1	cB; pL; pmE 134° 4; psbM..	3
2812	1152	II. 137		12 9 17.5	3.065	1.:	82 31 31.9	20.03	1	pF; pL; R; r (? R.A. 10 ^m)...	3
2813	1153	II. 136		12 9 21.7	3.063	2	81 45 38.9	20.03	2	pB; pS; iE; gb, not M; r...	4
2814		II. 109		12 9 23.0	3.057	1	76 8 1.9	20.03	1	r.....	1*
2815	1154			12 9 25.1	3.084	1	101 31 49.9	20.03	1	F; eS; R; *170°, 60'.....	1
2816	1155			12 9 28.9	3.005	1	42 12 32.9	20.03	1	F; S; iE; f of 2.....	1
2817	1156	II. 518		12 9 31.4	3.030	2	55 42 6.9	20.03	2	F; vS; vE; psbM; sp of 2...	4
2818	1157			12 9 32.3	3.026	1	52 53 41.9	20.03	1	vF; L; R; gbM.....	1
2819	1158	II. 519		12 9 36.9	3.030	2	55 39 37.9	20.03	2	cF; vS; iE; psbM; nf of 2...	4
2820	3386			12 9 41.0	3.160	1	144 31 20.9	20.03	1	Cl; F; pL; iF; st 13...15...	1
2821	1159	II. 17		12 9 58.6	3.063	2	82 1 40.9	20.03	2	pB; pL; pmE; iM; p of 2...	5
2822	1160			12 9 59.5	3.067	1	85 32 30.9	20.03	1	pB; L; R; gbM.....	1
2823	1161	II. 496		12 9 59.5	3.063	1	81 35 59.9	20.03	1	pF; R; vsbMSN.....	2
2824	1162	II. 11		12 10 3.8	+3.054	2	73 54 23.9	+20.03	2	pB; pL; iE; vgbM; r.....	5

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1880, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s		° ' "	"			
2825	1163	V. 51		12 10 4.1	+2.894	1	19 48 0.9	+20.03	1	vF; eL; mE 160°±; vgbM..	3
2826	1164	III. 851		12 10 5.7	2.939	1	25 48 56.9	20.03	1	vF; pS; iR; vglbM.....	2
2827		III. 719		12 10 13.6	2.999	1	41 44 31.9	20.03	1	vF; vS; n of D neb	1
2828		III. 720		12 10 13.6	2.999	1	41 45 31.9	20.03	1	vF; vS; s of D neb.....	1
2829	1165	III. 480		12 10 17.5	3.063	2	82 32 50.9	20.03	2	vF; L; vgbM; *7 s	3
2830	1166	III. 725		12 10 20.5	3.003	2	43 35 49.9	20.03	2	vF; cL; iR; vgbM; r.....	4
2831	1167	V. 41		12 10 27.9	3.019	2	51 24 39.9	20.03	2	pB; vL; eE 43°±; vgbM ...	3
2832	1168	I. 74		12 10 32.5	3.033	3	59 36 54.9	20.03	3	cB; pL; vLE; smbM; r	5
2833		III. 91		12 10 43.7	3.063	1	82 6 1.9	20.03	1	eF	1
2834	1169	II. 742		12 10 48.8	2.995	2	41 49 7.9	20.03	2	vF; S; pinE; psbM.....	4
2835	1170	I. 264		12 10 57.2	2.864	1	18 25 17.9	20.03	1	pB; S; R; pgbM	2
2836	1171	I. 89		12 11 3.5	3.034	2	61 2 42.9	20.03	3	vB; S; E; vsvmbMN; *6.7 f 90°.	4
2837	1172	III. 702		12 11 30.9	3.029	1::	59 23 0.9	20.03	1::	vF; vS; R	2
2838	1173		M. 99	12 11 41.8	3.052	3	74 48 7.6	20.02	4	!!; {(H. h.)B; L; R; gbm; r} {(L) 3-branched spiral }	6+
2839	1174	II. 846		12 11 56.0	2.898	1	23 19 23.6	20.02	1	pB; L; cE 38°±; bMBN ...	2
2840			D'Arrest, 88	12 12 1	3.06	[1]	83 29 42	20.02	[1]	vF pS; R; *18 s 2'	0
2841	1175	V. 43		12 12 1.7	2.988	3	41 55 40.6	20.02	4	vB; vL; vmE 0°; sbMBN...	8+
2842	1176	II. 139		12 12 4.6	3.063	1	83 23 51.6	20.02	1	F; pS; R; gbm	4
2843	1177	II. 138		12 12 12.9	3.063	2	83 9 22.6	20.02	2	pB; E; psbM	5
2844	1178			12 12 13.0	3.064	1	83 53 0.6	20.02	1	neb; "1st of 5"	1
2845	1179	II. 110		12 12 22.0	3.051	1	74 20 51.6	20.02	1	B; S; R; r	3
2846		III. 535		12 12 26.0	3.090	1	101 28 31.6	20.02	1	vF; pL; iF	1*
2847	1180	II. 140		12 12 26.3	3.063	1	83 22 30.6	20.02	1	F; pS; R; gbm	4
2848	1181	II. 166		12 12 38.0	3.053	2	76 26 0.6	20.02	2	pB; vS; R; vsmbM.....	3
2849			D'Arrest, 89	12 12 41	3.06	[2]	83 12 42	20.02	[2]	pF; S; R; *9 f 1.7, n 85"...	0*
5070				12 12 41		...	83 46 42		...	See No. 5070	0
2850	1182	III. 299		12 12 45.0	3.024	1	58 52 49.6	20.02	1	cF; S; iR; gmbM	3
2851	1185	I. 75		12 12 47.4	3.025	1	59 36 15.6	20.02	1	vB; vL; E 90°±; mbMN ...	3
2852	1183	II. 568?		12 12 48.2	3.064	1	83 53 1.6	20.02	1	B; L; E; gbm	1*
2853		II. 804		12 12 48.3	2.946	1	32 29 1.6	20.02	1	pB; pL; iF	1
2854	1184	II. 376		12 12 49.1	3.029	1	61 36 0.6	20.02	2	F; S; vLE; gbm; *15 nr ...	3
2855	1186	I. 90	} =	12 13 3.4	3.025	2	59 56 25.6	20.02	2	vB; pL; R; mbM; r; p of 2	5*
		II. 322									
2856		II. 571??		12 13 8.7	3.062	1	82 51 1.6	20.02	1	4 neb sc about. Place of the last (see note).	1*
2857	1187	II. 573	} =	12 13 12.5	3.063	1	83 50 34.6	20.02	1	vB; vL; R; pgbM; "3 more seen."	3*
		II. 569?									
2858	1188	II. 323		12 13 18.6	3.024	2	59 54 20.6	20.02	2	B; S; R; bM; 2nd of 3	4
2859		II. 377		12 13 23.0	3.025	1	60 0 39.6	20.02	1		1*
2860		III. 798		12 13 25.7	2.933	1	31 6 1.6	20.02	1	cF; IE; p of 2.....	1
2861		III. 300		12 13 35.7	3.023	2	59 51 31.3	20.01	2	vF	2
2862	1189	II. 570?		12 13 35.9	3.063	1::	83 54 0.6	20.02	1::	vF; S	1*
2863	1188, a		R. nova?	12 13 38.6	3.024	...	59 51 20.6	20.02	...	Most probably = H. III. 300	0?
2864	1191	III. 726		12 13 41.0	2.980	3	42 56 17.3	20.01	3	vF; pS; R; vgbM; r	5
2865	1190	II. 571?		12 13 46.0	3.063	1	83 52 34.3	20.01	1	vB; R; central of 4.....	1*
2866	1193	II. 805		12 13 48.1	2.930	1	31 7 37.3	20.01	1	pB; L; R; gmbM	2
2867	1195	V. 5		12 14 4.2	3.042	2	70 50 17.3	20.01	2	F; vL; E; lbM; r	3
2868	1192	I. 275		12 14 4.8	2.720	3	13 51 7.3	20.01	4	pB; vS; R; lbM; 3 st f	6
2869	1194			12 14 6.1	3.063	2	83 49 57.3	20.01	2	B; pL; IE; bM; 4th of 4.....	3*
2870	1196			12 14 6.1	3.064	1	84 38 4.3	20.01	2	F; S; R; vglbM; B*340°, 60'	2+
2871	1197	II. 61		12 14 8.4	3.053	4	77 42 56.3	20.01	4	F; L; mE 135°±; bi-N; p of 2	6
2872		III. 92		12 14 23.6	3.061	2	82 34 1.3	20.01	2	vF; vS	2
2873		III. 93		12 14 23.6	3.061	2	82 34 1.3	20.01	2	cF; eS	2
2874	1198	II. 111		12 14 24.9	3.048	2	74 36 52.3	20.01	2	F; L; E 0°±; vgbM, p of 2	4
2875	1200	II. 62		12 14 32.7	3.053	2	77 43 1.3	20.01	2	F; L; IE; vgbM; f of 2	4
2876	1201	II. 572		12 14 33.1	3.063	1	83 50 23.3	20.01	1	F; IE; vgbM	2
2877	1199	II. 112		12 14 33.2	+3.047	2	74 36 53.3	+20.01	2	L; vmE 0°±; f of 2	4

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
2878	1202	I. 139	M. 61	12 14 45.1	+3.064	2	84° 44' 55".3	+20.01	2	vB; vL; vsbM*; biN	5*
2879	3387			12 14 51.8	3.132	1	122 41 54.3	20.01	1	vF; vL; R; vglbM; r	1
2880	1203			12 14 56.7	3.050	1	76 29 35.3	20.01	1	vF; R	1
2881	1204	I. 76		12 15 23.5	3.016	1	59 19 46.0	20.00	1	cB; L; E 150° ±; sbM; * np	3
2882	1205	II. 378		12 15 24.5	3.017	1::	60 0 34.0	20.00	2	B; cL; lE; np of 2	3
2883	1206			12 15 24.5	3.017	1	60 1 8.0	20.00	1	F; sf of 2	1
2884	1202, a		R. nova?	12 15 30	3.06	...	84 35 40	20.00	...	F; E; 10' nf h. 1202	0*
2885	1207	II. 63		12 15 30.2	3.051	2	77 26 25.0	20.00	2	vF; L; E 135° ±; r	4
2886	1209	II. 628		12 15 31.6	3.044	(1)	73 40 57.0	20.00	1	pB; cL; E; gbM	2
2887		II. 324		12 15 34.9	3.011	1	58 11 0.0	20.00	1	F; S	1
2888	1210	I. 276		12 15 36.3	2.686	3	13 53 59.0	20.00	3	pB; pS; vLE; sbM	5
2889	1208			12 15 37.4	3.057	1	81 1 20.0	20.00	1	eF; *8 n 5'	1
2890	1211		M. 100	12 15 50.6	3.043	3	73 23 54.0	20.00	4	!!; { (H, h) pF; vL; R; vg, psbMrN (L) 2-branched spiral }	5
2891	1212	II. 85		12 16 1.2	3.041	1	72 30 5.0	20.00	1	pB; S; R; psbM	2
2892			D'Arrest, 90	12 16 2	3.06	[4]	83 58 42	19.99	[4]	pB; R or lE; bM	0*
2893	1213	II. 141		12 16 2.0	3.060	1	83 8 51.0	20.00	1	vF; S; R; bM; 1st of 3	2
2894		II. 84		12 16 4.6	3.043	1	73 25 0.0	20.00	1	F; S; R; r	1
2895	1216	II. 847		12 16 6.6	2.843	1	23 22 43.0	20.00	1	pF; S; vLE; vgbM	2
2896	1214			12 16 8.6	3.093	1	101 45 29.0	20.00	1	vF; vS; R; bMN	1
2897	1217	II. 806		12 16 8.9	2.905	2	30 47 2.0	20.00	2	pB; S; E; gbM	3
2898	1220	III. 942		12 16 11.2	2.637	1	13 3 7.0	20.00	1	eF; E 0° ±	2
2899	1215	II. 142		12 16 12.5	3.060	1	83 11 1.0	20.00	1	F; pS; R; bM; 2nd of 3	2
2900	1218			12 16 16.0	3.058	1	81 45 7.0	20.00	1	pF; S; R; * v nr	1
2901	1219	II. 406		12 16 21.1	3.035	2	69 48 9.0	20.00	2	vF; pL; iR; biN?	4
2902	3388			12 16 23.3	3.231	1	147 20 30.0	20.00	1	Cl; pRi; lC; st 12...14	1
2903	1221	II. 86		12 16 23.7	3.040	1	72 31 5.0	20.00	1	cB; vS; mE; vsbM	2
2904	1222	II. 143		12 16 26.6	3.059	3	83 8 10.0	20.00	3	B; pL; R; bM; 3rd of 3	4
2905		III. 95		12 16 29.8	3.058	2	82 14 31.0	20.00	2	eF; vS; R	2
2906		III. 96		12 16 29.8	3.058	2	82 14 31.0	20.00	2	eF; vS; R	2
2907	1223	III. 94		12 16 30.1	3.060	1	82 16 59.0	20.00	1	pF; S; E; ? D	3
2908	1224	III. 31		12 16 32.7	3.038	1	71 41 1.0	20.00	1	vF; pS; R; vglbM; Δ 2 st ...	2
2909	1200, a		R. nova	12 16 34.7	3.053	::	77 43 1.3	20.00	::	vF; vmE	0
2910	1225	I. 210		12 16 34.8	2.960	5	42 14 8.0	20.00	6	vF; S; mE 100° ±; vsmbMBN	8+
2911	1226	II. 625		12 16 43.1	3.077	1	92 40 18.0	20.00	1	F; pL; E 70° ±; vlbM	4
2912	3389		Δ. 292	12 16 49.2	3.262	3	151 7 11.0	20.00	3	Cl; vB; vL; lC; st 12...14 ...	3
2913		III. 481		12 17 3.3	3.055	1	80 42 0.7	19.99	1	vF	1
2914	1230	III. 799		12 17 9.0	2.896	1	30 50 31.7	19.99	1	cF; cS; lE (? 18 ^m R.A.)	2
2915	1228	I. 123		12 17 9.3	3.061	1	84 17 31.7	19.99	1	B; S; * 8.9 sf 3'	3
2916	1229	III. 648		12 17 10.3	3.006	1	57 42 34.7	19.99	1	cF; pmE 90°; vlbM	2
2917	1231	I. 65		12 17 16.6	3.107	1	108 0 3.7	19.99	1	vB; L; R; vsmbMN; r	2
2918	1233	III. 800		12 17 17.0	2.895	1	30 51 31.7	19.99	1	vF; cS; R; r	2
2919		III. 938		12 17 17.6	2.659	1	14 17 0.7	19.99	1	eF; pL; iF	1
2920		III. 801		12 17 18.5	2.894	1	30 48 59.7	19.99	1	cF; cS; R	1
2921	1232	I. 30		12 17 21.3	3.057	2	81 54 17.7	19.99	3	cB; pL; vLE; gl; smbM	5
2922		III. 97		12 17 23.3	3.057	1	81 50 0.7	19.99	...	eF	2
2923		III. 38		12 17 35.0	3.050	1	78 37 59.7	19.99	1	vF; vS	1
2924	1234	I. 166		12 17 40.3	2.981	2	49 51 4.7	19.99	2	cB; S; R; mbMN; r	4
2925	1235	I. 22		12 17 45.8	3.048	3	77 31 34.7	19.99	3	B; pS; R; gbM	6
2926	1236	II. 144		12 17 46.2	3.057	2	81 46 52.7	19.99	2	pF; pS; lE; bM	3
2927	3390		Δ. 67??	12 17 50.1	3.412	1	161 53 19.7	19.99	1	⊕; pF; L; R; st 12...16 ...	2
2928	3391			12 17 55.9	3.162	1	128 58 12.7	19.99	1	pB; S; R; pgvmbM	1
2929	1227	II. 64		12 17 56.9	3.050	1	77 59 51.7	19.99	2	cF; cS; lE	3
2930	1237		M. 84	12 17 57.6	3.045	1	76 20 8.7	19.99	1	vB; pL; R; psbM; r	2
2931	1238	II. 379		12 17 59.0	2.994	2	60 40 3.7	19.99	2	F; S; R; bM; * nf 90"	3
2932	} 1237, a		R. 9 novæ	12 18 ±	+3.045	...	76 20 ±	+19.99	...	"Twelve knots exam." (see h. 1237, 1244, 1250).	0
2933											
2934											
2935											
2936											

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and ll.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	ll.		h m s	s		° ' "	"			
2937	1237, a		R. 9 novæ (continued)	12 18 ±	+3.045	...	76 20 ±	+19.99	...	"Twelve knots exam." (see h. 1237, 1244, 1250).	0
2938											
2939											
2940											
2941		II. 530		12 18 2.6	3.060	1	83 29 59.7	19.99	1	F; S;.....	1
2942	1239	I. 12		12 18 7.3	3.041	1	74 27 43.7	19.99	1	B; S; R; smbM	4
2943	1240	II. 87??		12 18 13.0	3.039	1	73 35 30.7	19.99	1	pS; R; psbMN	2
2944	1241			12 18 17.0	3.051	1	79 13 0.4	19.98	1	vF; S; R; lbM	1
2945		II. 743		12 18 19.0	2.940	1	40 23 59.4	19.98	1	F; S;.....	1
2946	1242		M. 85	12 18 19.5	3.033	2	71 2 10.4	19.98	2	vB; pL; R; bM; * np	3
2947	1243	III. 879		12 18 22.3	2.910	2	34 43 1.4	19.98	2	cF; S; iR	3
2948	1247	I. 277		12 18 36.7	2.609	2	13 42 3.4	19.98	2	B; cL; lC; psmbM	4
2949	1244			12 18 41.2	3.045	1	76 34 33.4	19.98	1	vF; E; p of 2	1
2950	1245	II. 749		12 18 41.2	2.950	3	43 32 16.4	19.98	3	pB; pL; iE; vglbM	5+
2951		II. 87		12 18 43.3	3.039	1	73 41 59.4	19.98	1	S; bM; r	1*
2952	1248	III. 852		12 18 44.3	2.819	2	24 17 14.4	19.98	2	cF; S; R; sbM; ** sp	3
2953	1249	III. 729		12 18 48.4	2.951	1	43 25 20.4	19.98	1	cF; S; R; vgbM	2
2954	1246	III. 361		12 18 48.5	3.010	1:	61 39 49.4	19.98	1	vF; vL; iF; B * p	2
2955	1250	II. 167		12 18 51.3	3.044	1::	76 29 1.4	19.98	1	Northern of 2; no description	2
2956	1250	II. 168		12 18 51.3	3.044	1::	76 29 1.4	19.98	1	Southern of 2; E	2
2957	1251	II. 55		12 18 52.0	3.032	2	71 0 52.4	19.98	2	pB; lE; bM	3
2958	1252	V. 29.1		12 18 52.0	2.993	1	55 40 35.4	19.98	1	eF; vL; np of D neb	3+
2959	1252, a		R. 2 novæ	12 18 ±	2.993	...	55 40 ±	19.98	...		0
2960											
2961											
2962											
2961	1253		M. 86	12 19 0.2	3.044	1	76 17 9.4	19.98	1	vB; L; R; gbMN; r	5*
2962	1252	V. 29.2		12 19 1.1	2.993	2	55 42 28.4	19.98	2	vF; vL; pvlbM; sf of D neb	3+
2963		III. 755		12 19 2.5	3.086	1	96 54 29.4	19.98	1	vF; vS; E	1
2964		III. 756		12 19 2.5	3.086	1	96 54 29.4	19.98	1	vF; vS; E	1
2965			Auw. N. 30	12 19 2.6	3.041	A	76 6 32.4	19.98	A	F; L; mE 90° (Auwers, Mar. 5, 1862).	0
2966	1254	II. 88		12 19 3.4	3.037	1	73 3 10.4	19.98	1	pF; S; R; vsbM; r	2
2967		III. 39		12 19 5.0	3.050	1	78 42 59.7	19.98	1	vF; B * nr	1
2968	1255			12 19 11.1	3.044	1	76 34 59.4	19.98	1::	f of 2 neb	1
2969	1256			12 19 14.6	3.052	1	80 12 44.4	19.98	1	eF; vL; R; gbM	1
2970		III. 17		12 19 17.0	3.065	1	86 43 59.4	19.98	1	vF; pS; r	1
2971	1257	II. 34		12 19 26.2	3.063	2	85 15 53.4	19.98	2	F; pL; R; gbM; r	6
2972	1258	I. 77		12 19 29.4	2.997	1	57 30 19.4	19.98	2	vlB; L; E; g, vsmbM*	5+
2973		III. 482		12 19 30.1	3.053	1	80 47 0.4	19.98	1	eF	1
2974	1259	II. 169		12 19 32.4	3.044	1	76 39 31.4	19.98	1	cF; S; gbM; 2 st n, np	2
2975	1260			12 19 37.2	3.054	1	81 18 20.1	19.97	1	vF; L; R; * sp 5'	1
2976	1261	III. 492		12 19 44.6	3.072	1	90 6 39.1	19.97	2	{ H. vF; cL; mE } { h. F; S; R; * nr }	3*
2977	1262	II. 113		12 19 49.3	3.038	1	74 10 51.1	19.97	1	B; pmE 135° ±; sbM	3
2978	1263	II. 23		12 19 50.7	3.065	2	86 43 44.1	19.97	2	F; pL; lE; r; (?=III. 17)...	4
2979		II. 155		12 19 56.1	3.052	1	79 37 0.7	19.97	1	F; pL; E; lbp	1
2980	1265	III. 114		12 19 59.7	3.083	2	95 3 10.1	19.97	2	F; vS; R; psbM; 2S st nr	5
2981	1264	II. 89		12 19 59.8	3.037	2	73 45 10.1	19.97	2	pB; pL; pgbM; B * np	5
2982	1266	II. 145		12 20 1.6	3.058	1	83 20 42.1	19.97	1	vF; vS; E	3
2983	1267	II. 170		12 20 5.0	3.043	2	76 30 10.1	19.97	2	pF; S; R; bM	3
2984	1268	II. 171		12 20 8.2	3.044	1	76 55 44.1	19.97	1	vF; vS; cE; gbM	2
2985	1269			12 20 16.0	3.088	1	97 24 14.1	19.97	1	vF; pL	1
2986	1270	II. 146		12 20 19.4	3.057	2	82 58 0.1	19.97	2	cF; L; R; gbM	3
2987	1271	II. 65		12 20 20.7	3.046	2	78 7 49.1	19.97	2	B; L; cE; psbM; * 10 nf	3
2988	1272	II. 172		12 20 22.2	3.043	1	76 53 54.1	19.97	1	cF; S; gbM	2
2989		II. 497		12 20 25.0	3.053	1	81 0 59.1	19.97	1	pF; vS	1
2990	1273			12 20 26.1	3.089	1	97 30 49.1	19.97	1	pF; pL; lE	1
2991	1274	I. 28, 1		12 20 33.7	3.041	2	76 9 14.1	19.97	3	vB; cL; R; p of 2	2
2992	1274, a		R. nova	12 20 ±	3.041	...	76 9 ±	19.97	...	See note	0*
2993	1276	II. 173		12 20 35.2	3.043	1	76 55 49.1	19.97	1	B; pS; R; bM; r	2
2994	1275	I. 28, 2		12 20 36.7	3.041	3	76 13 17.1	19.97	3	B; cL; vLE; r; f of 2	5
2995	1275, a		R. nova	12 20 ±	+3.041	...	76 13 ±	+19.97	...	See note	0*

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	"			
2996	1277			12 20 39.3	+3.070	1	89 5 41.1	+19.97	1	F; eE 75°; *10 nf; place... that of *).	1
2997	3392		Δ. 300?	12 20 45.1	3.287	1	149 19 6.8	19.96	1	Cl; S; st 11...12	1
2998		II. 848		12 20 48.0	2.795	1	24 25 19.8	19.96	1	F; S; iR; bM	2
2999	1279	II. 156		12 20 58.0	3.048	1	79 24 53.8	19.96	1	vb; pL; R; smbM	2*
3000	3393			12 21 7.5	3.191	1	132 29 6.8	19.96	1	eF; L; R; vgbM	1
3001	1280	I. 91		12 21 15.0	2.999	3	60 36 32.8	19.96	4	B; L; E 90°; sbM	6
3002	1281	I. 213		12 21 17.5	2.944	4	45 8 8.8	19.96	4	vB; cL; mE 15°; rrr; * 9, 5'	5
3003	1282	{ II. 56		12 21 24.9	3.030	6	72 8 33.8	19.96	7	B; L; R; gymbM*; r; B* nr	10*
		= II. 90									
3004	1283	II. 26		12 21 39.0	3.055	1	82 42 49.8	19.96	1	F; pS; bM; r	2
3005	1284	II. 180		12 21 40.2	3.075	1	91 10 10.8	19.96	1	F; L; R; gbM; er	5
3006	1285	II. 355		12 21 41.6	3.014	2	66 24 25.8	19.96	2	F; L; E; gbM; 2 Bst nf....	3
3007	3394			12 21 43.2	3.147	1	119 19 41.8	19.96	1	eeF; vS; * 13 att	1
3008		I. 23		12 21 49.1	3.043	2	77 32 29.8	19.96	2	pB; S; vinE	2*
3009	1286	II. 35		12 21 50.2	3.063	3	85 39 10.8	19.96	3	eB; pS; R; smbMN	5
3010	1287	II. 121		12 21 51.6	3.039	1	75 59 6.8	19.96	1	pB; S; R; bM; p of 2	2
3011	1289	{ I. 212		12 21 52.8	2.936	1	44 20 29.8	19.96	1	B; pL; E 45° ±; psbM	4*
		= II. 750									
3012	1288	I. 161		12 21 54.1	3.037	2	75 14 55.5	19.95	2	pB; pL; iR; bM; r; * 8 sf 2'	3
3013	1290	{ II. 122		12 21 57.1	3.039	1	76 1 41.5	19.95	2	pF; S; R; bM; f of 2.....	4*
		= II. 174									
3014	3396	III. 764		12 22 2.0	3.128	2	112 23 40.5	19.95	2	pB; pS; E 130°; vbM	3
3015	3395			12 22 3.2	3.350	1	154 1 1.5	19.95	1	Cl; P; vIC	1
3016	1291			12 22 4.3	2.778	1	24 25 41.5	19.95	1	pB; R; gbM	1
3017		II. 630		12 22 15.1	3.037	1	75 15 58.5	19.95	1	cL	1
3018	1292	III. 483		12 22 15.4	3.051	1	81 4 3.5	19.95	1	F; vS; R; pgbM.....	2
3019		II. 157		12 22 25.8	3.049	1	80 23 58.5	19.95	1	pF; pL; mE; bM; r	1
3020	1293	{ II. 18		12 22 30.4	3.049	2	81 24 5.5	19.95	2	F; pL; iR; bM	6
		= II. 498									
3021	1294		M. 49	12 22 39.3	3.051	4	81 13 44.5	19.95	5	vB; L; { H. E h. R } ; mbM	6*
3022	1294, a		R. 3 novæ	12 22 ±	3.051	...	81 13 ±	19.95	...	"Four found" (one being h. 1294).	0
3023				12 22 ±	3.051	...	81 13 ±	19.95	...		0
3024					12 22 ±	3.051	...	81 13 ±	19.95	...	
3025		II. 115		12 22 43.3	3.016	2	75 36 0.5	19.95	2	vB; cL.....	2
3026	1295	{ II. 117		12 22 48.2	3.036	1	75 9 20.5	19.95	1	pF; R; r	3*
		= II. 629									
3027	1297	III. 362		12 22 48.7	2.998	1	61 58 28.5	19.95	2	eF; pL; R	5
3028	1296	II. 123		12 22 51.8	3.040	2	76 54 3.5	19.95	2	F; S; R; bM; 1st of 3	3
3029		II. 116		12 22 58.9	3.037	2	75 39 0.5	19.95	2	pB; pL.....	2*
3030		II. 114		12 23 1.6	3.037	2	75 48 33.5	19.95	2	vF; r	2
3031	1298	II. 124		12 23 11.1	3.040	4	76 54 41.2	19.94	5	pB; S; R; psbM; 2nd of 3...	7
3032	1299	II. 531		12 23 17.7	3.060	1	84 58 43.2	19.94	1	pF; pS; E; bs	3
3033		III. 40		12 23 33.4	3.043	3	78 28 58.2	19.94	3	eF; pL	1
3034	1300			12 23 37.1	3.100	1	100 51 52.2	19.94	1	pF; S; R; gbM	1
3035	1301		M. 87	12 23 44.0	3.039	4	76 50 39.2	19.94	4	vB; vL; R; mbM	7
3036		II. 776		12 23 44.2	3.091	1	97 18 28.2	19.94	1	F; vL; er.....	1
3037	1302	III. 484		12 23 44.6	3.049	1	80 51 38.2	19.94	1	vF; vS; iE	2
3038	1303	II. 91		12 23 48.1	3.027	3	72 27 55.2	19.94	3	pF; cS; R; gbM.....	4
3039	1304	III. 41		12 23 50.0	3.041	1	77 45 3.2	19.94	1	F; L; R	2
3040	1305	II. 499		12 23 53.2	3.050	1	81 9 14.2	19.94	1	pF; pL; vglbM; 2 st nr	2
3041	1306	I. 197		12 24 17.8	2.938	1	47 36 37.9	19.93	1	B; pS; iR; p of 2	2†
3042	1308	I. 198		12 24 23.6	2.938	1	47 39 25.9	19.93	1	vB; vL; mE 130°; rr	2†
3043	1307	I. 83		12 24 24.3	2.998	1	63 26 55.9	19.93	1	vB; pL; R; vsmbMN.....	2*
3044	1310	III. 301		12 24 26.4	+2.987	1	60 5 7.9	+19.93	2	pF; cS; R; pslbM	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
3045	1309	II. 36		12 24 29.8	+3.060	3	85° 17' 47.9	+19.93	3	F; cL; biN or D neb	6
3046		III. 42		12 24 33.4	3.040	1	77 37 57.9	19.93	1	vF	1
3047	3397			12 24 40.2	3.195	2	129 12 18.9	19.93	2	vF; L; R; vglbM	2
3048	1311	I. 234		12 24 41.5	2.827	1	31 16 14.9	19.93	1	B; cS; E; pgbM; * 9 f 30'' ..	3
3049	1312		M. 88	12 24 54.5	3.031	3	74 48 26.9	19.93	3	B; vL; vmE; p of D neb.....	8
3050		II. 118		12 24 +	3.031	...	74 48 -	19.93	...	F; S; f of D neb (not ob- by h.).	
3051		III. 69		12 24 57.5	3.024	1	72 25 57.9	19.93	1	vF; vS.....	1
3052	1313	II. 66		12 24 58.5	3.040	2	78 3 15.9	19.93	2	pB; S; R; gbM	3
3053	1314	II. 92		12 24 59.4	3.025	1	72 32 23.9	19.93	1	vF; S	2
3054	3398	II. 771		12 25 4.1	3.090	1	96 46 40.6	19.92	1	pB; cL; iE; gvlbM; er	3
3055	1315	III. 18		12 25 6.8	3.060	1::	85 14 53.6	19.92	1	vF; cL; r; f of 2.....	2
3056	1316	II. 631		12 25 6.8	3.034	1	75 48 23.6	19.92	1	cF; pmE90°±; gbM; *9 f8°	2
3057	3399			12 25 9.9	3.197	2	129 8 6.6	19.92	2	pB; S; R; psmbM*16	2
3058	1317			12 25 9.9	3.055	1	83 24 1.6	19.92	1	vS; R; sbM*13	1
3059	1318			12 25 14.6	2.974	1	57 7 59.6	19.92	1	vF; S; R; lbM	1
3060	1196, a		R. nova	12 25 30	3.058	???	84 47	19.92	???	Query R.A.; vF; 10's of scar- let *.	0
3061	1319	III. 834		12 25 33.9	2.833	1	32 45 54.6	19.92	1	pF; vS; iR; vgbM	2
3062	1321			12 25 45.5	2.748	1	25 29 50.6	19.92	1	pB; S; R; psbM.....	1
3063	1320	III. 302		12 25 50.1	2.980	2	59 30 52.6	19.92	2	cF; vS; R; bM	3
3064	1322			12 26 3.1	3.049	1	81 22 31.6	19.92	1	F; S; R; bM	1
3065	1323	III. 78		12 26 3.1	3.029	1	74 38 42.6	19.92	1	F; pS; R; r.....	3
3066		IV. 5		12 26 3.3	3.070	3	89 8 56.6	19.92	3	cB; vL; vmE95°±; B * in cont.	3
3067	1324	II. 93		12 26 4.0	3.024	1	72 56 23.6	19.92	1	F; vS; bM*	2
3068		II. 158		12 26 20.2	3.046	3	80 33 36.3	19.91	3	F; pL; R; bM; r	3
3069		II. 849		12 26 23.0	2.732	1	25 12 55.5	19.91	1	pB; vS; iE; sbMSN	1
3070		II. 757		12 26 24.2	3.091	2	96 36 57.3	19.91	2	vF; S; 2 vS st inv	2
3071	1326			12 26 26.9	2.741	1	25 37 18.3	19.91	1	pB; S; pmE; pgbM; * 9 inv ..	1
3072	1325			12 26 32.9	3.044	1	80 2 47.3	19.91	1	cF; pL; iE; vlbM	1
3073	1327			12 26 38.0	3.104	1	101 14 26.3	19.91	1	vF; iF; bM.....	1
3074	1328	II. 325		12 26 53.3	2.975	1::	58 57 57.3	19.91	1::	F; pL; iR; bM	3
3075	1329 {	I. 31	{	12 26 56.3	3.048	3	81 31 57.3	19.91	3	vB; vL; mE120°±; psmbM; L * f; * 9 p.	5*
		I. 38									
3076	1330	II. 37		12 27 0.4	3.063	2	86 34 26.3	19.91	2	pB; L; pmE60°±; mbM ...	4
3077	1331	II. 67		12 27 2.1	3.037	3	77 54 52.3	19.91	4	pF; cS; R; bM; * 9 f 30° ...	5
3078		III. 26		12 27 2.2	3.009	1	68 41 56.3	19.91	1	eF; L	2*
3079	1332		8 Canum	12 27 7.7	2.925	4	47 52 34.0	19.90	4	Nebulous *	4*
3080		II. 500		12 27 8.5	3.047	1	81 1 56.0	19.90	1	vL; er	1
3081	1333	II. 175		12 27 10.6	3.032	2	76 9 20.0	19.90	2	F; pL; R; vgbM	3
3082	1334	II. 147		12 27 11.7	3.051	1	82 46 13.0	19.90	1	pB; pL; pmE; vgbM; r	2
3083	1336	II. 410		12 27 13.1	2.952	2	53 42 9.0	19.90	2	cF; L; iE; vglbM; r	4
3084	1335 {	II. 94	{	12 27 14.9	3.024	2	73 40 25.0	19.90	2	F; pS; bM; r	5
		II. 119									
3085	1337	V. 2		12 27 19.6	3.064	1	87 2 47.0	19.90	1	B; vL; mE110°; sbM; er ...	5†
3086	1338			12 27 32.8	3.015	1	71 1 23.0	19.90	1	pB; pmE	1
3087	1341			12 28 6.8	2.860	1	38 25 19.7	19.89	1	eF; pL; R	1
3088	1340			12 28 12.2	3.052	1	83 6 37.7	19.89	1	pF; cS; R; bM	1
3089	1346	II. 850		12 28 13.8	2.721	1	25 42 2.7	19.89	1	F; L; iR; vgbM; S * nf.....	2
3090	1342	III. 493		12 28 15.1	3.071	1::	89 28 3.7	19.89	1:	eF; S; R; gbM	3
3091	1344	III. 802		12 28 15.7	2.783	2	30 19 21.7	19.89	2	vF; pS; E; vgbM; * 9 f 2'; p of 2.	4
3092	1339	I. 160		12 28 17.8	3.081	4	93 1 7.7	19.89	4	vB; cL; pmE63°±; vsmbMN.	6
3093	1345	II. 120		12 28 22.7	3.026	3	74 43 45.7	19.89	3	B; L; iE; lbM	5
3094	1347	III. 807		12 28 23.5	2.781	2	30 17 43.7	19.89	2	eF; pS; E; f of 2	3
3095	1343	I. 36		12 28 24.2	3.033	2	77 0 46.7	19.89	3	pB; S; vLE; sp of 2.....	4
3096	1349	I. 37		12 28 32.3	+3.033	2	76 58 38.7	+19.89	2	pB; S; R; bM; nf of 2	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. II.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
3097	1348		M. 89	12 28 35.2	+3.032	1	76° 40' 32.7	+19.89	4	pB; pS; R; gmbM	5*
3098	3400			12 28 36.9	3.211	1	128 39 49.7	19.89	1	F; vLE; glbM	1
3099	1350	II. 343		12 28 44.3	2.983	2	62 42 18.7	19.89	2	B; pS; iR; vsmbM * 12	3
3100	1351	II. 380		12 28 50.3	2.981	1	62 18 47.7	19.89	2	F; pL	3
3101	1352	I. 92		12 28 59.6	2.980	3	61 16 7.4	19.88	3	vB; vL; mE 150°; gbM; 3 st f	4†
3102	1354			12 29 0.3	2.980	1	62 16 6.4	19.88	1	vF; nf of 2 or 3	1
3103	1353	I. 119		12 29 0.4	3.046	1::	81 33 21.4	19.88	1:	cB; pL; R; gbM	2*
3104	1355	II. 407		12 29 5.7	3.008	2	69 54 20.4	19.88	2	pB; pL; vLE; lbM; r	4
3105	1356	II. 68		12 29 21.6	3.034	2	77 47 21.4	19.88	2	pB; S; iE; psbM	4
3106	1357	V. 24		12 29 22.9	2.993	4	63 14 30.4	19.88	4	B; cL; eE 136° 1; vsbMN = * 10, 11.	5†
3107	1360	III. 880		12 29 26.1	2.821	1	35 0 1.4	19.88	1	pF; S; iR; gbM	2
3108	{ 1358 = 1363 1359	IV. 8		12 29 26.5	3.035	4	77 59 26.4	19.88	4	vF; L; np of D neb } pos 160° ±	6*†
3109	{ 1363 = 1363	IV. 9		12 29 28.0	3.035	2	78 0 26.4	19.88	2	vF; L; sf of D neb } pos 160° ±	6*†
3110	1361	I. 32		12 29 45.6	3.047	5	81 59 3.1	19.87	5	cB; pS; mE 0° ±; sbMrN ...	9
3111			M. 90	12 29 52.8	3.028	2	76 4 18.1	19.87	2	pL; bMN	2*
3112	1364	III. 939		12 29 55.6	2.392	1	14 59 52.8	19.86	1	eF; S	2
3113	1362	III. 602		12 29 59.1	3.024	1	74 58 8.1	19.87	1	vF; L; E; vgbM; cB * att... ..	2†
3114	3401			12 30 6.6	3.241	1	132 51 14.1	19.87	1	vF; S; * 10 n 30'	1
3115	3402			12 30 17.6	3.199	1	124 44 3.1	19.87	1	vF; L; iE; vglbM	1
3116	3403			12 30 19.4	3.224	1	129 45 59.1	19.87	1	F; S; pmE; 2 st p	1
3117		III. 13		12 30 23.9	3.051	1	83 9 55.8	19.86	1?	vF; vS	1
3118	1365	II. 15		12 30 26.2	3.039	1	79 40 5.8	19.86	1	pF; pS; R; sbMN; * np ...	3
3119	1366			12 30 26.7	3.039	1	79 45 21.8	19.86	1	F; R; bM (? = IL 15 + 5' P.D.) ..	1
3120	1367		M. 91 ??	12 30 30.8	3.025	1?	75 26 55.8	19.86	1?	np this place is a F neb; not M. 91, whose existence? ..	1
3121	1368		M. 58	12 30 36.6	3.031	3	77 24 52.8	19.86	3	B; L; iR; vmbM; r	6
3122	1369	I. 124		12 30 40.5	3.053	1	83 51 55.8	19.86	1	pB; L; vgbM	3
5071				12 31 0.5		...	89 2 46.5		...	See No. 5071.	
3123	1370	III. 495		12 31 14.1	2.945	2	55 46 15.8	19.86	2	cF; S; iE; bM	3
3124			D'Arrest, 91	12 31 17	3.02	[3]	76 7 12	19.85	[3]	vF; S; R	0
3125	1371	I. 125		12 31 18.5	3.056	2	84 54 40.5	19.85	2	pB; L; E; psbM	4
3126		III. 98		12 31 45.7	3.047	1::	82 25 53.5	19.85	1::	vF; eS	1
3127	1374	I. 273		12 31 53.3	2.369	3	15 2 24.5	19.85	4	cB; L; iE; pgmbM	7*
3128	3404		M. 68	12 32 5.1	3.166	1	115 58 45.2	19.84	1	⊕; L; eRi; vC; iR; rrr; st 12, red.	4
3129	1372	III. 504		12 32 6.7	3.050	1	83 12 29.2	19.84	1	vF; cS	3
3130	1373	II. 31		12 32 8.0	3.072	1	89 16 17.2	19.84	1	F; L; E 90° ±; vgbM	4
3131	1375	II. 183		12 32 26.6	3.088	2	94 34 23.2	19.84	2	pB; cL; E; sbMN = *	4
3132	1376	I. 43		12 32 44.2	3.110	1	100 50 14.2	19.84	1	!; vB; vL; eE 92°; vsmbMN.	3†
3133	1377	II. 632		12 32 49.6	3.016	3	73 56 8.2	19.84	4	pF; pL; R; gbM	5
3134	1378	I. 24		12 32 50.8	3.034	5	79 2 56.2	19.84	5	B; pS; R; gmbM; r; 3 st f... ..	7
3135		II. 636		12 32 53.7	3.089	1	95 2 22.9	19.83	1	F; vL; bM	1
3136		III. 105		12 33 11.7	3.040	2	80 51 21.9	19.83	2	eF; L; R; vlbM	2
3137		III. 509		12 33 13.9	3.065	1	88 0 51.9	19.83	1	vF; vS	1
3138	1379	II. 577		12 33 14.1	3.059	2	86 6 33.9	19.83	2	F; S; R; 2 st 8 f	3*
3139	3405			12 33 23.5	3.242	1	130 8 57.9	19.83	1	eF; L; R; psbM; p of 2	1
3140	1380	II. 184		12 33 26.5	3.088	1	94 21 50.9	19.83	1	F; L; E; vglbM	3
3141	3406			12 33 31.0	3.242	1	130 12 2.6	19.82	1	F; L; R; vgbM; r	1
3142	1381	I. 254		12 33 42.8	2.689	1	27 36 49.6	19.82	1	B; L; vmE 118° 6; glbM ...	2
3143	1382	III. 43		12 33 53.3	3.027	2	77 20 9.6	19.82	3	vF; pS; E; 2 or 3 vS st inv ..	5
3144	1383	II. 69		12 34 10.3	3.033	2	79 3 58.6	19.82	2	pB; pL; R; psbM; r; * 12 np 1'.	4
3145	3407		Δ. 272	12 34 13.2	3.463	1	152 12 8.6	19.82	1	Cl; pL; pC; cE; st 10	1
3146		I. 7		12 34 16.3	3.042	1::	81 24 51.6	19.82	1::	vB; vL (no doubt a comet) ..	1
3147		II. 19		12 34 20.3	3.042	1	81 30 51.6	19.82	1	F; vL	2
3148	1384	II. 148		12 34 23.9	+3.043	3	81 55 10.3	+19.81	3	pB; S; R; psmbM	6*

No. of Catalogue.	References to			Right Ascension for 1800, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1800, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. II.'s Catalogues of Nebulæ.	Sir W. II.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s						
3149	3408			12 34 44.6	+3.247	1	129 54 10.3	+19.81	1	eF; vS; R; * att nf; p of 2...	1
3150		II. 744		12 34 46.0	2.815	2	38 48 50.3	19.81	2	pF; S; iR; er	2
3151	1385 {	I. 178	}	12 34 48.0	2.886	4	48 4 47.3	19.81	4	{ B; L; R; mbM } D neb; { pb } pos 160°	5†
3152		I. 179									
3153	1387			12 34 56.3	3.021	1	76 17 9.3	19.81	1	vF; S; R; vgbM	1
3154	1388	II. 411		12 34 56.5	2.922	3	54 9 54.3	19.81	3	F; pS; R; lbM; * 8.9 f	5
3155	1386		M. 59	12 34 56.9	3.026	3	77 34 1.3	19.81	3	B; pL; iE; vsvmbM; 2 st p ..	5
3156	3409			12 35 3.9	3.249	1	129 58 30.3	19.81	1	pF; S; R; psbM; f of 2	1
3157	1389	II. 149		12 35 3.9	3.041	1	81 33 55.3	19.81	1	cF; pL; E; psbM; r	5
3158	1390			12 35 7.3	3.059	1	86 10 20.3	19.81	1	B; E	1
3159	1391	II. 659		12 35 8.7	2.935	1	56 39 40.3	19.81	1	F; S; R; np of 2	2
3160	1392	II. 660		12 35 9.8	2.883	2	47 56 36.3	19.81	2	pF; S; R	3
3161	{ 1393 = 3410 }	II. 772		12 35 10.4	3.095	2	96 16 10.3	19.81	2	vF; cS; iE; glbM	4
	{ 1394 = 3411 }										
3162		II. 773		12 35 11.3	3.095	2	96 11 25.3	19.81	2	cF; S; E; gbM	4
3163			D'Arrest, 92	12 35 21	3.07	[1]	91 2 12	19.80	[1]	pB; pL; E; lbM; ? biN	0
3164	1395	II. 532		12 35 21.7	3.055	1	85 16 24.0	19.80	1	cF; S; R; lbM	3
3165	1397	V. 42		12 35 22.0	2.934	2	56 41 21.0	19.80	2	!; vB; vL; eE70°±; bMN; B*nr.	3†
3166	1396	I. 14		12 35 23.8	3.070	1	89 18 56.0	19.80	1	pB; L; E45°±	4
3167	1398	III. 603		12 35 38.1	3.015	2	74 55 47.0	19.80	2	vF; L; mE135°±; vgbM ...	3
3168	1400			12 35 39.8	2.992	2	69 17 27.0	19.80	2	vF; L; vglbM	2
3169	1399	II. 38		12 35 41.1	3.060	2	86 32 30.0	19.80	2	B; L; iR; vgvmbM; r	4
3170	1401			12 35 41.9	3.055	1	85 32 15.0	19.80	1	vB; cS; R; smbM	1*
3171	1402 {	II. 70	}	12 35 43.8	3.026	1::	77 49 6.0	19.80	1	F; R; gbM	3
		= II. 176									
3172	1402, a		R. nova	12 35 43±		...	77 49 ±		...	Makes a D or biN neb with h. 1402.	
3173	1403	II. 125		12 35 49.4	3.019	1	75 58 44.0	19.80	1	pB; S; E; r; * 12 sf 1'	2
3174		II. 20		12 35 56.9	3.043	1	81 53 10.3	19.79	1	vS	1*
3175		III. 494		12 36 3.7	3.072	2	89 53 20.7	19.79	2	vF; cS; E	2
3176	1402	I. 10		12 36 12.1	3.062	1	87 14 46.7	19.79	1	cB; pS; iE; mbM	5
3177	1406 {	II. 794	}	12 36 15.9	2.754	1	34 4 23.7	19.79	1	vF; S; R; gbM	1*
		No. 1									
3178	3412			12 36 21.6	3.262	1	130 58 46.7	19.79	1	pB; S; psbM	1
3179	1407 {	II. 794	}	12 36 23.0	2.756	1	34 22 23.7	19.79	1	F; S; 4 vS st sp	2*
		No. 2									
3180	1405	III. 44		12 36 25.5	3.025	4	77 39 25.7	19.79	3	vF; pL; iE115°±; np of D neb	6*†
3181	1410	I. 274		12 36 33.0	2.258	4	14 48 37.7	19.79	5	pB; cS; R; gbM; * p	8
3182	1408		M. 60	12 36 34.8	3.025	5	77 40 38.4	19.78	6	vB; pL; R; f of D neb	8†
3183	3413			12 36 41.2	3.256	1	129 57 35.4	19.78	1	vF; R; bM; r	1
3184	1409	II. 12		12 36 42.3	3.005	3	72 50 28.4	19.78	4	cB; L; E90°; gbM; r	6
3185	1413			12 36 48.8	2.698	1	30 16 19.4	19.78	1	pF; pL; gbM; 2 B st f	1
3186		III. 662		12 36 50.2	3.072	1	89 47 49.4	19.78	1	vF; pL	1
3187	1411	II. 126		12 36 54.0	3.018	2	76 7 8.4	19.78	2	F; vL; pmE; ?D; 3 st nr ...	4
3188	1412	II. 661		12 36 54.5	2.876	2	48 12 25.4	19.78	3	vF; vS; stellar; * 15, 16 f ...	3
3189	1414	I. 176		12 37 7.8	2.929	4	57 3 48.4	19.78	5	!; pB; L; vmE34°3; sp of 2	6*†
3190	1415	I. 177		12 37 16.6	2.929	3	57 1 24.1	19.77	3	!; pF; L; E90°±; nf of 2 ...	6*†
3191	3414	II. 558		12 37 23.1	3.109	1	99 18 53.1	19.77	1	vF; L; E; * 16 att; * 9 p ...	2
3192	1416	II. 127		12 37 27.8	3.016	1	75 43 35.1	19.77	1	F; cS; R; bM; r	3
3193	1417	II. 71		12 37 29.4	3.025	1	78 2 34.1	19.77	1	vB; S; vsvmbMN	2
3194	3415			12 37 37.9	3.264	1::	130 19 53.1	19.77	1::	F; pL; R; gbM	1
3195	1418	II. 643		12 37 42.5	2.898	1	52 5 58.1	19.77	1	vF; pL; R; gbM; r	2
3196		II. 39		12 37 46.2	3.057	1	86 0 49.1	19.77	1	pB; 2 S st in M; S * p	1
3197	1419	I. 142		12 37 58.3	3.057	2	86 11 14.8	19.76	2	B; pL; iR; mbM; * 10 sp ...	3
3198	1420	I. 15		12 37 59.0	3.072	1	89 41 29.8	19.76	1	B; vL; mE45°±; psbM ...	4
3199	1421			12 38 12.3	+3.023	1	77 47 31.8	+19.76	1	B; S; R; psbM	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos	Other Authorities.								
	h.	II.		h m s	s						
3200		III. 663		12 38 15.2	+3.072	1	89 46 47.8	+19.76	1	vF; S; iF.....	1
3201	1422	III. 328		12 38 23.9	2.952	3	62 6 37.8	19.76	4	pF; cS; R; bM; r; p of 2 ...	6
3202	1423	II. 774		12 38 32.4	3.098	1	96 18 3.8	19.76	1	pF; S; R; psmbM	3
3203	3416			12 38 36.0	3.273	1	130 56 55.5	19.75	1	eF; S; R; vgbM.....	1
3204	1424	III. 329		12 38 43.7	2.951	1	62 10 24.5	19.75	2	F; vS; R; sbM*10; f of 2 ...	4
3205	3417			12 38 47.3	3.104	1	97 52 44.5	19.75	1	vF; cS; R; glbM	1
3206		III. 778		12 39 11.1	2.733	2	34 28 43.5	19.75	2	cF; S; iE.....	2*
3207	1425	II. 326		12 39 17.7	2.928	1	58 30 6.2	19.74	1	vF; pmE; ?biN	3
3208	3418			12 39 18.0	3.274	1	130 49 55.2	19.74	1	eF; iE; vgbM.....	1
3209	3419			12 39 36.0	3.263	1	128 48 5.2	19.74	1	eeF; pL; R.....	1
3210	3420			12 39 36.4	3.118	1	100 52 34.2	19.74	1	eF; S; 1 or 2 st inv	1
3211	3421			12 39 45.5	3.291	1	132 34 50.2	19.74	1	pF; S; R; gbM	1
3212	3423	III. 523		12 39 58.2	3.111	1	99 17 29.9	19.73	1	cF; L; E 45°±; gvlbM	2
3213	3422			12 40 2.9	3.279	1	130 47 24.9	19.73	1	eF; pS; R; vgbM; S * sp ...	1
3214	1426	II. 181		12 40 6.3	3.081	1	91 58 8.9	19.73	1	B; pL; pmE 25°	5*
3215	1427	III. 398		12 40 13.3	2.984	3	69 46 52.9	19.73	3	F; S; R; sbM*; rr.....	4
3216	1428	II. 795		12 40 16.6	2.728	1	34 41 11.9	19.73	1	pF; vS; vmE; vsmbM	3*
3217	1430			12 40 39.3	2.897	1	53 52 53.6	19.72	1	vF; vS; R; psbM	1
3218	1429	III. 543		12 40 43.4	3.051	1	84 53 57.6	19.72	2	eF; pL; *9.10 p 10°	3
3219	1431	II. 128		12 40 44.5	3.010	3	75 28 34.6	19.72	3	pB; vL; E; vglbM; r.....	5
3220		III. 664		12 40 48.3	3.076	1	90 55 44.6	19.72	1	vF; S	1
3221	1432	II. 182		12 40 58.2	3.081	1	92 33 48.6	19.72	1	pB; pL; E 90°±; mbM.....	5
3222	1433	II. 381		12 41 2.8	2.943	2	62 0 41.6	19.72	3	F; cS; R; bM.....	4
3223		III. 906		12 41 5.5	2.332	1	18 3 44.6	19.72	1	vF; pL; E	1
3224	1435	II. 796		12 41 9.5	2.731	1	34 51 21.6	19.72	1	cF; pS; vIE; mbMN	3*
3225	1434	II. 72		12 41 11.2	3.021	6	78 14 47.6	19.72	6	pF; S; vIE	7
3226	3424		Δ. 510 ?	12 41 12.2	3.282	1	130 32 29.3	19.71	1	pB; L; R; gbM; r	1
3227	1436	I. 39		12 41 21.8	3.094	2	95 2 2.3	19.71	2	vB; L; iE 45°±; smbMrN ..	5
3228		I. 8	}	12 41 28.1	3.032	6	80 44 53.3	19.71	6	cB; pL; iR; bM; r.....	6*
		III. 6									
3229	{ 1437 = 3425 1438 }	I. 129		12 41 46.4	3.107	2	97 54 3.3	19.71	2	vB; R; vmbMrN; r	3
3230	{ 1438 = 3426 }	III. 524		12 41 49.6	3.119	2	100 38 16.3	19.71	2	F; L; mE40°; vlbM; B*p ...	4
3231		II. 578		12 42 2.1	3.054	1	85 50 43.0	19.70	1	F; S.....	1
3232		III. 514		12 42 5.8	3.109	2	98 21 44.0	19.70	2	eF; cS; pmE	2
3233	1439	II. 662		12 42 8.9	2.842	2	47 18 39.0	19.70	2	cF; S; R; gbM	3
3234	3427			12 42 15.5	3.288	1	130 31 49.0	19.70	1	vF; vS; R; psbM	1
3235		III. 815		12 42 19.7	2.753	1	38 2 43.0	19.70	1	S; stellar	1
3236		III. 722		12 42 22.8	3.118	1	100 20 43.7	19.69	2	eF; S	1
3237	3428		Δ. 511	12 42 26.6	3.289	1	130 36 13.7	19.69	1	pB; cS; R; gbM.....	1
3238	1440	III. 610		12 42 29.3	3.092	1	94 26 6.7	19.69	1	cF; pL; iE	2
3239	1451, a		R. nova	12 42 35.9	2.95	::	63 45 13.4	19.69	::	E	0
3240	1441	II. 95		12 42 39.1	3.000	2	74 4 17.7	19.69	2	cB; pL; vmE 28°5; sbMN ..	4+
3241	1442			12 42 39.4	2.948	1	63 45 43.7	19.69	1	vF; pL	1
3242	1443	II. 412		12 42 40.1	2.888	1	53 54 17.7	19.69	1	F; S; E; glbM; er	4
3243	1444	I. 140		12 42 51.5	3.045	4	83 55 22.7	19.69	4	pB; L; vIE; glbM	6
3244	1445	III. 536		12 43 1.6	3.130	2	102 33 51.7	19.69	2	F; pS; R; gbM	3
3245	1446			12 43 19.9	3.093	1	94 30 56.4	19.68	1	eF; vS; bet 2 st	1
3246	1448	III. 424		12 43 22.9	2.900	1	56 4 35.4	19.68	1	vF; stellar	2
3247	1475, a		R. nova	12 43 26	2.92	::	60 23 ±	19.68	::	E; bMN	0
3248	1447	III. 611		12 43 26.4	3.088	1	93 23 9.4	19.68	1	eF; S; bM	2
3249	1451	I. 84		12 43 32.0	2.945	1	63 44 13.4	19.68	1	vB; vL; E; vg; vsmbMeBN ..	2+
3250	1449	III. 280		12 43 34.1	3.135	1	103 34 27.1	19.67	1	F; vS; R; stellar; np of 2 ...	2
3251	1450	II. 298		12 43 37.5	3.135	1	103 34 57.1	19.67	1	F; pL; R; lbM; sf of 2	2
3252	3430			12 43 38.8	3.293	1::	130 19 50.1	19.67	1::	neb; 1st of 3	1
3253	3431			12 43 38.8	3.293	1::	130 19 50.1	19.67	1::	2nd of 3	1
3254	1452	I. 41		12 43 45.4	+3.098	1	95 38 6.1	+19.67	1	vF; pL; E	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
3255	h.	II. 814		h m s 12 43 53.8	s +2.720	1	36 20 41.1	+19.67	1	F; S; vsmbM	1
3256	1453	II. 73		12 44 3.4	3.019	4	78 19 45.1	19.67	5	cF; pL; { ^R _{mE90°} } r; *12p.	6*
3257	1454			12 44 5.1	3.047	1	84 23 15.1	19.67	1	vF; vS; R	1
3258	1456		M. 94	12 44 17.2	2.838	5	48 6 31.8	19.66	5	vB; L; iR; vsvmbMBN; r...	10†
3259	1457	III. 496		12 44 18.7	2.890	1::	55 5 14.8	19.66	1	eF; vS; pmE	2
3260	1455	III. 515		12 44 20.7	3.108	1	97 39 9.8	19.66	1	F; pL; IE; pglbM	2
3261	1458	III. 721		12 44 30.1	2.776	1	41 33 55.8	19.66	1	vF; S; R; psbM	2
3262	3432	I. 133		12 44 30.5	3.118	1	99 41 33.8	19.66	1	cB; vS; vbmN=*9; *10 sf	2
3263	3429			12 44 31.3	3.300	1	130 37 52.8	19.66	1	F; R; gbM	1
3264	3433			12 44 37.2	3.297	1	130 18 42.8	19.66	1	F; L; E; gbM; 3rd of 3.....	1
3265	1460			12 44 52.3	3.012	1	77 9 54.5	19.65	1	pB; mE; r	1
3266		II. 344		12 44 54.5	2.940	1	63 28 40.5	19.65	1	F; pL; IE	1
3267	1459	III. 537		12 44 55.1	3.132	1	102 38 47.5	19.65	1	F; vS; iR; gbM	2
3268		III. 907		12 44 55.5	2.244	1	17 36 39.8	19.66	1	vF; cL; E 135°±	1
3269	1451, b		R. nova	12 45 0.0	2.95	::	63 23 0.4	19.65	::	E 0°	0
3270	1463	IV. 78		12 45 0.2	2.175	1	16 21 38.5	19.65	1	pB; L; R; vg, vsbM	2
3271	3434			12 45 6.7	3.313	1	131 54 27.5	19.65	1	B; pS; R; vg, vsmbM	1
3272		III. 82		12 45 10.2	3.005	1	75 44 40.5	19.65	1	vF; S; E; r	1
3273	1461	I. 16		12 45 11.6	3.075	1	90 26 26.5	19.65	1	cB; L; vIE; vglbM.....	5
3274	1462	{ I. 25 =		12 45 15.0	3.015	5	77 55 36.5	19.65	5	B; pL; R; psbM; p of 2.....	7
3275	3435		Δ. 301	12 45 22.2	3.533	2	149 35 20.2	19.64	3	Cl; vL; st vB (κ Crucis).....	3†
3276	1464	III. 281		12 45 35.3	3.143	1::	104 38 49.2	19.64	1::	vF; pS; r.....	2
3277	1465	III. 70		12 45 46.1	2.992	3	73 23 18.2	19.64	3	vF; pL; E?	4
3278	1466	II. 75		12 45 53.4	3.015	4	78 0 38.2	19.64	4	pB; vmE 34°0; 3B sts; f of 2	6†
3279		III. 489		12 45 55.7	3.152	1	106 13 39.0	19.63	1	vF; S; lbM	1
3280	1467	III. 544		12 46 6.4	3.048	1	84 46 58.9	19.63	1	F; cS; R; gbM	2
3281	3436			12 46 11.3	3.295	2	128 58 3.9	19.63	2	B; pS; IE; mbM.....	3
3282		III. 525		12 46 12.7	3.115	1	98 45 39.9	19.63	1	vF; vS	1
3283	1468	II. 535		12 46 13.2	3.063	1	87 58 25.9	19.63	1	F; pL; mE; *9 p 90°	2
3284		III. 516		12 46 22.4	3.111	1	97 54 38.9	19.63	1	vF; S	1
3285	1469	II. 24		12 46 22.4	3.058	2	87 3 50.9	19.63	2	pF; pS; R; mbM	5
3286	1471	III. 618		12 46 26.5	2.862	1	52 25 5.9	19.63	1	eF; cS; R; bM	2
3287	1470	II. 186		12 46 29.7	3.101	1	95 51 34.9	19.63	1	F; cL; R; vglbM; r	3
3288	3437	II. 559		12 46 46.5	3.114	1	98 26 38.6	19.62	1	F; S; R; vlbM; p of D neb...	2
3289		III. 517		12 46 47.5	3.112	1	98 1 38.6	19.62	1	vF; S	1
3290	3438			12 46 47.5	3.114	1	98 26 17.6	19.62	1	vF; S; R; vlbM; f of D neb	1
3291	1472	III. 106		12 46 48.1	3.020	2	79 31 37.0	19.60	2	vF; pL; R; r	3
3292		I. 134		12 47 4.9	3.120	1	99 46 38.3	19.61	1	cB; vL; mE	1
3293		I. 135		12 47 15.1	3.131	2	101 49 8.3	19.61	2	pF; pS; R; mbM; p of D neb	2
3294		I. 136		12 47 16.1	3.131	2	101 50 8.3	19.61	2	pF; pS; R; mbM; f of D neb	2
3295		III. 526		12 47 16.9	3.121	1	99 51 38.3	19.61	1	eF; cS	1
3296	3439			12 47 17.2	3.384	1	137 58 54.3	19.61	1	vF; S; R; glbM	1
3297		II. 187		12 47 21.4	3.102	2	96 6 8.3	19.61	2	pB; pS; mbM; r.....	2
3298	1473	II. 345		12 47 29.9	2.925	3	62 9 58.3	19.61	2	F; R; *9 att 1'n	4
3299		II. 560		12 47 31.8	3.120	1	99 28 38.3	19.61	1	pF; pS; iR	1
3300	1475	I. 93		12 47 54.0	2.911	2	60 17 53.0	19.60	2	pB; pS; IE; *8nf 1'	4
3301		II. 538		12 47 55.2	3.132	2	101 52 8.0	19.60	2	vF; S; 2 or 3 st near	2
3302	1474	II. 21		12 47 59.1	3.029	3	81 10 55.0	19.60	3	pF; pL; R; bM; r	7
3303	1477	II. 382		12 48 6.7	2.921	2	61 49 20.0	19.60	2	pF; pS; gbM	3
3304	1476	III. 548		12 48 7.8	3.054	1	86 20 18.0	19.60	1	cF; S; vS* att	2
3305	1478	I. 211		12 48 9.1	2.765	1	42 43 7.0	19.60	1	pB; cS; R; psbM; *14 p ...	5
3306	1479	III. 816		12 48 26.6	2.683	1	36 8 27.7	19.59	1	eF; S; IE.....	2
3307		IV. 40		12 48 33.9	3.136	1	102 17 37.7	19.59	1	S; att to pB*	1
3308	3440			12 48 39.2	3.230	1	118 45 0.7	19.59	1	F; cS; R; gvlbM	1
3309	} 1509, a		R. 2 novæ	12 48 45	3.05	::	86 42 ±	19.58	::	{ F; D neb; E at right angles to each other.	0
3310											
3311	1480	I. 141		12 48 53.4	+3.047	1::	84 56 47.4	19.58	1::	{ H. vB } ; cL; E 135°± ...	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s		131° 2' 20.4	+19.58	1	eF; cS; R; gbM; p of 2.....	1
3312	3441			12 49 6.0	+3.326	1	131 3 50.4	19.58	1	eF; S; R; gbM; f of 2	3
3313	3442			12 49 6.1	3.326	1	61 29 48.4	19.58	1	vF; pL.....	2
3314	1481	II. 383		12 49 9.7	2.915	1	30 54 17.4	19.58	2	B; pS; vE; vgbM	2
3315	1483	I. 243		12 49 10.9	2.590	2	96 3 28.1	19.57	1	F; S; R; bM	2
3316	1482	II. 777		12 49 20.0	3.104	1	154 11 48.1	19.57	3	Cl; pL; pRi; iF; st 10 ...18	3
3317	3443			12 49 23.7	3.680	3					
3318	1484 = 3445	II. 549		12 49 33.0	3.112	2	97 46 5.1	19.57	2	pB; L; pmE 0°; gbM	3
3319	1485	II. 384		12 49 48.9	2.917	1	62 3 50.8	19.56	1	F; cL	2*
3320		II. 563		12 49 51.1	3.140	1	102 54 35.8	19.56	1	pB; iF; bM.....	1
3321	1486		M. 64	12 49 51.8	2.951	3	67 33 15.8	19.56	3	!; vB; vL; vmE 120°±; bMSBN=? ?.	10†
3322	3447			12 50 5.6	3.220	1	116 32 13.8	19.56	1	F; S; R; gbM	1
3323	3446			12 50 6.4	3.321	1	129 59 36.8	19.56	1	pF; vS; R; sbM*17; *10, 70°3	1
3324	1487	II. 346		12 50 6.5	2.917	1	62 14 32.8	19.56	1	vF; pL; iF	3
3325	3444		Δ. 164	12 50 7.9	3.899	2	160 6 53.5	19.55	2	⊕; B; L; R; g, vsbM; st 12	2
3326	1488	III. 817		12 50 11.1	2.680	1	36 56 44.8	19.56	1	vF; S; iR; bM	2
3327	3448			12 50 15.0	3.375	2	135 30 1.5	19.55	3	F; pL; mE; vgbM	3
3328	1489			12 50 23.4	2.725	1	40 26 12.5	19.55	1	Neb; ?	1
3329	1490			12 50 42.3	3.138	1	102 17 56.2	19.54	1	vF; 3Sst sp	1
3330	1491	II. 536		12 50 42.5	3.060	1	87 40 23.5	19.55	1	pF; pL; pmE; vgbM; *nf 30°	2
3331	1493	II. 387		12 50 45.5	2.906	2	60 46 1.2	19.54	2	pF; pL; R; vS* att	3
3332	1492	III. 613		12 50 47.5	3.087	1	92 51 26.2	19.54	1	cF; E; er; *sf 30"	2
3333	1494	II. 386		12 50 55.4	2.912	1::	61 49 52.2	19.54	1	F; pL; R.....	2
3334	1495			12 51 9.4	2.837	1	51 52 3.2	19.54	1	eF	1
3335	3449		Δ. 311	12 51 45.5	3.571	2	148 50 34.2	19.54	2	Cl; L; pRi; iR; st 10... ..	2
3336	1496	II. 385		12 51 48.6	2.908	2	61 38 24.6	19.52	2	F; S; R; psbM	3
3337	1497	I. 68		12 51 59.8	3.151	1	104 17 25.6	19.52	2	B; R; psmbM; *13 np	3*
3338		II. 299		12 52 1.7	3.152	1	104 31 34.6	19.52	1	pB; pL; mbM.....	1*
3339		III. 908		12 52 1.7	2.172	1	19 1 32.6	19.52	1	eF; vS; iR; vlbM	1
3340	1499	IV. 30		12 52 22.9	2.852	2	54 23 0.3	19.51	2	vF; pL; vmE 30°±; bet 2 st	5†
3341		II. 644		12 52 24.2	2.832	1	51 56 32.3	19.51	1	pB; S; R; mbM	1
3342	1498	I. 162		12 52 28.6	2.990	6	75 4 23.3	19.51	6	B; pL; mE 90°; sbMN; S* inv.	7
3343	1500			12 52 41.1	2.903	1::	61 16 45.3	19.51	1::	1st of 5; *7 n	1
3344	1501	II. 388		12 52 49.6	2.903	2	61 17 56.0	19.50	3	cF; S; R; *7 n; 2nd of 5 ...	4
3345		III. 758		12 52 58.9	3.102	1	95 20 33.0	19.50	1	vF; vS; p of 2	1
3346		III. 759		12 52 58.9	3.102	1	95 20 33.0	19.50	1	vF; vS; F of 2	1
3347	1502	II. 389		12 53 0.1	2.902	1::	61 16 45.0	19.50	1::	3rd of 5	2
3348	1503	III. 83		12 53 9.2	2.999	2	76 46 17.0	19.50	2	cF; pL; R; vglbM; r.....	3
3349	1504			12 53 17.4	3.106	1	96 5 50.7	19.49	1	vF; S; E	1
3350	1505	II. 778		12 53 20.0	3.102	1	95 19 0.7	19.49	1	pF; cS; E; psbM; *np.....	2
3351	1507	II. 391		12 53 24.0	2.901	2	61 15 23.7	19.49	2	pB; pmE; bM; *7 n; 4th of 5	3
3352	1506	III. 614		12 53 24.1	3.094	1	93 50 3.7	19.49	1	cF; S; iR; bM	2
3353	1508	II. 390		12 53 28.3	2.908	1	62 21 3.7	19.49	1	vF	2
3354	1510	III. 363		12 53 29.7	2.900	1	61 17 14.7	19.49	2	pF; S; R; *7 n; 5th of 5 ...	3
3355		II. 300		12 53 33.9	3.147	2	103 11 32.7	19.49	2		2
3356	1509	I. 143		12 53 33.9	3.055	2	86 45 0.7	19.49	3	cB; cE; *10 att 135°±	4*†
3357	1512			12 53 35.2	2.724	1	42 1 50.7	19.49	1	pF; S; R; gbM	1
3358	1511	I. 69		12 53 37.9	3.150	1	103 45 56.7	19.49	1	pB; pL; iR; st nr	2*
3359	3450			12 53 44.3	3.257	1	120 12 2.4	19.48	1	vF; cS; R; * att; p of 2.....	1
3360	3451			12 53 51.3	3.257	1	120 10 2.4	19.48	1	vF; vS; R; slbM; f of 2.....	1
3361		II. 517		12 53 51.5	3.069	2	89 18 31.4	19.48	2	pB; pS; R; bM	2
3362	3452			12 54 9.0	3.361	1	132 0 55.1	19.47	1	cF; 3 or 4 st 11, 12, f	1
3363		V. 3		12 54 9.5	3.060	1	87 35 32.4	19.48	1	eF; vL; rr	1*
3364		II. 392		12 54 10.0	2.899	1::	61 23 44.4	19.48	1::	1st of 3	1
3365	1514	II. 645		12 54 10.5	2.823	2	51 54 16.4	19.48	2	pB; cS; R; smbM; *17 np..	3
3366	1513	IV. 47		12 54 12.7	3.094	1	93 47 23.4	19.48	1	pB; S; R; bM; stellar?	3
3367	1515			12 54 26.8	2.719	1	42 1 52.1	19.47	1	eF; S; E; bM	1
3368	1516	II. 393		12 54 39.4	+2.897	1	61 21 47.1	+19.47	2	F; pL; 2nd of 3	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
3369	1518	II. 394		12 54 42.9	+2.897	1	61° 24' 48".1	+19.47	1	vF; 3rd of 3.....	2
3370	1517			12 54 50.7	3.154	2	104 13 10.8	19.46	2	eF; L; vIE 45°±	4
3371	1519	II. 779		12 54 51.3	3.112	1	96 57 43.8	19.46	1	eF; S	2
3372		III. 364		12 55 1.9	2.895	1	61 12 29.8	19.46	1	vF	1
3373	3453	II. 190		12 55 43.0	3.115	1	97 19 31.2	19.44	1	F; pS; vIE; glbM	4
3374		III. 760		12 55 51.7	3.115	1	97 22 30.2	19.44	1	eF; vS; R	1
3375	3454			12 56 14.8	3.358	1	130 39 45.9	19.43	1	vF; R; Δ2st 8, 9, f	1
3376		III. 818		12 56 20.2	2.663	1	38 47 27.9	19.43	1	eF; S; R; vglbM	1
3377		II. 191		12 56 27.9	3.136	1	100 44 59.9	19.43	1	pB; pL; iR	2
3378	3456			12 56 38.5	3.263	2	119 46 26.6	19.42	2	pB; S; R; bM; *f 6s	1
3379	3455			12 56 42.6	3.425	1	136 28 2.6	19.42	1	eeF; S; R; p of 2	1
3380	1521			12 56 48.3	2.646	1	37 55 20.6	19.42	1	eF; R; psbM	1
3381	3458	II. 561		12 56 56.5	3.129	1	99 35 40.6	19.42	1	pB; L; R; gmbM	2
3382	3457			12 56 58.2	3.426	1	136 29 22.3	19.41	1	F; S; R; f of 2	1
3383	1520	I. 40		12 56 59.1	3.101	2	94 48 23.6	19.42	2	pF; L; E; gbMBN; r	3
3384		III. 761		12 56 59.5	3.113	1	96 55 28.6	19.42	1	vF; S	1
3385	1522	II. 395		12 57 6.0	2.887	1	61 3 42.6	19.42	1	F; S; R; bM; *9 nf 1'	2
3386	3459		Δ. 411	12 57 14.1	3.455	1	138 32 6.3	19.41	1	B; vL; vME 38°-7	1
3387	3460			12 57 32.1	3.388	2	132 50 51.0	19.40	2	B; pS; R; gpmbM; p of 2... ..	2
3388	3461			12 57 34.1	3.306	1	124 35 6.0	19.40	1	F; pL; R; vglbM	1
3389	3462			12 57 40.2	3.387	2	132 45 46.0	19.40	2	eF; S; R; psbM; f of 2.....	2
3390	1523	II. 188		12 57 43.2	3.107	(2)	95 45 18.0	19.40	2	F; pL; iE; r	3
3391	1524	II. 396		12 58 15.4	2.877	4	60 7 28.7	19.39	5	F; S; R; psbM*11	6
3392	3463			12 58 18.4	3.329	1	126 48 31.4	19.38	2	vF; pS; am 3Sst.....	2
3393	1527			12 58 23.9	1.656	2	13 50 36.4	19.38	2	vF; S; R; vglbM.....	2*
3394	3464			12 58 26.1	3.263	1	119 0 14.4	19.38	1	F; cS; R; gbM	1
3395	1525	II. 413		12 58 26.5	2.825	2	54 4 20.7	19.39	2	pB; cS; R; smbM	4
3396	1526	II. 397		12 58 27.1	2.888	1	61 40 48.4	19.38	1	F; S; R	2
3397	3465	I. 130		12 58 31.6	3.116	1	97 15 49.4	19.38	1	vB; pS; E 0°±; bMBN.....	3
3399	1528			12 59 2.4	2.840	1	56 4 6.1	19.37	2	eF; S; R	2
3400	1529	II. 398		12 59 4.5	2.885	1	61 30 58.1	19.37	*1	F; S; iF	2
3401		III. 303		12 59 9.8	2.874	1	60 10 26.1	19.37	1	eF; vS.....	1
3402	1530	II. 663		12 59 27.3	2.754	2	47 31 10.8	19.36	2	F; vS; R; stellar; vS*s.....	3
3403	1532	III. 779		12 59 28.4	2.537	1	32 55 57.8	19.36	1	eF; S; iE.....	3
3404	3466			12 59 33.6	3.255	2	117 28 37.8	19.36	2	vF; vL; cE; vgbM	2
3405	1531	III. 304		12 59 35.3	2.873	2	60 12 10.8	19.36	3	vF; vS; vIE; vglbM; *sp... ..	4
3406	1533	III. 783		12 59 35.4	2.588	1	35 40 40.8	19.36	1	vF; S; E; * att	2
3407	3467			12 59 37.3	3.221	1	112 55 45.8	19.36	1	F; pL; R; glbM	1
3408		III. 765		12 59 57.7	3.224	1	113 15 24.5	19.35	1	vF; pL; iF	1
3409		III. 937		13 0 17.3	1.670	1	13 57 22.5	19.35	1	vF; S; iR; bM	1
3410		III. 781		13 0 18.4	2.582	1	35 38 40.2	19.34	1	vF; S	1
3411		III. 782		13 0 32.3	2.580	1	35 36 40.2	19.34	1	vF; S	1
3412	1534			13 0 36.8	3.099	1	94 16 0.9	19.33	1	vF; vS; R; psbM	1
3413	3468			13 0 40.4	3.480	1	138 45 23.9	19.33	1	B; pL; R; gmbM	1
3414		III. 780		13 0 41.9	2.540	1	33 34 22.2	19.34	1	eF; S	1
3415	1535			13 0 57.8	2.948	2	70 50 8.9	19.33	2	F; vS; R; sbM; stellar	2*
3416		III. 346		13 1 8.2	2.902	1	64 29 21.6	19.32	1	eF; pL; iE	1
3417	3469			13 1 29.8	3.264	1	117 53 57.3	19.31	1	eF; cS; R	1
3418	1537	II. 189		13 1 32.2	3.111	1	96 1 35.3	19.31	1	B; pL; R; *9 sf	4
3419		III. 365		13 1 32.5	2.873	1	60 56 21.3	19.31	1	vF	1
3420	1536	II. 301		13 1 33.1	3.168	2	104 45 42.3	19.31	2	B; pL; R; psmbM	3*
3421		II. 185		13 1 40.9	3.102	1	94 36 21.3	19.31	1	F; S; iF; pB*nr.....	1*
3422	1539	III. 654		13 1 50.4	2.743	1	47 34 57.3	19.31	1	vF; vS; R; lbM	2
3423	1538	III. 401		13 1 51.4	2.811	2	54 3 8.3	19.31	2	vF; S; R; stellar	3
3424	1542	II. 815		13 1 54.5	2.599	1	37 18 52.3	19.31	1	vF; vS; stellar	2
3425	3470			13 1 54.7	3.404	1	132 21 15.0	19.30	1	vF; S; E; r.....	1
3426			Auw. N. 31	13 2 1.0	3.099	...	94 38 51.0	19.30	...	A neb (Markree Obs. Apr. 9, 1852).	0*
3427	1541			13 2 6.3	2.993	1	77 37 1.0	19.30	1	vF; S; iE; 2Sst s	1
3428		III. 766		13 2 11.9	3.225	1	112 38 21.7	19.29	1	vF; vS.....	1
3429	3471			13 2 15.9	+3.219	1	111 48 8.7	+19.29	1	pF; cS; R; slbM; am st.....	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	"			
3430	$\left\{ \begin{array}{l} 1540 \\ = \\ 3472 \end{array} \right\}$	I. 42		13 2 23.2	+3.118	2	97 5 2.7	+19.29	2	pB; pL; R; vgpmbM; * 8 np	6
3431		III. 819		13 2 29.4	2.616	1	38 34 19.7	19.29	1	vF	1
3432	1543	II. 537		13 2 32.9	3.057	1	87 35 35.7	19.29	1	cF; pL; R; lbM; er	2
3433	1544	III. 366		13 3 5.6	2.863	1	60 20 39.4	19.28	1	cF; pS; lE	2
3434	1545			13 3 37.6	2.558	1	35 44 28.1	19.27	1	pF; S; iR; gbM	1
3435		III. 655		13 4 15.2	2.704	1	47 27 18.5	19.25	1	vF; pS; lbM	1
3436	1546	III. 305		13 4 20.5	2.852	1	59 36 40.5	19.25	2	vF; vS; vLE	3
3437	1547	I. 96		13 4 28.5	2.781	2	52 11 9.2	19.24	2	vB; vL; vmE 25°; vsbMN...	4
3438		III. 848		13 4 34.7	2.340	1	27 7 16.5	19.25	1	vF; vS	1
3439		D'Arrest, 93		13 4 37	3.05	[1]	63 51 42	19.24	[1]	pF; pL; R	0
3440	1550	III. 820		13 4 45.5	2.610	1	39 9 53.2	19.24	1	vF; R; bet 2 vS st	2
3441	1549	I. 85		13 4 48.8	2.907	2	66 20 21.9	19.23	2	pF; cL; E 17° 0; biN; * 9 f	3
3442	1548			13 4 50.5	3.175	1	105 3 9.9	19.23	1	vF; R; bM; * 10 np 5'	1
3443	3473			13 4 51.7	3.419	3	132 21 4.9	19.23	3	pB; cS; R; am 4 st	3
3444	1551	II. 414		13 5 2.2	2.787	1	52 58 43.9	19.23	1	pF; S; E; psbM	2
3445	1552	II. 637		13 5 7.5	3.097	(1)	93 36 2.6	19.22	1	F; cL; iR; lbM	2
3446		II. 356		13 5 7.9	2.897	1	65 12 15.6	19.22	1	pB; S	1
3447	1553	III. 669		13 5 28.3	3.183	1	106 0 52.6	19.22	1	vF; R; bM	2
3448	1554	II. 746		13 5 30.7	3.203	1	108 46 4.3	19.21	1	cB; S; R; mbMpBN	2
3449	1555	III. 545		13 5 36.5	3.036	1	84 31 7.3	19.21	1	cF; vS; R; er	2
3450	1556	II. 129		13 5 42.5	2.982	3	76 39 28.3	19.21	3	cF; cL; vLE; lbM	4
3451	1557			13 5 46.2	2.663	1	43 4 1.3	19.21	1	pF; cS; R; * 12 nf 90"	1
3452	1559	II. 664		13 5 55.8	2.692	2	45 12 45.3	19.21	2	pF; L; mE 20°; vlbM	4
3453	1558		M. 53	13 6 2.0	2.941	5	71 5 3.0	19.20	5	!; ⊕; B; vC; ir; vvmB; st 12...	12
3454	1560	III. 649		13 6 10.1	2.826	3	57 26 40.0	19.20	2	vF; S; lE; * 13 n	4
3455	3474			13 6 13.8	3.425	3	132 12 57.0	19.20	3	pB; pL; R; gbM; * 7 nf ...	3
3456	1561			13 6 16.8	3.027	2	83 11 34.0	19.20	2	vF; S; R; pgbM	2
3457	1562			13 6 27.1	2.646	1	42 10 39.0	19.20	1	F; vS; R; gbM	1
3458	1563	III. 367		13 6 45.9	2.861	2	61 27 30.7	19.19	2	vF; pL; iR	3
3459	1564	I. 97		13 6 59.7	2.775	2	52 39 40.4	19.18	2	vB; pl; E 166° 8; smbMvBN; * np.	3
3460		III. 909		13 7 19.4	1.918	1	18 36 12.4	19.18	1	vF; vS; R	1
3461	1565	II. 510		13 7 33.0	3.185	2	105 51 1.8	19.16	2	cF; pS; vLE; bM	3
3462		II. 816		13 7 49.8	2.568	1	37 58 10.8	19.16	1	F; S; ir; vgmbM	1
3463	3477			13 7 55.8	3.243	1	113 14 29.5	19.15	1	F; L; R; vglbM; * 9 p ...	1
3464	3476			13 7 56.4	3.744	1	149 19 9.5	19.15	1	Cl; P; E; sc st 11... ..	1
3465	1566	II. 511		13 7 56.9	3.184	1	105 38 49.5	19.15	1	pB; pL; R; bM	3
3466	3475			13 7 59.4	3.845	1	152 40 25.2	19.14	1	Cl; vL; vRi; st 11... ..	1
3467	3478			13 8 26.1	3.283	1	117 40 52.2	19.14	1	pF; R; sp of 2	1
3468		III. 670		13 8 26.7	3.185	1	105 43 13.2	19.14	1	vF	1
3469		II. 512		13 8 31.7	3.185	2	105 37 43.2	19.14	2	cF; S	2
3470	3479			13 8 46.5	3.283	1::	117 35 33.9	19.13	1::	Neb; nf of 2	1
3471	1567			13 9 0.0	2.837	1	59 33 42.9	19.13	1	vF	1
3472	1569	VI. 7		13 9 30.5	2.938	1	71 35 10.3	19.11	2	Cl; vF; pL; iR; vgbM; st 15...	3
3473	1568	II. 513		13 9 31.1	3.188	1	105 53 46.3	19.11	1	F; pS; iR	3
3474	1570		M. 63	13 9 31.9	2.699	1	47 13 45.3	19.11	1	vB; L; pmE 120° ±; vsmbMBN.	3
3475	1571	III. 306		13 9 38.0	2.823	1	58 18 9.3	19.11	1	cF; cS; R; p of 2	2
3476	1572	III. 307		13 9 52.3	2.821	1	58 13 16.0	19.10	1	cF; cS; R; f of 2	2
3477	3480	I. 138		13 10 27.4	3.274	1	116 6 47.7	19.09	1	vB; S; R; vsmbM; * 10 f...	4
3478	3482			13 10 33.7	3.358	1::	124 41 30.4	19.08	1	eF; vS; E; r	1
3479	3481			13 10 33.9	3.357	1	124 35 15.4	19.08	1	eF; vS; R; * nr	1
3480	3483			13 10 46.3	3.519	1	137 10 22.1	19.07	1	B; S; R; psbM	1
3481	1573	III. 308		13 10 57.3	2.817	1	58 10 2.4	19.08	1	vF; cS	3
3482		III. 312		13 11 14.2	3.227	1	110 22 7.8	19.06	1	F; L; iR; bM	1
3483	1574	III. 282		13 11 54.7	3.178	1	104 6 53.5	19.05	1	vF; pL; pmE 135° ±	2*
3484	1575	III. 309		13 11 55.9	+2.810	1	57 47 16.5	+19.05	1	eF; vS	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	"			
3485	$\left\{ \begin{array}{l} 1576 \\ = \\ 3489 \end{array} \right\}$	III. 117		13 12 6.9	+3.162	2	102 0 4.2	+19.04	2	vF; cS; R; 1st of 3	5
3486	$\left\{ \begin{array}{l} 1577 \\ = \\ 3490 \end{array} \right\}$	II. 193		13 12 9.0	3.161	2	101 55 11.2	19.04	2	pB; S; vIE; sbM; 2nd of 3...	5
3487	3484	II. 566		13 12 9.6	3.285	1	116 50 8.2	19.04	1	pB; pS; cE; psbM; *7.8 f...	3
3488	$\left\{ \begin{array}{l} 1578 \\ = \\ 3491 \end{array} \right\}$	III. 118		13 12 14.9	3.162	2	101 57 46.0	19.04	2	cF; pS; vIE; 3rd of 3.....	5
3489	1578, a		R. nova	13 12 ±	3.162	...	101 57 ±	19.04	...	No description, one of a group of four.	0
3490	3485			13 12 26.3	3.285	1	116 40 17.9	19.03	1	vF; S; R; 1st of 4	1
3491	1579	II. 313		13 12 42.4	3.236	1	111 4 37.6	19.02	1	cB; cS; vIE 90°±; bf	2
3492		II. 780		13 12 44.0	3.258	1	113 40 5.6	19.02	1	F; L; R; vglbM.....	1
3493	3486			13 12 55.4	3.469	1	132 59 25.6	19.02	1	eF; vS; R; 2nd of 4	1
3494		III. 724		13 12 55.9	3.226	1	109 52 4.6	19.02	1	cF; vS; iF	1
3495	1580	II. 327		13 13 3.1	2.818	2	59 1 33.6	19.02	2	pF; pL; gbM	3
3496	3487			13 13 4.2	3.470	2	132 58 29.3	19.01	2	pB; pL; R; 3rd of 4	2
3497	3488			13 13 8.5	3.470	2	132 59 51.3	19.01	2	cF; S; vIE; 4th of 4	2
3498	1583	III. 633		13 13 20.1	2.702	1	48 51 23.3	19.01	1	vF; S; R; lbM	2
3499	1581	III. 539		13 13 22.1	3.174	1	103 20 52.0	19.00	1	cF; vS; R; gbM.....	2
3500	1582			13 13 24.5	3.085	1	91 34 7.0	19.00	1	vF; iR; * 11 sp	1
3501	1584	III. 650		13 13 39.1	2.787	1	56 11 1.0	19.00	1	vF; cS; R; bM; sp of 2.....	3
3502	1585			13 13 46.8	2.786	1	56 7 15.0	19.00	2	vF; S; bet 2 st; nf of 2	2
3503	3493	II. 567		13 14 3.7	3.289	2	116 41 36.7	18.99	2	cB; pS; iE; psbM *	4
3504	3492			13 14 4.5	3.385	1	125 54 9.7	18.99	1	vB; pS; R; svmbM	1
3505		II. 665		13 14 9.9	2.660	1	46 10 1.7	18.99	1	pB; cS; E	1
3506		II. 22		13 14 34.0	3.002	1	80 46 29.4	18.98	??	vF; vS	1*
3507	1586	III. 619		13 15 6.1	2.717	1	50 42 22.8	18.96	1	vF; S; cE 0°±	2
3508	3494			13 15 12.3	3.342	1::	121 36 35.5	18.95	1	eeF; p of 2	1
3509	1588	III. 808		13 15 14.4	2.367	1	31 37 0.8	18.96	1	cF; S; cE	2
3510	1587	III. 119		13 15 32.9	3.168	2	102 14 3.2	18.94	2	cF; cS; iR; glbM	2
3511	1589	II. 646		13 15 38.3	2.712	1	50 31 29.5	18.95	1	F; L; iR; vglbM	2†
3512		II. 826		13 16 2.4	2.359	1	31 34 28.2	18.94	1	F; S; E	1*
3513	3495			13 16 12.7	3.345	1	121 36 34.6	18.92	1	F; iE; psbM; f of 2	1
3514	1590	III. 368		13 16 19.0	2.841	2	62 17 27.6	18.92	2	pF; pS; pmE; glbM; r	3
3515	1592			13 16 19.0	2.827	1	60 56 38.6	18.92	1	vF; L; Δ 2 st 11 mp.....	1
3516	1591	III. 925		13 16 24.5	3.018	1	82 52 33.6	18.92	1	vF; S; R; gbM	3
3517	3497			13 16 35.7	3.164	1	101 33 30.3	18.91	1	pB; S; iE	1
3518	3496			13 16 37.7	3.939	1	152 40 50.0	18.90	1	Cl; eR; mC; st 12...16	1
3519	3498			13 16 46.4	3.409	2	126 57 18.3	18.91	2	cB; P; R; psmbM; r	2
3520	1594	II. 666		13 16 59.8	2.646	1	46 10 43.3	18.91	1	pF; S; R; gmbM	2
3521	3499			13 17 1.9	3.327	2	119 34 56.0	18.90	2	vF; S; vIE	2
3522	1593			13 17 2.8	2.990	1	79 33 7.0	18.90	1	pF; S; R; gbM	1
3523	3500			13 17 5.0	3.333	1	119 36 57.0	18.90	1	vF; vS	1
3524	1596	II. 328		13 17 15.0	2.790	2	57 42 15.0	18.90	3	pB; pL; R; gmbM; * p.....	5
3525	3501		Δ. 482	13 17 15.1	3.481	4	132 17 10.7	18.89	4	!!; vB; vL; vME 122°5; bifid	4†
3526	1595	II. 653		13 17 15.9	2.955	3	75 17 16.7	18.89	3	pB; vS; R; gmbM; * f.....	4
3527	1597	II. 314		13 17 43.5	3.240	1	110 23 21.4	18.88	1	F; pS; iE; vgbM	2*
3528	3502			13 17 56.3	3.325	1	119 6 14.4	18.88	1	pB; S; E	1
3529	1598	III. 84		13 17 59.6	2.953	1	75 31 20.4	18.88	1	eF; vS; R; psbM	3
3530	3503		Δ. 312?	13 18 19.5	3.811	2	148 16 44.8	18.86	2	Cl; Ri; iC; st 11... ..	2
3531	3504		Δ. 440	13 18 24.0	3.554	2	136 34 49.8	18.86	2	!!; ⊕; ∞ Centauri	2†
3532	3505			13 18 27.3	3.370	2	123 9 1.8	18.86	2	vF; S; R; glbM.....	2
3533	1599	III. 402		13 18 29.8	2.730	2	52 53 5.8	18.86	2	cF; cS; R; vsmbM*; *12sp; sp of 2.	3
3534	1600	III. 403		13 18 39.8	2.729	2	52 50 44.8	18.86	2	F; cS; R; vsmbM*; n of 2	3
3535	1600, a		R. nova	13 18 39.8	2.729	::	52 49 ±	18.86	:::	vF	0
3536		IV. 70		13 18 48.6	1.743	2	18 45 22.8	18.86	2	○?; cB; S; R; g, sbM.....	2
3537	1602	II. 667		13 19 4.9	+2.632	1	46 0 37.5	+18.85	1	pB; vS; vIE; glbM	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. II.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s						
3538		III. 115		13 19 8.3	+ 3.167	2	101° 35' 25.2	+ 18.84	2	vF; vS; stellar	2
3539	1601	II. 25		13 19 11.5	3.050	2	87 9 53.2	18.84	2	pB; pL; vIE; vsmbM * 12...	6
3540	1604	III. 404		13 19 47.8	2.730	2	53 19 49.9	18.83	2	cF; pS; E; bM; sp of 2.....	3
3541	1603			13 19 49.1	2.926	2	72 23 55.6	18.82	2	vF; S; R; * nf	1
3542	3507			13 19 51.7	3.328	2	118 50 27.6	18.82	2	cF; S; R; psbM; * f 2'.....	2
3543	3508			13 20 2.9	3.329	1::	118 54 24.3	18.81	1::	vF; S; R; p of D neb.....	1
3544	3509			13 20 4.1	3.329	1	118 53 52.3	18.81	1	pF; S; f of D neb	1
3545	1605	III. 405		13 20 7.4	2.728	2	53 16 40.6	18.82	2	vF; pL; R; nf of 2.....	3
3546	3506			13 20 8.6	3.977	1	152 41 27.0	18.80	1	Cl; vRi	1
3547	3510			13 20 17.7	3.594	2	138 10 46.0	18.80	2	pB; cS; iE; glbM; r	2
3548	1606	III. 651		13 20 49.5	2.773	3	57 17 26.7	18.79	3	F; pS; vIE; bM	4
3549	1607			13 20 59.9	2.916	1	71 29 31.4	18.78	1	vF; R	1
3550			D'Arrest, 94	13 21 13	3.02	[2]	83 16 42	18.78	[2]	pF. See note	0*
3551	3511			13 21 19.2	3.373	1	122 26 25.1	18.77	1	pF; L; vmE; pgbM; rr	1
3552		III. 821		13 21 34.6	2.438	1	36 28 48.1	18.77	1	cF; stellar	1
3553	1609	III. 784		13 21 38.4	2.371	1	33 47 3.1	18.77	1	cF; S; iR.....	2
3554	1608			13 21 47.4	2.770	3	57 14 35.8	18.76	3	pF; pL; iE; lbM; f of 2.....	3
3555	3512			13 22 3.3	3.898	2	150 12 3.1	18.77	2	Cl; vF; S; vRi; st 15.....	2
3556	1611			13 22 8.6	2.560	1	42 38 21.5	18.75	1	vF; pS; R	1
3557	1610	V. 22		13 22 17.6	3.221	2	107 14 29.2	18.74	2	cF; L; mE 128° 8; pgbM ...	4
3558	1613			13 22 28.5	2.920	2	72 13 18.2	18.74	2	F; pL; R; gbM	2
3559	1614	III. 672		13 22 28.5	2.558	2	42 41 26.2	18.74	2	F; vS; R; stellar.....	5
3560	1612	III. { 45	}	13 22 29.1	2.974	2	78 16 4.2	18.74	2	{ vF; pL } D neb	4
3561		46								{ vF; pL }	
3562	1615	III. 71		13 22 37.1	2.922	1	72 26 57.9	18.73	2	vF; S; R; am 3 st; * 7 nf...	3
3563	1616			13 22 49.1	2.953	2	75 58 15.9	18.73	2	vF; S; R	2
3564	3513			13 22 51.6	3.321	2	117 25 21.6	18.72	2	vF; pL; vIE; * 7 nf 10'.....	2
3565	1617	II. 679		13 22 54.8	3.081	1	90 59 58.6	18.72	1	F; cS; iE; gbM; p of 2.....	3
3566	1618	II. 680		13 22 59.8	3.081	1	90 56 3.6	18.72	1	pF; pL; iR; bM; f of 2.....	3
3567	1619	III. 642		13 23 9.3	2.952	1	75 54 40.6	18.72	1	vF; S; iR	2
3568	1620	III. 652		13 23 21.2	2.774	1	58 8 37.3	18.71	2	vF; vS; R; glbM	3*
3569	3515			13 23 31.3	3.400	2	124 3 45.0	18.70	2	F; pL; vIE; vglbM	2
3570	3514		Δ. 252?	13 23 41.0	4.129	1	155 15 11.7	18.69	1	l; B; pL; cE; bM curved axis; 4 st inv.	1+
3571	1621			13 23 50.0	2.908	2	71 8 20.0	18.70	2	cF; S; R; bM; ** f	2
3572	1622		M. 51	13 23 55.4	2.539	5	42 5 4.0	18.70	4	!!!; nucl & ring (h); spiral (R)	10+
3573	3516			13 23 57.1	3.382	1	122 30 27.7	18.69	1	pB; S; R; g; psbM.....	1
3574	1623	I. 186		13 24 4.4	2.536	4	42 0 50.7	18.69	3	B; pS; R; vgbM; f of 2.....	6
3575	1625	IV. 63		13 24 12.2	2.264	1	30 51 32.7	18.69	1	pB; cL; iR; gmbM; r	2
3576		II. 689		13 24 14.0	2.546	3	42 35 13.7	18.69	2	pB; pL; R; mbM	3
3577	1624	III. 406		13 24 21.5	2.726	2	54 26 31.4	18.68	2	vF; vS; iE	3
5072				13 24 32.0		...	89 18 29.7		...	See No. 5072.	
3578		II. 797		13 24 39.9	2.409	2	36 12 12.4	18.68	2	pF; cS; R; vglbM	1
3579	3517	III. 507		13 24 52.2	3.142	1	98 3 15.8	18.66	1	vF; cS; R; gbM; r	2
3580	3518			13 25 18.7	3.610	1	137 24 41.2	18.64	1	F; pL; R; vgbM	1
3581	1626	III. 643		13 25 21.9	2.945	1	75 22 53.5	18.65	1	F; S; cE; * 11 att np.....	2
3582	1627	III. 9		13 25 26.6	3.003	3	81 57 44.2	18.64	4	F; vS; R; psbM; p of 2.....	6
3583	1628	III. 10		13 25 42.8	3.003	2	81 57 21.2	18.64	2	F; vS; R; stellar; f of 2.....	4
3584	1629	III. 99		13 25 48.2	3.004	1	82 6 15.9	18.63	1	F; S; R; psbMN	4
3585	1630			13 25 54.1	3.076	1	90 18 50.9	18.63	1	pB; S; R; psmbM	1
3586	1631			13 26 21.9	3.003	1	81 59 5.6	18.62	1	eF	1
3587	1632	III. 656		13 26 44.5	2.615	1	47 24 36.3	18.61	1	vF; S; R; lbM	2
3588	1633	III. 926		13 27 7.3	3.009	1	82 47 49.7	18.59	1	vF; S; * 9 nf inv? (? 28" R.A.)	3*
3589	3519			13 27 10.2	3.397	1	122 45 39.4	18.58	1	eF; cS; * S and * p	1
3590	1635	II. 841		13 27 14.9	2.075	1	26 33 53.7	18.59	1	pB; S; vIE	3
3591	1634			13 27 18.2	2.904	1	71 25 25.4	18.58	1	vF; S; R; bM.....	1
3592	1636	II. 842		13 27 21.3	2.072	1	26 30 43.7	18.59	1	pB; pL; R; gbM	3
3593	3520			13 27 47.5	3.583	1?	135 11 16.8	18.56	1::	vF; S; R; * n, nr	1
3594	3521			13 27 59.8	3.399	1	122 44 13.8	18.56	1	vF; S; R; * 10 f.....	1
3595	1637	III. 86		13 28 4.5	2.942	2	75 27 49.8	18.56	2	vF; S; vIE; 1st of 3	4
3596	1638	III. 85		13 28 4.9	+ 2.943	3	75 32 20.8	+ 18.56	3	cF; S; R; bM; 2nd of 3.....	5

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
3597	1639	III. 87		13 28 5.2	+2.943	1	75° 37' 21.8"	+18.56	2	cF; pL; R; glbM; 3rd of 3..	4
3598	1640	III. 407		13 28 6.2	2.713	3	54 34 50.8	18.56	3	F; cS; R; *10 p; p of 2 ...	5
3599		III. 822		13 28 9.6	2.422	1	37 46 37.8	18.56	1	cF; pS; iR; lbM.....	1
3600	1641	III. 928		13 28 15.0	3.054	(1)	87 53 16.5	18.55	1::	vF; S; R.....	2
3601	1642	III. 408		13 28 16.7	2.711	2	54 29 21.8	18.56	2	vF; vS; R; f of 2	4
3602	1643			13 28 31.7	2.941	1	75 35 24.2	18.54	1	F; L; E; vgbM	1
3603	1645	III. 425		13 28 51.6	2.711	1	54 36 47.9	18.53	1	F; S; R; vS *nr	2
3604	3522			13 28 52.3	3.666	1	139 6 59.6	18.52	1	ceF; S; IE	1
3605	1644	III. 100		13 28 58.3	3.008	3	82 41 27.9	18.53	3	vF; pS; vLE; *9 sp	4
3606	3523		{ M. 83 = Δ. 628 }	13 29 9.0	3.360	4	119 9 31.6	18.52	4	{ (H, h) vB; vL; E 55° 1; esbMN (L) 3-branched spiral. }	6+
3607	3524			13 29 20.8	3.539	4	132 8 2.3	18.51	4	F; pL; cE; vglbM	4
3608	1646	III. 101		13 29 27.9	3.000	1	81 54 44.3	18.51	1	vF; pL; R; er.....	4
3609		III. 823		13 29 26.9	2.410	1	37 39 33.6	18.52	1	cF; pL; R; vlbM	1
3610		III. 409		13 29 29.9	2.696	1	53 43 3.3	18.51	1	vF; pL; R; lbM.....	1
3611	1647			13 30 0.9	3.041	1	86 30 48.0	18.50	1	eF; eL	1
3612	1648	III. 620		13 30 4.6	2.654	2	50 55 41.0	18.50	2	cF; pL; E 65°; biN?	3
3613	3525			13 30 17.1	3.591	2	135 9 0.4	18.48	2	vF; S; R; vglbM; *13 att...	2
3614	1649	II. 297		13 30 30.2	3.234	2	107 10 3.1	18.47	2	{ (H, h) cF; vL; vg, psmbMLN (L) 2-branched spiral. }	4+
3615	1650	I. 34		13 30 35.1	2.985	1	80 23 32.1	18.47	1	B; L; E 150°; psbMrN	3
3616	1651	III. 72		13 30 48.6	2.916	3	73 18 27.1	18.47	3	vF; S; R; bM.....	4
3617		II. 817		13 30 49.8	2.409	1	38 1 30.1	18.47	1	pB; S; R; vgbM	1
3618	1652	III. 369		13 30 53.0	2.795	1	61 51 45.1	18.47	1	vF; S; vLE	2
3619	1653	III. 505		13 31 11.6	3.025	(1)	84 46 14.8	18.46	1	vF; S; R; bM.....	4
3620	3526	II. 638	Δ. 623	13 31 59.9	3.390	1	120 55 34.6	18.42	1	B; pL; E 45° ±; psmbM ...	2
3621	3527			13 32 12.4	3.174	1	100 46 55.6	18.42	1	pB; L; pmE; glbM	1
3622		III. 803		13 32 21.9	2.236	2	32 10 56.6	18.42	2	vF; vS	2
3623	1656	III. 673		13 32 32.9	2.465	1	40 59 35.6	18.42	1	eF; vS; R; gbM.....	2
3624	1654	II. 895		13 32 45.5	3.059	1	88 26 55.0	18.40	1	vF; S; R; bM; p of D neb...	2
3625	1655	II. 896		13 32 49.9	3.058	1	88 27 15.0	18.40	1	F; S; iR; f of D neb	2
3626	1657			13 33 13.6	3.019	2	84 13 5.7	18.39	2	vF; R; am pB st.....	2
3627	1660			13 33 14.4	0.995	1	14 14 14.0	18.40	1	eF; S	1
3628	1658	III. 370		13 33 26.7	2.775	1	60 53 12.4	18.38	1	cF; S; mE 0° ±; *9 sp.....	2
3629	3528			13 33 43.5	3.374	1	119 12 28.8	18.36	1	vF; pL; R; vlbM	1
3630	1659	III. 410		13 33 58.5	2.660	2	52 25 32.8	18.36	2	F; cS; vLE; er.....	3
3631	3529			13 34 27.7	3.665	2	137 27 41.9	18.33	2	B; pL; vLE; vglbM; 3 st nr..	2
3632	1661			13 34 32.8	2.628	4	50 30 0.2	18.34	4	F; S; R; gbM; S * np	4
3633			Auw. N. 32	13 34 43.9	3.196	...	103 9 3.6	18.31	...	A nebula (Markree Obs. 1855)	0
3634	3530			13 35 3.6	4.112	2	152 11 47.3	18.31	2	Cl; P; L; iF; st 12	2
3635	1662			13 35 5.5	3.026	1	85 1 44.6	18.32	1	eF; S; bet 2 st.....	1
3636	1663		M. 3	13 35 40.8	2.769	5	60 55 6.0	18.30	5	!!; ⊕; eB; vL; vsmbM; st 11..	14
3637	1664	I. 98		13 35 56.1	2.670	3	53 37 45.7	18.29	3	cB; pL; R; g, psmbM	4
3638	1664, a		R. nova	13 36 8.6	2.670	::	53 37 45.7	18.29	::	F; S.....	0
3639	1665	II. 798		13 36 24.8	2.248	2	33 37 25.4	18.28	2	F; E 73° 0; D or biN; B *nf	3
3640	3531		Δ. 273	13 36 54.7	4.131	2	152 11 45.2	18.24	2	Cl; B; S; pC; iR; st 10...12	2
3641	3532			13 37 15.3	3.985	1	148 29 24.2	18.24	1	Cl; L; vRi; st 7...16	1
3642	3533		Δ. 388	13 37 37.5	3.757	2	140 40 6.6	18.22	2	⊕; vB; pL; R; rrr; st 15....	2
3643	3534			13 38 48.2	4.237	2	153 59 16.1	18.17	2	Ch; S; C; iR; st 14.....	2
3644	1666	II. 668		13 39 9.3	2.561	1	47 47 33.4	18.18	1	vF; vS; IE 90° ±; sbM	2
3645		I. 170		13 39 17.9	2.556	2	47 33 53.1	18.17	2	cB; pL; E 90° ±; bMN.....	2
3646	3535			13 39 25.8	3.397	1	119 41 12.8	18.16	1	vF; R; vlbM; * p	1
3647	3536			13 39 41.2	3.405	1	120 13 12.2	18.14	1	pF; S; R; 2 st nr	1
3648		V. 6		13 39 55.7	2.899	1	72 59 14.2	18.14	1	eF; vL; r.....	1
3649	1667	III. 785		13 39 57.1	+2.232	1	34 0 24.5	18.15	1	eF; 2 st att or inv	2
3650		III. 946		13 40 20.5	-0.151	1	9 52 6.7	18.09	1	vF; vS; R	1*
3651	1668, a		R. nova	13 40 +	+2.511	::	45 28 +	18.13	::	R; bM; is sp h. 1668	0
3652	1668	I. 180		13 40 31.6	2.511	1	45 27 31.9	18.13	1	cB; L; pmE 142°; gbM.....	2
3653	3538			13 40 39.5	+3.401	1	119 44 21.3	+18.11	1	F; S; R; gbM.....	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	"			
3654	3537			13 40 51.7	+4.064	1	149 14 42.0	+18.10	1	Cl; vL; vRi.....	1
3655	1669	II. 533		13 41 9.9	3.026	2	85 21 35.0	18.10	2	vF; vL; IE; vgbM	4
3656	1670	II. 688		13 41 13.7	2.456	1::	43 9 25.1	18.07	1	eF; L; vmE	3
3657	3539			13 41 22.7	3.405	1	119 47 20.4	18.08	1	F; S; R; gbm	1
3658	1672	III. 681		13 41 40.0	2.605	1	51 0 17.4	18.08	1	pF; cS; IE; F* inv	2
3659	1673	III. 621		13 41 51.2	2.612	2	51 28 9.4	18.08	1	eF; S; R	3
3660	{ 1671 = 3540 }	II. 306		13 41 51.7	3.139	2	96 31 46.8	18.06	2	vF; vS; R; r	3
3661	3541			13 42 11.8	3.782	1	140 30 34.2	18.04	1	O, or vF; eS; D neb	1+
3662	1674	I. 255		13 42 19.9	1.999	1	28 18 58.8	18.06	1	B; pL; mE 57°.4; psbMBEN	2*
5073				13 42 38.3		...	89 14 2.4		...	See No. 5073.	
3663	1675	II. 710		13 42 57.5	2.571	1	49 19 5.9	18.03	1	eF; cS; R; sbM; p of 2	3
3664	1676	III. 422		13 43 37.6	2.671	2	55 41 13.3	18.01	2	vF; R; stellar; 1st of 3	3*
3665	1677	II. 711		13 43 45.0	2.567	2	49 19 42.0	18.00	2	pB; pS; vLE; glbM; f of 2 ...	3
3666	3542		Δ. 282	13 44 7.0	4.155	1	151 9 39.1	17.97	1	Cl; pL; pC; st 11... ..	1
3667	1678			13 44 8.2	3.014	1	84 18 29.4	17.98	1	vF; vL; R; vgbM	1
3668	1679	III. 423		13 44 22.4	2.667	1	55 36 31.1	17.97	2	F; S; R; psbM; 2nd of 3 ...	3*
3669	1682	II. 669		13 44 27.1	2.539	1	47 56 13.1	17.97	1	eF; pL; R; gbm	2
3670	1680			13 44 28.9	2.668	1	55 39 38.1	17.97	1	eF; pL; R; vsmbM*	2
3671	1684	I. 256		13 44 33.4	2.012	1	29 6 16.1	17.97	1	vB; pL; iR; psmbM	2
3672	1689	II. 899		13 44 38.6	0.426	1	12 28 21.7	17.99	1	vF; pS; IE 0°+	2
3673	1681	II. 307		13 44 43.6	3.129	1	95 21 14.5	17.95	1	eF; L; iR; bM	3
3674	1685	II. 712		13 44 51.4	2.570	2	49 44 8.8	17.96	3	eF; S; vLE; sbM	4
3675	1683	II. 685		13 44 52.1	3.089	1	91 30 32.5	17.95	1	F; pS; R; 2 st p	3
3676	3543	III. 923		13 44 56.4	3.388	2	117 46 54.5	17.95	2	pB; S; R; slbM	3
3677	1686	III. 549		13 45 5.2	3.042	1	86 58 26.2	17.94	1	F; vS; R; psbM	2
3678	1687	III. 929		13 45 8.1	3.044	1	87 12 15.2	17.94	1	vF; S; E 0°; rr	3
3679	3544			13 45 39.7	3.737	1	137 48 33.6	17.92	1	vF; vS; R; *8 f	1
3680		III. 665		13 45 44.8	3.078	2	90 25 29.6	17.92	2	eF; vL; R; lbM; r.....	2
3681	1688			13 45 49.7	3.036	1	86 28 58.6	17.92	1	F; iR	1
3682	1690	II. 670		13 46 20.2	2.494	1	46 3 51.0	17.90	1	eF; pL; R; psbM	2
3683	1691	III. 698		13 46 26.0	2.561	1	49 37 37.0	17.90	1	vF; S; iR; B * p.....	3
3684	1694	III. 849		13 46 43.8	2.006	1	29 25 55.7	17.89	1	eF; vS	2
3685	1692	II. 308		13 46 50.5	3.147	1	96 54 11.4	17.88	1	vF; S; R; lbM	3
3686	1693	II. 686		13 47 2.3	3.081	1	90 44 38.1	17.87	1	F; S; R; bM	3
3687	1695	II. 424		13 47 6.9	2.661	1	55 49 32.1	17.87	1	pF; cL; R; lbM	2
3688	1696	II. 713		13 47 25.5	2.544	2	48 56 20.8	17.86	2	eF; pL; bM; B * p; 1st of 4.	5
3689	1697	II. 697		13 47 26.3	2.588	2	51 23 43.8	17.86	2	eF; L; IE 90°; vgbM	3
3690	{ 1697, a }		R. 2 novæ	13 47 ±		...	51 23 ±		...	{ 2 neb in a line with } h. 1697; bM.	0
3691											
3692	1700	II. 415		13 47 32.2	2.617	2	53 10 5.5	17.85	2	eF; S; R; lbM; * nf 90" ...	3
3693	1698	II. 714		13 47 32.9	2.545	3:	49 2 24.5	17.85	1	pB; S; R; 2nd of 4	5
3694	1699	II. 715		13 47 33.2	2.545	3:	49 1 24.8	17.86	1	pF; S; R; 3rd of 4	5
3695	1702	III. 699		13 47 45.1	2.543	2	48 57 33.2	17.84	3	vF; pS; 4th of 4	3
3696	1701	III. 506		13 47 56.1	3.008	5	83 58 28.9	17.83	5	F; pL; vmE 15°; r	7
3697	3546			13 47 58.1	3.422	1	119 38 52.6	17.82	1	pF; S; R; glbM; bet 2 st 10.	1
3698	1703, a		R. nova	13 48 4.7	3.009	::	84 2 32.4	17.81	::	F; S; E (nisi=III. 506).....	0
3699	3545			13 48 30.8	4.752	2	159 43 18.7	17.79	2	Cl; vL; lRi; lC; st 11... ..	2
3700		III. 682		13 48 38.9	2.574	1	50 52 52.3	17.81	1	eF; cS; E	1
3701		II. 671		13 48 58.8	2.520	1	48 3 22.0	17.80	1	pB; pL; E	1
3702	1703	I. 6		13 49 5.3	3.009	4	84 3 32.4	17.78	4	B; pL; R; psbM; *8 nf.....	8
3703	1703, b		R. nova	13 49 5.3	3.009	:	84 17 32.4	17.78	:	vF; L	0
3704	1705	II. 534		13 49 12.0	3.011	(1)	84 18 20.4	17.78	2	eF; L; R; gbm	4
3705	3547			13 49 15.5	3.657	2	133 14 39.1	17.77	2	pB; cS; R; pgbM; am st ...	2
5074				13 49 18.6		...	89 31 9.0		...	See 5074.	
3706	3548			13 49 20.9	3.581	3	129 17 44.1	17.77	4	l; vB; vL; vl; vsmbM* ...	6+
3707	1706	III. 786		13 49 24.0	2.192	2	34 58 35.4	17.78	2	F; cS; R; stellar; *16 nf ...	3
3708	1704	III. 285		13 49 33.3	3.125	1	94 48 28.8	17.76	1	vF; vS; R	2
3709	1708	II. 843		13 49 38.6	1.943	1	28 37 31.4	17.78	1	F; S	2
3710	1707	II. 716		13 49 46.8	+2.531	1	48 49 39.8	+17.76	1	pB; L; R; bMFN	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
3711	1709	III. 809		13 49 48.6	+2.031	1	30° 39' 35".1	+17.77	1	cF; S; E; ? * inv	2
3712	1710	II. 889		13 50 29.0	2.999	4	83 12 55.9	17.73	4	cF; pL; R; vgbM; *11 np...	5
3713	1711			13 50 36.8	2.715	3	60 8 59.9	17.73	3	pB; pL; R; lbM	3
3714		II. 844		13 50 38.7	1.987	1	29 47 45.9	17.73	1	pB; cL	1
3715		I. 238		13 50 40.4	1.986	3	29 48 15.9	17.73	3	cB; pL; vLE; vgmbM	3
3716	1712	I. 187		13 50 41.1	2.382	4	42 4 55.9	17.73	4	B; L; mE 40° 4; smbMN ..	5
3717	1713			13 50 47.4	2.576	1	51 31 33.6	17.72	1	pB; IE; vglbM	1+
3718	1714	II. 698		13 50 56.6	2.579	3	51 42 17.3	17.71	3	F; cS; R; smbM	4
3719	3549			13 50 57.8	4.119	2	148 54 29.7	17.69	2	Cl; Ri; vC; pL; st 11...12...	2
3720		I. 239		13 51 6.5	1.973	1	29 35 14.3	17.71	1	pB; pS; E; mbM	3
3721	1715	III. 546		13 51 15.6	2.996	1	83 3 41.7	17.69	2	vF; vS; r; stellar	3
3722	1715, a		R. nova	13 51 ±		...	83 0 ±		...	F; S; R	0
3723	1717	I. 181		13 51 17.9	2.496	1	47 28 19.0	17.70	1	cB; cL; R; gbM	2
3724	1721			13 51 19.2	0.414	1	13 7 58.9	17.73	1	Cl; P; S	1
3725	1716	III. 547		13 51 22.6	2.996	1	82 58 36.7	17.69	2	vF; vS; biN; r; stellar	3
3726	1719	I. 240		13 51 30.0	1.968	1	29 33 27.0	17.70	1	pB; pL; E; mbMN	3
3727	1718			13 51 31.1	2.524	1	48 52 36.7	17.69	1	F; L; vgbM; *9 nf.....	1
3728	1720	III. 666		13 52 10.9	3.101	1	92 31 21.8	17.66	1	vF; cS; R; gbM	2*
3729	3550			13 52 33.0	3.414	1	118 11 17.2	17.64	1	vF; S; R; glbM	1
3730	1722	I. 191		13 52 34.9	2.575	2	51 52 3.5	17.65	2	cF; S; np of 2	3
3731	1723	I. 190		13 52 38.2	2.575	3	51 53 52.5	17.65	3	cF; cL; E 15° 0; lbM; sf of 2	4
3732		III. 125		13 52 39.4	2.709	1	60 11 32.5	17.65	1	vF; S; iR; sbM*	1
3733	3551			13 52 55.7	3.492	1	123 16 22.6	17.62	1	vF; S; R; gbM	1
3734	3552			13 53 10.9	3.479	1	122 23 20.3	17.61	1	pB; pL; R; vgbM	1
3735	1724	III. 411		13 53 24.5	2.618	2	54 32 37.3	17.61	2	eF; vS; pmE 90°	3
3736		III. 667		13 53 31.9	3.097	2	92 10 41.0	17.60	2	vF; cS	2
3737	1725	III. 412		13 53 39.8	2.592	1	53 4 6.0	17.60	1	cF; cS; E	2
3738	1727	III. 810		13 53 41.9	1.944	1	29 28 38.3	17.61	1	vF; vS; R	2
3739	1726	III. 683		13 53 51.9	2.557	1	51 8 1.0	17.60	2	vF; pL; iF	4
5075				13 53 58.0		...	89 13 53		...	See No. 5075.	
3740	1728	II. 699		13 54 24.3	2.541	2	50 24 2.1	17.57	2	F; pS; R; lbM	5
3741	1732	III. 684		13 54 50.2	2.535	(1)	50 9 17.8	17.56	1	vF; vS; R; bM; in Cl	2
3742	3553			13 54 50.7	3.632	1	130 44 6.2	17.54	1	eF; E bet 2 vS st.....	1
3743	1729	II. 672		13 55 0.0	2.498	1	48 19 36.5	17.55	1	pF; pS; bM	2
3744		III. 56		13 55 0.2	2.958	1	79 54 37.2	17.54	1	eF; vS; E; r	1
3745	1733			13 55 10.5	1.647	1	24 24 32.5	17.55	1	pF; pS; R; psbM	1
3746	1730	III. 11		13 55 17.4	2.974	4	81 17 11.9	17.53	4	cF; S; R; psbM; *p	5
3747	1731			13 55 20.0	2.978	1	81 38 22.9	17.53	1	vF; R; bM	1
3748	3554			13 55 24.3	3.501	3	123 17 44.6	17.52	3	pB; pL; R; gpmbM	3
3749	1736	I. 230		13 55 44.4	2.117	2	34 9 20.6	17.52	2	pB; S; pmE 45° ±; vsymbMN.	4
3750	1734	II. 309		13 55 59.7	3.134	1	95 18 37.0	17.50	1	pF; cL; R; gmbM; np of 2 ..	3+*
3751	1735	II. 310		13 56 3.7	3.134	1	95 21 25.7	17.49	1	pF; cL; R; sf of 2	3*
3752	1738	II. 827		13 56 14.1	1.945	1	29 58 35.3	17.51	1	pB; S; iE; mbM	2
3753	1737	III. 653		13 56 26.8	2.646	2	56 49 5.4	17.48	2	vF; cS; iE 0°; bM	3
3754	1739	II. 416		13 56 57.7	2.606	1	54 33 31.8	17.46	1	pF; cS; iE; bM; *11 sp.....	3
3755	1740			13 57 18.2	2.607	1::	54 40 40.5	17.45	1	vF; S	3
3756	1741	II. 417		13 57 19.7	2.598	3	54 11 30.5	17.45	3	pB; pL; ivLE; vsmbM	4
3757	1742	III. 413		13 57 25.7	2.599	1	54 18 33.2	17.44	1	F; *13 p	2
3758		II. 799	H. O. N.	13 57 26.0	2.080	1	33 30 33.5	17.45	1	pF; L; E	2
3759		III. 57		13 57 30.0	2.953	1	79 42 30.9	17.43	1	cF; cS	1
3760	1744, a	III. 787?	R. nova?	13 57 30.5	2.137	...	35 5 18.3	17.44	...	B; S; R; gmbM; conn with M. 101.	0
3761	1743	II. 691		13 57 30.6	2.293	1	40 9 16.2	17.44	1	pB; cL; vmE 90° ±; smbMN	2
3762	1744, b		R. nova	13 57 32.8	2.134	...	35 0 28.3	17.44	...	vF; pL; gvlbM } all conn	0*
3763	1744, c		R. nova	13 57 33.0	2.137	...	35 5 51.3	17.44	...	F; pS; iR; glbM } with	0*
3764	1744, d		R. nova	13 57 41.7	+2.131	...	34 57 41.3	17.44	...	vF; pL; iR; vlbM } M. 101.	0*
3765	1747	III. 947		13 57 43.7	-0.245	1::	11 5 51.1	17.47	1	vF; pL; iR; vglbM	2
3766		III. 787		13 57 50.1	+2.133	1	35 3 26.9	17.43	1	vF; vS	1+*
3767	1744, e		R. nova	13 57 59.7	2.131	...	35 2 3.3	17.43	...	F; pL; iE; vlbM } conn with	0*
3768	1744, f		R. nova	13 58 2.8	+2.134	...	35 8 16.3	+17.43	...	pB; pS; R; psbM } M. 101.	0*

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
3769	h.	II.	D'Arrest, 95	h m s	s	[2]	77° 27' 4"	+17.40	[2]	F; pS	0
3770	1744		M. 101	13 58 12.9	2.127	1	34 57 22.3	17.41	1	pB; vL; iR; g, vsmbMBSN..	5*
3771	1744, g		R. nova	13 58 14.1	2.131	...	35 2 42.3	17.41	...	vF; pL; R; vlbM; conn with M. 101.	0*
3772	3555		Δ. 431	13 58 39.9	3.807	3	137 38 52.1	17.37	3	Cl; vL; vLC; st 8...	3
3773	1744, h	III. 788?	R. nova?	13 58 49.8	2.125	...	34 59 40.3	17.39	...	B; pS; R; psbM } conn with	0*
3774	1744, i	III. 789?	R. nova?	13 58 53.9	2.122	...	34 56 57.3	17.38	...	pB; pL; iR; gbM } M. 101.	0*
3775	3556			13 59 1.0	3.449	1	119 20 28.8	17.36	1	pF; S; R; psbM	1
3776	1746	VI. 9		13 59 8.7	2.700	2	60 48 11.1	17.37	2	Cl; L; vRi; vmC; st 11...	4
3777	1745	III. 286		13 59 14.1	3.129	1	94 46 56.5	17.35	1	F; L; R; vgbM	3
3778		III. 788		13 59 19.3	2.120	1	34 59 23.1	17.37	1	vF; vS; 2nd of 3 } inv in {	1†
3779		III. 789		13 59 26.2	2.118	1	34 57 22.8	17.36	1	vF; vS; 3rd of 3 } M. 101. {	1†
3780		III. 58		13 59 30.1	2.955	1	79 57 25.5	17.35	1	eF; S; IE	1
3781	1745, a		R. nova	13 59 30.2	3.129	::	94 46 56.5	17.36	::	pB; S; E	0
3782	1748	I. 231		13 59 50.7	2.096	1	34 25 50.5	17.35	1	pB; S; R; gbM	3
3783		I. 214		13 59 59.3	2.149	2	35 40 25.5	17.35	2	pB; L; bM	2
3784	1750	II. 800		14 0 20.7	2.061	1	33 35 24.6	17.32	1	pB; S; pmE; bM	2
3785	1749			14 0 34.6	2.994	1	83 17 26.0	17.30	1	F; mE; vglbM	1
3786	1751	III. 287		14 0 49.1	3.138	(2)	95 25 38.7	17.29	1:	F; pS; iR	3
3787		III. 790		14 0 54.3	2.104	1	34 52 19.0	17.30	1	vF; pL	1
3788		III. 762		14 0 55.8	3.085	1	91 1 21.4	17.28	1	vF; vS	1
3789		II. 692		14 1 24.6	2.225	1	38 37 18.1	17.27	1	F; pS; vgbM; np of 2	1
3790		II. 693		14 1 42.5	2.223	1	38 37 47.8	17.26	1	F; vS; smbM; stellar; sf of 2	1
3791		III. 59		14 1 43.2	2.958	1	80 24 17.5	17.25	1	eF; S	1
3792	3557			14 1 46.3	3.704	1	132 39 20.2	17.24	1	pF; vL; R; vgbM	1
3793		III. 791		14 2 26.5	2.073	1	34 19 14.9	17.23	1	vF; S; R; p of 2	1
3794		I. 232		14 2 26.5	2.073	1	34 19 14.9	17.23	1	cB; R; vgbM; f of 2	1
3795		II. 801		14 2 34.8	2.071	2	34 16 45.6	17.22	2	F; pL	2
3796	3558			14 2 45.6	3.520	1::	122 58 35.7	17.19	1::	F; R; *8 s nr	1
3797	3559			14 3 10.0	3.775	2	135 25 33.1	17.17	2	vF; S; R; bM	2
3798	1752	III. 32		14 3 17.4	2.846	1	71 47 18.4	17.18	2	cF; cS; R; sbMF*	4
3799	1753	II. 890		14 3 59.6	2.988	3	82 58 10.5	17.15	3	pB; pS; R; gbM; r	5
3800	1754	II. 876		14 4 1.8	2.817	1	69 43 45.5	17.15	1	pB; vS; E	3
3801	1755	IV. 46		14 4 12.2	3.126	1	94 22 49.9	17.13	1	pB; vS; R; psmbM*; *18 inv.	2
3802	3560			14 4 16.2	3.475	1	119 59 46.6	17.12	1	pB; L; R; gbM; rr	1
3803	3561			14 4 23.5	3.419	1	116 26 51.6	17.12	1	vF; S; R; bM; *sf	1
3804		III. 674		14 5 5.6	2.265	2	40 46 9.3	17.11	2	cF; cS; iR	1
3805	1756			14 5 13.3	3.050	1	88 5 4.7	17.09	1	vF; S; rr	1
3806	1757	II. 687		14 5 59.1	3.105	1	92 32 43.5	17.05	1	pB; L; E 20°±; lbM	3
3807	1757, a		R. nova	14 6 ±	3.105	...	92 33 ±	17.05	...	3' dist from h. 1757	0
3808	1758	IV. 49		14 6 5.9	3.104	1	92 29 28.5	17.05	1	cF; S; R; stellar	3
3809	1759	II. 877		14 6 36.4	2.801	1	68 54 40.9	17.03	1	pB; pL; iR	2
3810	1760	III. 685		14 6 51.3	2.482	3	50 1 53.6	17.02	3	vF; S; vIE	5
3811	3562			14 6 54.8	3.848	2	137 27 28.0	17.00	2	pF; S; R; psbM; S *nf	2
3812		III. 676		14 7 49.9	2.196	2	38 59 31.4	16.98	2	F; S; IE; stellar	2
3813	1761			14 8 20.7	3.010	1	84 56 4.5	16.95	1	F; S; R; bM	1
3814		III. 644		14 8 23.6	2.871	1	74 13 59.5	16.95	1	vF; vS; E; a D neb	1
3815	1762, a		R. 3 novæ	14 8 ±	2.725	...	64 0 ±	16.94	...	3 "knots" near h. 1762	0
3816											
3817											
3818	1762	III. 134		14 8 31.0	2.725	3	64 0 55.2	16.94	3	F; pL; pmE 90°; *10 np	5
3819	1764, a		R. nova	14 8 47.0	2.53	::	52 56 57.7	16.93	::	eeF	0
3820	1763	III. 804 = III. 835		14 9 17.5	1.899	1	31 34 24.3	16.91	1	vF; S; E; r	3*
3821	1764, b		R. nova	14 9 18.9	2.53	::	53 6 33.7	16.89	::	vF	0
3822	1764	III. 414		14 9 40.1	2.533	1	53 7 33.7	16.89	1	cF; pL; vmE 110°3; vgvmbM.	2
3823	3563			14 9 41.8	3.742	1	132 43 31.1	16.87	1	l; vF; pmE; esvmbM *12	1
3824			D'Arrest, 96	14 9 53	2.92	[2]	78 28 0	16.87	[2]	F; S; R; III. 47 f 10°4	0
3825	1765	III. 47		14 10 2.1	+2.926	1	78 32 18.1	+16.87	1	vF; vS; R; gbM; r	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and h.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
3826	h. 1766	II. 418		h m a s	+2.549	3	53° 59' 51".1	+16.87	3	pB; R; vsmbM; 2 or 3 st inv	4
3827	1768	III. 731		14 10 40.3	2.462	1	49 51 34.2	16.84	1	cF; vS; R; p of 2	2
3828	1767			14 10 43.5	2.960	1	81 10 12.2	16.84	1	F; pL; iF; gbM	1
3829		III. 805		14 10 44.8	1.789	3	29 20 6.5	16.85	3	cF; vS; R; stellar	3
3830	1770, a		R. nova	14 10 45.3	2.968	::	81 39 52.2	16.85	::	vF	0
3831	1769	III. 732		14 10 48.4	2.459	3	49 45 41.2	16.84	4	cF; S; R; gbM	5
3832	1770, b		R. nova	14 10 53.9	2.968	:	81 46 46.3	16.83	:	vF	0
3833	1771	II. 419		14 11 8.5	2.521	2	52 46 41.6	16.82	3	F; pS; E 80°; D or biN	4
3834	1771, a		R. nova	14 11 ±	2.521	...	52 46 ±	16.82	...	Makes D or biN neb with h. 1771.	0
3835	1770			14 11 13.9	2.968	2	81 46 56.3	16.81	2	pB; cS; gbM	2
3836		III. 551		14 11 29.6	+2.970	2	81 59 ±	16.80	2		2*
3837		III. 948		14 11 36.1	-0.747	1	10 44 46.7	16.79	1	eF; vS; E 0° ±	1
3838	1773	II. 194		14 11 38.1	+2.720	2	64 12 39.0	16.80	2	cF; pS; R; vsvbmM*	4
3839	1772	III. 552		14 11 43.6	2.970	1	81 58 40.7	16.79	1	vF; vS; R	2
3840	1774			14 11 43.8	2.897	1	76 27 52.7	16.79	1	vF; cS; pmE	1
3841	1775			14 12 12.9	2.701	1	63 4 49.8	16.76	1	vF; S; IE	1
3842	3564			14 12 24.5	3.475	1	118 36 20.5	16.75	1	eF; L; S * inv.	1
3843	1776	I. 99		14 12 32.5	2.517	2	52 51 11.8	16.76	2	cB; S; R; vsbM*	4
3844	1777	III. 347		14 12 50.8	2.723	1	64 32 41.9	16.73	1	vF; S; vIE; bM	2*
3845	1778	II. 579		14 13 1.0	3.013	2	85 21 16.6	16.72	2	pF; cL; E; gbM	3
3846	1779	I. 144		14 13 16.5	3.014	3	85 25 7.3	16.71	3	B; pL; R; psbM; r; *12 nf.	4*
3847	1780			14 13 20.8	2.541	1	54 13 56.6	16.72	1	pF; R	1
3848	1779, a		R. nova	14 13 28.5	3.014	...	85 24 7.3	16.72	...	Place from MS.	0
3849	1781	III. 12		14 13 31.7	2.967	(1)	81 50 21.0	16.70	2	F; S; iR	4
3850	1782	I. 145		14 13 52.1	3.023	1	86 6 53.4	16.68	1	pF; pS; IE; p of 2	2
3851	1783	I. 146		14 14 1.1	3.022	1	86 5 8.4	16.68	1	B; S; R; vsmbM; f of 2	2
3852	1784	III. 415		14 14 31.5	2.536	2	54 9 46.8	16.66	2	vF; cL; p of 2	3
3853	1785			14 14 45.4	2.535	1	54 8 52.2	16.64	1	pB; S; f of 2	1
3854	1786	II. 754		14 15 1.5	2.440	3	49 39 28.2	16.64	3	pB; pS; R; bMFN; *sp	5
3855	1783, a		R. nova	14 15 5.4	3.02	...	85 53 50	16.64	...	L; F; vmE	0
3856	1790	I. 235		14 15 18.1	1.909	1	32 38 5.9	16.63	1	pF; L; iR; vgbM; r	3
3857	1787	III. 110		14 15 30.3	2.879	1	75 26 38.0	16.60	1	F; cS; vIE; *8 sf	3
3858	1789			14 15 30.7	2.534	1	54 13 57.3	16.61	1	vF; R; gbM; *8 sf	1*
3859	1788	III. 416		14 15 30.8	2.531	2	54 5 13.3	16.61	2	vF; S; R; np of 2	3*
3860	1791	III. 417		14 15 43.3	2.532	3	54 9 17.0	16.60	3	cF; S; R; bM*; sf of 2	4*
3861	3565	III. 924		14 15 47.9	3.474	1	118 1 55.4	16.58	1	F; S; E; gvlbM; r	2
3862	3566		Δ. 357	14 16 13.4	4.121	1	144 9 51.1	16.55	1	Cl; vIRi; vIC; st 10	1
3863		III. 135		14 16 25.3	2.689	1	62 58 37.8	16.56	1	eF; vS; stellar	1*
3864	1792	III. 121		14 16 32.5	3.291	1	106 5 6.5	16.55	2	F; pL; R; vgbM; p of 2	4
3865	1795	III. 418		14 16 41.3	2.489	1	52 14 23.5	16.55	1	eF; S; R; stellar	2
3866	1793	III. 122		14 16 46.6	3.292	1	106 7 34.2	16.54	2	vF; L; vIE; vglbM; f of 2	4
3867	1796	III. 733		14 16 49.7	2.419	1	49 2 30.5	16.55	1	F; vS; R; bM	3
3868	1794	III. 927		14 16 53.9	2.978	4	82 45 50.2	16.54	4	F; S; IE	5
3869	1797	II. 177		14 17 7.8	2.866	1	75 43 22.6	16.52	1	pB; pS; gbM	3
3870		II. 694		14 17 17.0	2.134	1	38 49 32.9	16.53	1	pF; pS; IE; mbM	1
3871	1800	III. 734		14 17 24.4	2.414	2	48 58 44.6	16.52	2	cF; pS; R; gbM	3
3872	1799	III. 668		14 17 28.1	3.107	1	92 34 12.0	16.50	1	F; pS; R; vgbM*; r	2
3873	{ 1798 = 3569 }	III. 120		14 17 32.1	3.243	2	102 32 27.0	16.50	2	vF; pL; R; vgbM	3
3874	3568		Δ. 313	14 17 36.5	4.345	1	148 59 58.4	16.48	1	Cl; S; pC; st L & S	1
3875		II. 331		14 17 43.9	0.726	2	17 47 31.9	16.53	2	pF; cS; iR; bM; cr	2
3876	1801	II. 673		14 17 48.1	2.380	1	47 35 13.0	16.50	1	F; pL; IE; vglbM	2
3877	1802	III. 136		14 18 3.6	2.714	1	64 45 34.4	16.48	1	vF; S; pmE 0° ±; *9 f	4
3878	1803			14 18 6.3	2.566	1	56 18 59.4	16.48	1	F; S; R; bM	1
3879	3567			14 18 9.9	6.643	1	167 46 0.3	16.41	1	vF; E; gbM; r	1
3880	1804	II. 420		14 18 12.4	2.530	2	54 29 56.4	16.48	2	pB; S; R; smbM	3
3881	} 1804, a		R. 3 novæ	14 18 ±	+2.530	...	54 30 ±	+16.48	...	h. 1804 is D; 2 others near ..	0
3882											
3883											

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and l.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
3884	1805	III. 419		14 18 31.3	+2.496	1	52° 54' 5.8"	+16.46	1	vF; S; cE; vgbM; er.....	2
3885	3570		Δ. 302	14 19 23.2	4.417	3	150 5 16.7	16.39	3	Cl; L; pRi; pCM; st 8...	3
3886		III. 763		14 20 1.3	3.095	1	91 37 28.4	16.38	1	eF; S	1
3887	1806			14 20 15.9	3.000	1	84 33 53.8	16.36	1	vF; S; R; vgbM	1
3888		III. 319		14 20 17.6	0.680	1	17 44 24.0	16.40	1	eF; vS	1*
3889	1807	III. 14		14 20 54.8	2.952	1	81 7 26.2	16.34	1	eef; L; r.....	2
3890	1809	III. 677		14 21 3.9	2.177	1	40 48 47.2	16.34	1	vF; pS; vIE; vglbM	2
3891	1808	II. 329		14 21 12.5	2.552	1	56 7 10.9	16.33	1	cF; S; R; vsnbM; r	4
3892	1810			14 21 25.6	2.408	1	49 24 57.6	16.32	1	vF; S; R; gbM	1
3893	3571			14 21 35.0	3.508	1	119 7 31.7	16.29	1	eF; S; R	1
3894	1811			14 21 47.6	2.907	1	77 59 10.7	16.29	1	vF; vS; R; *9sp	1
3895	1812			14 21 59.6	2.685	2	63 31 30.7	16.29	2	pF; S; R; gbM	2
3896	1814	II. 674		14 22 4.2	2.374	4	48 6 50.7	16.29	4	F; S; E 90°±; gbM	5
3897	1820	I. 236		14 22 10.2	1.865	2	32 47 47.7	16.29	2	B; S; R; psbMrN	5
3898			Auw. N. 33	14 22 12.0	3.068	...	89 49 7.8	16.26	...	Neb *11f 150° (Bond, May, 1853).	0
3899	1818	I. 185		14 22 14.2	2.243	2	43 13 13.4	16.28	2	cB; pS; R; pglbM	4
3900	1813	I. 70		14 22 14.9	3.146	1	95 20 26.8	16.26	1	⊕; vB; cL; R; gbM; rrr; st 19; *17 sf.	3
3901	1815	III. 132		14 22 20.8	2.657	1	61 57 59.1	16.27	1	F; S; E; sbM	2
3902	1816	II. 580		14 22 34.6	3.020	1	86 5 58.2	16.24	1	eF; cL; R; np of 2	2
3903	1819	II. 357		14 22 35.7	2.728	2	66 11 4.5	16.25	2	vF; S; R; vgbM	3
3904	1817	II. 581		14 22 36.8	3.020	2	86 8 14.2	16.24	2	cB; pL; R; sf of 2	3
3905	1817, a		R. nova	14 23 ±	3.020	...	86 8 ±	16.24	...	Makes a BD neb with h. 1817	0
3906	1821			14 22 41.4	+2.602	1	58 58 19.5	16.25	1	vF; R; *7 p; *11s	1
3907		III. 949		14 23 2.9	-1.635	1	9 18 14.0	16.30	1	eF; S; IE	1
3908	1822	III. 126		14 23 11.4	+2.608	2	59 21 35.9	16.23	2	cF; S; * inv; *12 nf	3
3909	3572		Δ. 469	14 23 40.3	3.826	2	133 34 30.1	16.17	2	pB; L; R; vglbM; st inv ...	2
3910	1823	II. 150		14 23 43.8	2.964	2	82 5 56.7	16.19	2	cF; pL; iR; gbM	5
3911	1824	III. 645		14 23 54.1	2.867	1	75 22 18.4	16.18	1	eF; vS; np of 2	2
3912			Auw. N. 34	14 24 3.0	3.066	...	89 42 5.1	16.17	...	Neb R (Bond, May, 1853)...	0
3913	1825	II. 891		14 24 4.0	2.982	2	83 23 45.1	16.17	2	pB; pL; vIE; bM	3
3914	1826	II. 330		14 24 7.8	2.583	1	58 9 43.1	16.17	1	pF; pS; R; bM	2
3915	1828	III. 420		14 24 14.9	2.478	1	53 0 46.8	16.16	1	F; S; E?; * inv?	2
3916	1827			14 24 17.5	2.868	1	75 27 56.8	16.16	1	eeF; sf of 2	1
3917	1829	II. 421		14 24 34.5	2.499	3	54 2 59.5	16.15	4	pF; pL; R; mbM; r	5
3918			Auw. N. 35	14 24 48.0	3.067	...	89 45 4.8	16.16	...	Neb; F; E (Bond, May, 1853)	0
3919	1831			14 24 52.7	2.686	1	63 59 8.9	16.13	1	eF	1
3920	1832	II. 695		14 24 53.3	2.122	1	39 45 49.2	16.14	1	pB; L; iR; vgbM	2*
3921	1830	II. 892		14 24 59.9	2.977	2	83 7 34.6	16.12	2	vF; pS; iE	3
3922	3573		Δ. 342	14 25 7.9	4.256	2	145 56 31.0	16.10	2	Cl; L; pRi; IC; st 9...	2*
3923	1833	II. 27		14 25 31.6	2.951	2	81 18 15.0	16.10	2	pB; pL; R; gbM; r	5
3924	1834			14 26 19.7	2.916	1	78 51 33.5	16.05	1	vF; vS; R; stellar	1
3925		II. 807		14 26 22.6	1.686	1	29 54 6.1	16.07	1	pB; pS; E 0°	1
3926	1835	II. 574		14 26 22.9	3.002	2	84 55 54.5	16.05	2	F; pS; vIE; *14 inv	3
3927		II. 79		14 26 28.2	2.924	1	79 28 9.2	16.04	1	F; L; R; lbM; r	1
3928	3574			14 26 28.4	3.887	2	135 20 29.9	16.03	2	vF; S; cE; bet 2 st	2
3929		III. 882		14 26 35.4	0.855	1	19 44 5.4	16.08	1	vF; pL; R; bM	1
3930	1836	III. 310		14 26 39.7	2.563	1	57 43 21.2	16.04	1	vF; vL; iR; lbM; * p	3
3931	1838	II. 696		14 26 42.1	2.100	1	39 26 14.5	16.05	1	F; S; cE; *15 np	2
3932	1837	II. 893		14 26 53.5	2.988	4	83 55 1.9	16.03	4	cF; pS; R; gbM	5
3933	1839	II. 422		14 26 53.6	2.470	1	53 4 35.9	16.03	1	F; pS; E; bM	2
3934	1843	I. 237		14 27 51.2	1.758	1	31 27 43.7	15.99	1	B; L; IE 0°; vgmbM	3
3935	1842	I. 189		14 27 51.3	2.110	1	39 55 14.7	15.99	1	B; L; E 45°±; pgbM; r ...	2
3936	1840	III. 283		14 27 55.2	2.678	1	63 54 56.1	15.97	1	vF; vS; R; r; 3 st 9, 10 np...	2
3937	1841	II. 894		14 28 8.6	2.988	2	84 1 3.8	15.96	4	vF; S; R; *12 att	6
3938	1844	III. 421		14 30 6.5	2.454	4	52 50 57.8	15.86	4	F; cS; R; bM; p of 2	5
3939	1845			14 30 18.9	2.454	1	52 53 2.5	15.85	1	vF; S; R; f of 2	2
3940	1849	II. 808		14 30 26.8	1.902	1	34 53 55.5	15.85	1	pF; S; iF; r; *10 f	2
3941	3575			14 30 28.4	3.880	1	134 25 46.3	15.81	1	F; S; vgbM; am st	1
3942	1848	I. 188		14 30 29.8	+2.120	1	40 38 43.5	+15.85	1	cB; S; E 90°±; psmbM ...	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.			Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by H. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.	h	m	s							
3943	1848, a		R. 3 novæ	14	30	±	+2.120	...	40 38 ±	+15.85	...	3 novæ, one mottled	0
3944				14	30	39.7	3.032	1	87 6 6.6	15.82	3	vF; mE or biN 140° ±; *6.7 p	4
3945				14	30	42.5	3.070	1	89 46 46.6	15.82	1	pB; pS; iE; gbM	3
3946	1846	II. 582		14	30	42.5	3.070	1	89 46 46.6	15.82	1	pB; pS; iE; gbM	3
3947	1847	II. 681		14	31	38.1	2.448	2	52 49 33.4	15.78	3	pB; cS; R; bM; r	4
3948	1851	II. 423		14	31	38.1	2.448	2	52 49 33.4	15.78	3	pB; cS; R; bM; r	4
3949	1851, a		R. 2 novæ	14	32	±	2.488	...	52 50 ±	15.78	...	No description	0
3950				14	32	±	2.488	...	52 50 ±	15.78	...	No description	0
3951				14	31	38.2	2.321	(2)	47 36 0.4	15.78	1	cF; cS; R; lbM; r	3
3952	1853	II. 675		14	31	38.8	2.325	1::	47 45 53.4	15.78	1::	F; vS; R; bM; 4B st p	2
3953	1852	II. 700		14	31	38.9	2.404	2	50 55 57.4	15.78	3	cF; cS; iE; in Δ of st	4
3954	3576	II. 196		14	32	4.2	3.477	1	115 55 37.2	15.74	1	cB; cS; R; psbM; r; * nr ...	3
3955		III. 127		14	32	12.5	2.594	1	59 53 49.2	15.74	1	eF; vS	1
3956	1854	II. 575		14	32	12.8	2.986	4	84 1 23.2	15.74	4	cB; pS; R; mbM; *15 p ...	5
3957		III. 894		14	32	20.6	2.754	2	68 53 19.9	15.73	2	vF; vS	1
3958		III. 128		14	32	26.5	2.593	1	59 53 48.9	15.73	1	vF; vS; iR	1
3959	1855	II. 649		14	32	26.6	2.351	1	48 52 33.2	15.74	1	F; cS; iE 0° ±	4
3960	1859			14	32	46.2	2.351	2	48 56 49.6	15.72	2	F; pL; E 0° ±; gbM	2
3961	1856	III. 895		14	32	50.2	2.760	1	69 21 2.0	15.70	1	vF; S; vgbM; * f; p of 2 ...	2
3962	1858			14	32	55.7	+2.761	1	69 24 52.0	15.70	1	eF; vS; * att; f of 2	1
3963		III. 950		14	32	57.9	-1.318	1	10 33 44.4	15.78	1	vF; S; R; S Cl p	1
3964	1857	I. 182		14	32	59.9	+3.068	1	89 40 43.0	15.70	1	cB; pL; R; psmbM; r	3
3965	1861	III. 675		14	33	2.8	2.175	1	42 44 30.0	15.70	1	vF; pS; iE; * n; 1st of 3 ...	2
3966	3577		Δ. 333	14	33	15.1	4.353	1	146 56 49.8	15.66	1	Cl; L; pRi; CM; st 11...13...	1
3967		VI. 8		14	33	16.5	3.198	1??	98 30 48.4	15.68	1??	Cl; pL; eRi; vmC	1*
3968	1860	III. 671		14	33	19.0	3.326	2	106 52 1.1	15.67	2	vF; pL; R	3
3969	1864			14	33	29.2	2.172	2	42 42 29.7	15.69	1	vF; S; R; * nr; 2nd of 3 ...	2
3970	1862	III. 550		14	33	39.6	3.014	1	85 56 1.8	15.66	2	vF; S; R; vglbM; *8.9 nf...	3
3971	1863	II. 682		14	33	44.9	3.069	1	89 42 24.5	15.65	1	pF; S; iE; bM	3
3972	1865, a		R. 3 novæ	14	34	±	2.170	...	42 41 ±	15.67	...	h. 1865 is quadruple; ? F neb connecting.	0
3973				14	33	49.6	2.170	1	42 40 59.1	15.67	1	vF; S; R; psbM; 3rd of 3...	1
3974				14	33	53	3.03	[1]	87 12 42	15.64	[1]	○?; vF; S; disc; *15 n 95''...	0
3975	1865		D'Arrest, 97	14	33	53	3.03	[1]	87 12 42	15.64	[1]	○?; vF; S; disc; *15 n 95''...	0
3976				14	34	38.1	3.325	2	106 38 34.0	15.60	2	pF; pL; pmE 45° ±; mbM; *10s.	3*
3977	1866	I. 184		14	34	38.1	3.325	2	106 38 34.0	15.60	2	pF; pL; pmE 45° ±; mbM; *10s.	3*
3978	3578	III. 508		14	34	40.0	3.197	2	98 24 7.0	15.60	2	F; pL; E; r	3
3979	1867	III. 657		14	34	41.8	2.281	1	46 36 11.3	15.61	1	vF; cS; E 90° ±	2
3980	1868	III. 658		14	34	51.9	2.281	1	46 37 41.0	15.60	1	vF; cS; iE	2
3981	1869	III. 686		14	35	2.8	2.387	2	50 45 12.7	15.59	2	vF; S; R; lbM	3
3982	1870	III. 133		14	36	29.1	2.598	1	60 40 46.3	15.51	1	vF; L; iR; lbM	2
3983	1871	III. 896		14	36	42.8	2.773	1	70 31 5.7	15.49	1	vF; cS; R; vglbM	2
3984	1873	I. 171		14	37	12.6	2.297	1	47 34 26.1	15.47	1	pB; S; R; smbM; r; * nr ...	3
3985	1872	II. 538		14	37	17.6	3.039	1	87 43 15.8	15.46	1	pB; L; iR; gbM; r	2
3986	3579			14	37	23.1	3.276	1	103 20 50.5	15.45	1	vF; S; E; psbM	1
3987	1874	I. 126		14	37	48.7	3.035	1	87 26 59.9	15.43	1	B; L; vmE; bMBN	2
3988		III. 48		14	38	25.5	2.880	1	77 18 32.0	15.40	1	eF; S	1
3989	3580		Δ. 356	14	38	57.4	4.258	1	143 56 9.2	15.34	1	Cl; pL; pRi; IC; st 10...11...	1
3990	1875	I. 183		14	39	9.0	3.068	1::	89 38 40.5	15.35	1::	pF; pS; vIE; r	3
3991	1877	II. 809		14	39	24.3	1.892	1	35 59 5.8	15.36	1	F; S; vIE; Δ 2 st 10.11 ...	2
3992	1878	III. 687		14	39	47.5	2.368	3	50 40 19.9	15.33	3	cF; cS; R; bM	4
3993	3581			14	39	54.0	3.293	1	104 15 56.3	15.31	1	pB; pL; pmE; gpmbM	1
3994	1876	III. 690		14	39	54.2	3.362	1	108 29 45.3	15.31	1	vF; S; iR; lbM	2
3995	1879	III. 885		14	41	13.2	2.773	1	70 54 33.2	15.24	1	vF; vS; cE 90°; vglbM	2
3996	3582			14	43	44.7	4.208	1	142 5 25.1	15.07	1	Cl; vF; vS; vC	1
3997	1880			14	43	52.3	2.985	1	84 17 59.7	15.09	1	D neb; both eF	1
3998		III. 373		14	44	58.6	3.104	3	91 57 9.6	15.02	3	F; R; bMFN; S * s	3*
3999	1881	II. 576		14	46	13.3	3.003	1	85 27 35.5	14.95	1	cF; S; vIE; bM; ?biN	4*
4000	1882	III. 129		14	46	16.8	2.551	1	59 35 3.5	14.95	1	vF; S; R; pgbM	2
4001	1883			14	46	17.5	2.295	3	48 49 8.8	14.96	3	pB; pL; iE; psbM; *8 np...	3
4002	1884	III. 130		14	46	34.3	+2.551	1	59 37 52.9	+14.93	1	vF; S; R; pgbM	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. II.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s						
4003	1885, <i>a</i>		R. nova	14 46 50.9	+3.009	::	85 53 7.0	+14.91	:	eF; 2' p h. 1885	0
4004	1885	III. 554		14 46 58.9	3.009	1	85 53 7.0	14.90	1	F; pS; vmE 148° 4; gvlbM...	4
4005			D'Arrest, 98	14 47 33	3.02	[1]	86 27 8	14.87	[1]	vF; pL; vlbM; *8.9p; 225'' s	0
4006		III. 806		14 47 33.0	1.555	1	30 26 56.7	14.89	1	vF; vS; IE	1
4007	1886			14 48 52.2	3.344	1	106 40 4.7	14.79	1	F; S; R; bM; *16 sp.....	1
4008	1887	II. 676		14 49 5.8	2.229	2	46 52 39.0	14.80	2	pB; S; R; smbM; stellar ...	3
4009	3583			14 49 57.8	3.877	1	131 27 21.3	14.71	1::	F; mE; L * sf	1
4010	1888	II. 677		14 50 2.4	2.227	1	46 56 4.2	14.74	1	F; cS; R; psbM	2
4011	1890	III. 976		14 50 41.1	2.532	1	59 12 55.0	14.70	1	cF; pS; iF	2
4012	1889	III. 691		14 50 51.7	3.382	1	108 42 27.1	14.67	1	pF; S; R; stellar	2
4013		II. 683		14 51 12.5	3.081	1	90 31 47.8	14.66	1	pB; pL; R; mbM; cB*npatt	1
4014	1891			14 51 21.5	1.979	1	39 44 42.1	14.67	1	pF; S; vsbM*13; 1st of 3 ...	1
4015	1893	III. 678		14 51 46.4	1.977	1	39 44 55.2	14.64	1	F; S; vsbM*13; 2nd of 3 ...	2
4016	1892	III. 131		14 51 48.3	2.534	1	59 28 16.9	14.63	2	F; S; R; vgbM; *nf (? var)	4*
4017	3584			14 51 49.4	5.894	1	161 52 27.1	14.57	1	eF; S; R; bM	1
4018	3585			14 52 5.8	4.224	1	141 21 29.4	14.58	1-	Cl; pL; pRi; iC	1
4019	1895	III. 679		14 52 31.1	1.973	1	39 45 19.0	14.60	1	vF; vS; vsmbM*13; *6 nr; 3rd of 3.	2
4020	1895, <i>a</i>		R. nova	14 52 ±	1.973	...	39 45 ±	14.60	...	S	0
4021	1894	II. 539		14 52 53.9	+3.034	1	87 32 48.8	14.56	1	cB; cL; E 165° ±; sbMN ...	2
4022		III. 311		14 53 0.6	-0.023	1	16 24 35.3	14.61	1	vF; S; iR; bet 2 st	1
4023	3586			14 53 11.2	+3.296	1	103 36 47.9	14.53	1	vF; S; E; glbM	1
4024	3587	I. 71		14 53 28.2	3.185	2	96 53 29.6	14.52	2	cB; S; R; svmbM	4
4025		II. 756		14 53 51.4	1.781	2	35 37 32.1	14.48	2	pF; pS; iE; r	2*
4026	1896	I. 127		14 54 5.4	3.037	1	87 44 16.7	14.49	1	B; pS; R; psmbM	2
4027	1897			14 54 15.4	3.038	1	87 49 46.1	14.47	1	vF; vS; R	1
4028		III. 811		14 54 22.9	1.784	1	35 35 32.7	14.49	1	vF; S; E	1
4029	1898	II. 756?		14 54 29.9	1.781	1	35 32 4.1	14.47	1	B; R; sbM; splendid * f ...	1*
4030	1898, <i>a</i>		R. nova	14 54 50.6	1.781	::	35 32 4.1	14.47	::	vF	0
4031	3588			14 55 2.0	4.347	1	143 47 24.0	14.40	1	Cl; vL; Ri; iC; st 9...12	1
4032	3589			14 55 20.2	4.409	1	145 2 17.4	14.38	1	Cl; cL; Ri; iCM; st 13...14	1
4033	1899	II. 540		14 57 2.4	3.045	(1)	88 14 41.0	14.30	1	pB; S; mbM	2
4034		II. 332		14 57 32.3	0.165	2	17 45 19.6	14.32	1	pB; cL; iR; bp; r	2
4035	3590			14 57 32.5	6.045	3	162 19 4.3	14.21	3	F; cS; iE; glbM; am st	3
4036	1900			14 57 46.9	+3.675	1	122 34 35.2	14.24	1	eeF(?)	1
4037		III. 312		14 57 54.8	-0.262	1	15 36 18.3	14.31	1	eF; vS; IE; 2 st inv	1
4038		II. 542		14 58 21.5	+3.030	1	87 21 21.6	14.22	1	pB	1
4039		II. 541		14 58 21.9	3.037	1	87 49 21.6	14.22	1	F	1
4040	3592			14 58 41.7	3.755	2	125 47 8.4	14.18	2	vF; S; IE; vlbM; r	2
4041	3591			14 58 51.0	5.033	2	154 8 9.5	14.15	2	pB; pL; R; vglbM	2
4042		III. 511		14 59 1.9	3.036	1	87 46 20.4	14.18	1	vF; R; p of 2	1
4043	1901, <i>a</i>		R. 2 novæ?	14 59 ±	3.038	...	87 51 ±	14.16	...	2 of 6	*
4044											
4045	1901	I. 128		14 59 23.5	3.038	1	87 50 41.8	14.16	1	vB; pL; R; psbMN; f of 2...	2
4046			D'Arrest, 99	14 59 29	3.03	[1]	87 26 48	14.15	[1]	eF; S; v close * f 7'	0
4047	1902	II. 543		15 0 2.6	3.039	1	87 54 47.6	14.12	1	cF; S; iE; psbM	2
4048		III. 886		15 0 16.5	2.848	1	76 36 4.3	14.11	1	eF; vS; np of 2	1*
4049		III. 887		15 0 16.5	2.848	1	76 36 4.3	14.11	1	eF; vS; sf of 2	1*
4050	1903	II. 544		15 0 44.0	3.021	3	86 53 20.4	14.08	3	pB; S; vIE; lbM; am st	5
4051	1905	II. 751		15 1 1.8	2.725	(3)	69 54 1.8	14.06	1::	cF; cS; E; p of D neb	4*†
4052	1905	II. 752		15 1 9.2	2.725	(3)	69 56 1.8	14.06	1::	pF; pS; E; f of D neb	4†
4053	1904	IV. 71		15 1 14.4	2.746	1::	71 1 0.5	14.05	1	*6 in vL neb	2
4054	1906			15 1 29.3	2.179	2	46 49 50.5	14.05	2	F; S; R; psbM	2
4055		II. 192		15 1 37.4	3.255	2	100 46 50.6	14.02	2	F; L; E; r	2
4056	1907	II. 585		15 2 30.8	3.013	3	86 24 19.8	13.96	3	pF; cS; iE; gbM; *14 f ...	4
4057		II. 684		15 2 36.1	3.057	1	89 0 5.5	13.95	1	pB; S; iE	1
4058	1909	I. 215		15 2 36.3	1.639	1	33 41 39.4	13.98	1	vB; cL; pmE 146° 0; gbM...	2†
4059	1909, <i>a</i>		R. nova	15 2 ±	1.639	...	33 42 ±	13.98	...	vS	0
4060			D'Arrest, 100	15 2 41	3.05	[1]	88 55 48	13.95	[1]	eF; H. II. 545, s 3' 15''	0
4061	1908	II. 545		15 2 41.0	3.057	1	88 59 25.5	13.95	1	pF; S; E; psbM	5
4062		II. 755		15 5 10.7	+1.792	1	36 55 21.6	+13.82	1	pB; pL; iE	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. II.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s						
4063	3593	III. 736		15 6 3.3	+3.312	1	103 44 18.9	+13.73	1	pB; pL; pmE0°; psmbM; * inv	2
4064	1910	II. 757		15 6 9.4	1.548	1	32 27 30.8	13.76	1	cB; S; E; mbMRN; r	4
4065		II. 818		15 6 27.0	1.128	1	26 30 47.5	13.75	1	pF; cS; R; vgbM	1*
4066	3594			15 7 19.4	4.059	2	135 7 30.2	13.64	2	O; vS; R; quite sharp	2†
4067	3595	III. 116		15 7 30.1	3.238	1	99 33 7.2	13.64	1	F; cL; R; vgbM	2
4068	1911			15 7 38.4	2.199	1	48 14 58.2	13.64	1	F; vS; R; bM	1
4069	1912	III. 659		15 7 58.6	2.198	1	48 12 57.2	13.64	1	cF; vS; R; bM; r	2
4070	1912, a		R. nova	15 8 18.6	2.198	...	48 8 27.2	13.62	...	vF; place from MS.	0
4071	1913	II. 678		15 8 29.8	2.175	1	47 31 15.0	13.60	1	F; S; R; r; 3 st nr.	2
4072	1913, a		R. 2 nova	15 8 ±	2.175	...	47 31 ±	13.60	...	2 nova apparently connected.	0
4073											
4074		II. 763		15 8 37.8	1.353	2	29 40 9.0	13.60	2	pF; pS; E 0° ±	2
4075	3596	VI. 19		15 9 22.4	3.443	1	110 29 44.6	13.52	1	⊕; pF; L; viR; vgbM; rrr	2
4076	3597	III. 138		15 9 53.3	3.504	1	113 31 44.7	13.49	1:	F; S; R; gbM	4
4077	1914	II. 650		15 10 0.6	2.167	2	47 26 25.3	13.51	2	cB; pL; pmE; smbMN	5
4078	1915	III. 660		15 10 2.9	2.162	2	47 16 49.0	13.50	2	vF; S; vLE; gbM	3
4079	1915, a		R. nova	15 10 2.9	2.162	...	47 17 ±	13.50	...	Close to 1915 pos 0°	0
4080		III. 737		15 10 15.1	1.863	1	39 12 3.0	13.50	1	vF; vS; stellar	1
4081	3598	III. 139		15 10 24.3	3.505	2	113 31 3.5	13.45	2	cF; S; R; gpmbM	5
4082		II. 758		15 10 56.2	1.606	1::	34 0 29.5	13.45	1::	pF; pS; iR	1
4083	1916		M. 5	15 11 27.2	3.028	1	87 23 8.7	13.39	1	ll; ⊕; vB; L; eCM; st 11..15	11†
4084			R. nova	15 11 39	1.568	::	33 51 55	13.39	::	One of 2, 15' apart np & sf..	0
4085		II. 760		15 11 57.2	1.606	1::	34 6 25.7	13.39	1::	pF; pS; R	1
4086	1917, a		R. nova	15 12 ±	1.552	...	32 10 ±	13.38	...	A ray, vmE, par. to h. 1917	0
4087	1917	II. 759		15 12 11.7	1.552	2	33 10 5.4	13.38	2	cB; vL; vmE 155°0; vg, psbMN.	3†
4088			R. nova	15 12 53	+1.568	::	34 2 31	13.39	::	One of 2, 15' apart np & sf..	0
4089		III. 943		15 13 10.9	-0.870	1	14 7 6.1	13.37	1	vF; vS	1
4090		II. 400		15 13 15.7	+2.681	2	68 36 53.4	13.28	2	vF; S; er	2
4091		III. 944		15 13 35.7	-0.876	1	14 7 4.9	13.33	1	vF; vS	1
4092	{1915 = 3599}	III. 374		15 13 38.5	+3.109	2	92 4 2.5	13.25	2	vF; pL; vLE; r	3
4093											
4094	3600			15 13 51.6	3.297	1	102 34 48.6	13.22	1	B; S; R; glbM; p of 2	1
4095	3601			15 13 56.2	3.299	1	102 39 13.6	13.22	1	F; S; IE; glbM; f of 2	1
4096	3602			15 14 3.2	3.194	1	96 51 8.3	13.21	1	eF; vS; psbM	1
4097	1920			15 14 36.6	2.018	2	43 36 42.0	13.20	2	cF; L; pmE; glbM; * s.	2
4098	1919	I. 148		15 14 59.9	2.975	4	84 25 34.8	13.16	4	cB; cL; iR; vsbM * 12; am st	5
4099	1922	III. 661		15 16 12.3	2.158	(2)	47 50 32.7	13.09	1:	vF; S	2
4100	1921			15 16 13.2	2.156	4	47 46 29.7	13.09	4	cF; pL; vLE; vgbM	4
4101	3603		Δ. 357	15 17 15.2	4.480	3	144 1 39.4	12.98	3	Cl; vL; vRi; lC; st 11..14..	3
4102	3604		Δ. 389	15 17 55.5	4.300	2	140 10 47.2	12.94	2	⊕; cB; L; R; vgbM; rrr; st 15	2
4103	1923	II. 874		15 19 40.8	2.729	1	71 25 32.5	12.85	1	pB; cS; R; psbM; * 7 n.	2
4104	1924			15 21 5.7	2.141	1	47 50 27.1	12.77	1	vF; vS; sp of D neb	1
4105	1925	II. 651		15 21 8.7	2.141	4	47 50 15.1	12.77	4	pF; pS; R; nf of D neb	6
4106		II. 130		15 23 5.1	2.825	2	76 33 15.6	12.62	2	F; pL; iR; vgbM; r	2
4107	{1926 = 3606}	II. 401		15 23 29.0	3.116	2	92 20 47.7	12.59	2	pB; pS; R; vgbM; 3 st f.	3
4108											
4109	3605			15 23 30.4	5.466	1	156 22 34.9	12.53	1	F; S; am st	1
4110	3607			15 25 20.2	4.334	2	140 11 14.9	12.43	2	⊕; cB; pL; R; vglbM; rrr; st 16.	2
4111		II. 906		15 26 3.2	0.808	1	24 45 29.8	12.46	1	F; S; IE 45° ±; vglbM	1
4112		II. 654		15 27 17.7	2.782	1	74 32 25.2	12.34	1	F; pS; E 150° ±	1
4113	1927	II. 178		15 28 2.2	2.777	1	74 20 50.4	12.28	1	pB; cS; p of D neb	4
4114	1927	II. 179		15 28 2.2	2.777	1	74 20 50.4	12.28	1	pB; cS; f of D neb	4
4115		II. 399		15 28 57.7	2.483	1	60 51 21.6	12.22	1	pF; pL; iR; bM; r	1
4116		II. 761		15 29 26.4	1.445	1	33 1 16.3	12.21	1	pF; pS; iF	1
4117	1931	II. 762		15 29 57.4	1.431	(1)::	32 50 58.1	12.17	1	cF; cL; IE	2
4118	1928	II. 96		15 30 6.8	2.746	1	72 55 36.2	12.14	1	pF; pL; iLE; gbM	2
4119	1931, a		R. nova	15 30 31.4	+1.431	::	32 46 22.1	+12.14	::	No description	0

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
4118	1929			15 30 38.0	+2.953	1	83° 33' 9.7	+12.09	1	⊕; vF; vL; R; vgbM; rrr...	1†
4119	1930	III. 634		15 30 48.4	2.172	2	49 46 12.3	12.11	2	vF; S; R; gbM; 2 st 8 f.....	3
4120	3608			15 31 12.7	7.159	1	165 13 11.1	11.97	1	F; pL; R; vgbM	1
4121	3609			15 31 20.1	3.694	1	120 5 47.2	12.04	1	vF; L; R; gbM; r	1
4122		II. 76		15 32 10.2	2.834	1	77 24 9.7	11.99	1	pF; pL; R; rr.....	2
4123	1932			15 33 25.5	2.396	1	57 46 40.6	11.92	1	vF; vS; R; bM	1
4124	1934, a		R. nova?	15 34 29.2	1.214	...	30 6 57.2	11.76	...	R; psbM (by diagram)	0
4125	3610			15 34 53.7	5.014	2	150 46 24.5	11.75	3	!; ○; pF; vS; R; r? am 150 st	3†
4126	1933	II. 655		15 35 0.3	2.759	1	73 45 35.7	11.79	1	F; pS; E 0°.....	2
4127	1934, b		R. nova?	15 35 33.7	1.214	...	30 8 46.2	11.77	...	F; mE	0
4128	1934	II. 764		15 36 18.6	1.214	1	30 11 55.2	11.74	1	cB; S; R; psbM; r.....	2
4129		II. 656		15 36 21.3	2.790	1	75 20 47.7	11.69	1	pB; S; E 135° ±; bM	1
4130		II. 765		15 36 24.1	1.306	1	31 29 46.9	11.73	1	pF; cS.....	1
4131		II. 766		15 36 32.5	1.215	1	30 13 46.6	11.72	1	pB; cL; iE; r	1*
4132	3611		Δ. 552	15 36 53.4	3.904	2	127 19 24.9	11.63	2	!; ⊕; vB; L; R; vgbM; st 13...15.	2
4133	1934, c		R. nova	15 37 18.3	1.214	...	30 14 28.0	11.62	...	(? if not = II. 766).....	0
4134		III. 378		15 38 47.1	1.169	1	29 47 35.8	11.56	1	vF; vS.....	1
4135	1935	II. 425		15 39 16.4	3.019	2	87 8 54.4	11.48	2	vF; vS; R; gbM	5
4136	1936	III. 635		15 39 29.9	2.103	2	48 26 39.7	11.49	2	vF; vS; R; bM; sp of 2.....	3
4137	1937	III. 636		15 39 33.0	2.102	1	48 26 5.4	11.48	1	cF; vS; R; bM; nf of 2.....	2
4138	3613			15 40 38.9	3.334	1	103 19 13.4	11.38	1	eF; S; R; vS * p	1
4139	1938	II. 97		15 40 39.7	2.709	2	71 40 24.0	11.40	2	pF; cS; R; r; bet 2D st.....	4
4140		VII. 29		15 40 56.1	3.664	1	118 10 31.5	11.35	1	Cl; pL; pRi; st vS	1
4141	3612		Δ. 343	15 41 7.4	4.716	1	146 2 39.3	11.31	1	Cl; L; pRi; st 12...14	1
4142	3614			15 41 11.3	3.684	1	118 57 17.9	11.33	1	vF; S; R; sbM	1
4143		III. 371		15 42 3.8	2.459	1	60 54 51.7	11.29	1	vF; S; R.....	1
4144	3615		Δ. 334	15 44 34.9	4.795	3	147 0 57.8	11.06	3	Cl; pS; pRi; mC; st 16	3
4145	1939	II. 583		15 47 11.0	+3.054	1	89 1 54.3	10.91	1	pF; S; E 90° ±; gbM; r	3
4146		III. 313		15 47 35.7	-0.483	2	17 24 55.8	10.96	2	vF; S; E 90° ±; vS * f	2
4147		II. 657		15 47 49.1	+2.772	1	74 59 59.1	10.87	1	F; bet 2 B st	1
4148	1940			15 49 3.7	2.948	1	83 39 30.4	10.78	1	pB; pL; E	1
4149		III. 739		15 49 24.0	0.892	1	27 15 50.0	10.80	1	vF; pL; R; vgbM	1
4150	1941			15 50 19.6	2.946	2	83 35 40.4	10.68	2	!; vF; vS; R; g, smbM	2
4151	1942	III. 646		15 51 5.5	2.742	1	73 42 40.9	10.63	1	vF; S; iE; p of 2	2
4152	1943	III. 73		15 51 8.4	2.741	1	73 37 40.9	10.63	1	eF; vS; iE; f of 2	3
4153	3516		Δ. 304	15 51 50.9	5.056	4	150 6 0.6	10.52	4	Cl; B; vL; pRi; iC; st 7...	4
4154	3617			15 52 18.5	3.846	1	124 8 38.0	10.50	1	F; S; R; gpmbM; Δ of st np	1
4155	3618		Δ. 359	15 56 43.3	4.641	1	143 38 28.8	10.16	1	Cl; S; mC; st 11...14	1
4156	1944	III. 622		15 57 35.3	2.182	2	52 15 33.5	10.15	2	vF; S; R; * 10 sf	4
4157		III. 33		15 58 43.2	+2.658	1	70 16 11.9	10.14	1	eF; (?)	1
4158		II. 873		15 58 43.5	-0.284	1	18 55 6.9	10.13	1	F; R; bM	1
4159	1945			15 58 49.9	+2.901	1	81 31 23.2	10.04	1	* 7 in photosphere	1
4160	1946	III. 637		15 59 36.9	2.065	2	48 55 52.0	10.00	2	pF; vS; R; stellar	3†
4161		III. 140		16 0 51.0	2.629	1	69 4 1.0	9.90	1	vF; vS; r; pB * sf	1
4162	3619		Δ. 360	16 2 19.2	+4.675	2	143 50 42.2	9.74	3	Cl; vB; vL; vRi; iC; st 10...	3
4163		III. 973		16 2 40.3	-3.081	1	10 38 44.0	9.90	1	vF; vS; iE 0°; r	1
4164	1947	III. 553		16 2 48.9	+3.051	1	88 55 4.2	9.74	1	F; L; pmE; vgbM; r.....	2
4165		III. 883		16 3 32.5	-0.266	1	19 13 44.8	9.76	1	eF; vS.....	1
4166	3620			16 3 44.7	+3.922	1	125 52 42.5	9.65	1	pF; R; vglbM; r	1
4167	1948	III. 74		16 4 5.4	+2.713	1::	72 55 51.5	9.65	1	vF; S; r	2*
4168		III. 884		16 5 50.0	-0.147	1	19 59 34.4	9.58	1	vF; vS.....	1
4169	3621			16 6 33.5	+3.866	1	123 53 2.9	9.43	1	eF; S; E; lbM	1
4170	3622		Δ. 326	16 7 16.0	4.936	2	147 32 42.5	9.35	2	Cl; B; L; iC; st 7...10	2
4171		III. 812		16 7 18.1	1.199	1	32 8 30.2	9.44	1	vF; vS; iE	1
4172	1949	III. 889		16 7 18.4	2.293	1	56 35 35.9	9.33	1	vF; S; R; bM	2
417	3624		M. 80	16 8 41.9	+3.567	2	112 37 34.1	9.27	2	!!!; ⊕; vB; L; vmbM (var *); rrr; st 14.	2*
4174		III. 314		16 8 49.3	-0.740	2	17 10 21.4	9.38	2	vF; vS; iE	2
4175	3623		Δ. 68	16 9 54.3	+6.720	3	161 51 51.0	+ 9.10	3	⊕; pF; L; iR; vgbM; rrr; st 14...	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. II.'s Catalogues of Nebulæ.	Sir W. II.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s		° ′ ″	″			
4176	1950	III. 888		16 10 18.4	+2.222	2	57 41 3.1	+9.17	2	vF; S; R; vglbM	3
4177	1951	III. 688		16 11 21.9	2.206	2	53 56 19.0	9.10	2	vF; S; iR	3
4178	1952	II. 151		16 11 58.8	2.911	1	82 15 5.9	9.03	1	F; pL; iE; vgbM; r	2
4179	3625			16 13 42.5	4.589	1	141 36 46.8	8.86	1	Cl; eL; eRi	1
4180	1953	II. 402		16 14 31.8	3.114	1	91 56 56.9	8.83	1	vF; cL; cE 45°±; r	2
4181	1954			16 14 43.9	2.132	1	51 52 57.9	8.83	1	vF; eS; R	1
4182	1955	III. 623		16 14 50.5	2.133	2	51 54 7.9	8.83	2	vF; vS; R; * nf	4
4183			M. 4= B.A.C.5455	16 15 3.6	3.665	...	116 11 11.1	8.77	...	{ Cl; 8 or 10 L st in line, with 5 st; rrr. }	4
4184	3626		Δ. 514	16 16 1.3	4.097	1	130 20 6.7	8.69	1	Cl; B; L; pRi; ICM; st 9...11	2
4185		II. 810		16 16 4.4	1.154	1	32 2 48.5	8.75	1	pF; pS; iE	1
4186		III. 891		16 16 46.1	2.121	1	51 40 47.1	8.67	1	eF; vS; R; lbM	1
4187	3627		Δ. 412	16 17 20.5	4.456	2	138 49 29.1	8.57	2	Cl; eL; pRi; ICM; st 13...15	2
4188	1956	III. 624		16 18 3.7	2.121	2	51 44 50.1	8.57	2	F; S; iR; bM	4
4189	3628		Δ. 536	16 18 14.8	4.037	3	128 30 59.3	8.51	4	B; pL; R; psbM; rr	4
4190		III. 740		16 18 19.3	0.401	1	24 17 36.7	8.59	1	eF; pL; iR	1
4191		III. 892		16 18 23.5	2.144	1	52 26 49.2	8.54	1	eF; S; bM	1
4192		II. 811		16 18 42.4	1.320	1	34 34 35.2	8.54	1	pB; iR; vglbM	1
4193	3629	VI. 10		16 18 43.0	3.657	1	115 43 5.4	8.48	1	Cl; eL; mC; gbM; rrr	2
4194	1957			16 20 20.8	2.009	1	48 44 41.0	8.40	1	F; R; bM	1
4195	1958	III. 638		16 20 27.2	2.010	2	48 47 10.4	8.38	2	eF; vS; R; bM	3
4196	1958, a		R. 2 novæ	16 20 ±	2.010	...	48 47 ±	8.38	...	2 novæ, one eF; one S	0
4197											
4198	1959	III. 639		16 21 10.4	2.025	1	49 13 2.9	8.33	1	vF; vS; R	2
4199	3630			16 21 41.1	7.067	1	162 57 2.8	8.16	1	vF; vS; * 9 nr	1
4200	3631			16 21 49.9	4.654	1	142 18 59.3	8.21	1	Cl; L; IC; st L	1
4201		III. 680		16 22 6.9	1.623	2	39 48 49.8	8.26	2	vF; S; R; lbM; er	2
4202		II. 690		16 22 34.5	1.687	2	41 17 48.6	8.22	2	F; pS; iF; gbM	2
4203	3632			16 22 36.1	5.219	2	150 18 1.9	8.13	2	pF; pL; vIE; gbM	2
4204	1960	II. 652		16 23 0.8	2.004	2	48 45 15.4	8.18	2	eF; pL; R; gbM; r	3
4205		II. 647		16 23 2.7	2.057	2	50 13 17.4	8.18	2	F; S; iF	2
4206	3633			16 23 22.7	4.423	1::	137 48 2.7	8.09	1::	eF; (?); * f nr	1
4207	3634			16 23 39.8	4.426	1	137 51 20.8	8.06	1	F; cS; iE; vglbM; * p	1
4208	1961	II. 875		16 23 50.4	2.053	1	50 8 9.6	8.12	1	pF; S; vIE; vgbmB	2
4209	3635		Δ. 400	16 23 50.9	4.506	1	139 27 48.5	8.05	1	Cl; L; IC; iF	1
4210	3636			16 24 7.3	4.246	1	133 44 18.2	8.04	1	Cl; μ <i>Normæ</i> inv	1
4211	3637	VI. 40	Mechain	16 24 43.0	3.350	1	102 44 17.3	8.01	1	⊕; L; vRi; vmC; R; rrr ...	2
4212	1962	III. 640		16 24 58.0	2.004	1	48 52 16.6	8.02	2	eF; vS; R; bM	3
4213	1962, a		R. nova	16 25 ±	2.004	...	48 52 ±	8.02	...	No description; near h. 1962	0
4214	1963	III. 641		16 25 18.7	2.010	1	49 3 57.0	8.00	1	vF; vS; R	3
4215	1964	III. 890		16 25 34.0	2.203	1	54 38 1.1	7.97	2	vF; pL; iE; rr; * nr	3
4216	3638			16 25 36.3	4.316	2	135 19 19.6	7.92	2	Cl; B; S; st pL	2
4217	1964, a		R. nova	16 25 47.9	2.203	...	54 35 11.1	7.91	...	No description; 4' nf 1964...	0
4218		II. 753		16 26 37.0	2.624	2	69 55 19.4	7.88	3	pB; pL; vIE; pgmbM	2
4219		III. 813		16 26 50.0	1.264	1	34 9 56.7	7.89	1	vF; vS; iR	1
4220	3639			16 26 50.4	6.305	1	159 4 52.8	7.76	1	vF; eS; R; gbM	1
4221	1965			16 28 15.0	2.190	1	54 21 40.8	7.76	1	F; S; R; gbM; * 11 np	1
4222		III. 730		16 28 18.6	2.579	1	68 9 52.2	7.74	1	eF; vS; E	1
4223	3640			16 29 14.5	4.483	1::	138 43 48.6	7.62	1::	l; F; vL; vIE; B * inv	1
4224	3641		Δ. 483	16 30 29.3	4.233	2	133 5 18.9	7.53	2	Cl; eL; pRi; iR; st 11...14..	2+
4225	3642		Δ. 413	16 30 52.1	4.475	1	138 29 3.7	7.49	1	Cl; vL; iRi; IC; rrr; F neb inv.	1
4226	1966	III 893		16 31 43.6	2.058	2	50 41 28.4	7.48	2	vF; S; R; gbM; bet 2 st ...	4
4227	1967			16 32 19.0	2.155	1	53 31 20.9	7.43	1	vF; vS; sbM * 12	1
4228	3643			16 33 50.3	4.420	1	137 11 50.2	7.24	1	Cl; (in M. Way)	1
4229	3644		Δ. 442	16 36 6.6	4.404	2	136 45 36.8	7.06	2	Cl; pRi; eiCM; st 11...12...	2+
4230	1968		M. 13, Halley	16 36 41.2	2.140	3	53 16 19.4	7.08	3	ll; ⊕; eB; vRi; vgcCM; st 11...20.	14+
4231	1969	II. 701		16 38 12.4	2.125	(1)	52 54 7.5	6.95	1	pB; pL; E 45°±; vgmB...	2
4232	3646		Δ. 364	16 38 18.4	4.776	3	143 33 29.1	6.87	3	Cl; L; Ri; ICM; st 9...12 ...	2
4233	3645			16 38 23.8	+7.032	1	162 20 32.0	+6.80	1	vF; pL; vglbM	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by H. and II.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. II.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s		° ' "	"			
4234	1970		{ Σ.5 = Lal. 30510}	16 38 36.0	+2.513	1	65 56 10.0	+6.90	1	{ O; vB; vS; R; disc and border.	1+
4235	3647			16 39 3.5	+5.147	1	148 44 47.7	6.79	1	pF; R; vglbM; *5 p 79 ^s ...	1
4236		I. 280		16 39 16.1	-3.036	3	11 33 21.3	7.01	3	B; cL; lE; slbM	3
4237	3648		Δ. 454	16 39 19.8	+4.308	3	134 28 15.7	6.79	3	Cl; pS; pRi; pC; st 12...15.	3
4238	1971		M. 12	16 39 58.1	3.110	1	91 41 47.4	6.78	2	!; ⊕; vB; vL; iR; gmbM; rrr; st 10...	7
4239	3649			16 40 39.6	5.171	2	148 58 2.1	6.67	2	⊕; pB; cL; R; glbM; rr ...	3
4240	3650		Δ. 456?	16 40 41.4	4.311	1	134 28 33.4	6.68	1	Cl; vL; vRi; lbM; st 12...13	1
4241			D'Arrest, 101	16 41 24	0.72	[2]	28 9 18	6.72	[2]	F; S; R; mbM	0
4242			D'Arrest, 102	16 41 47	0.68	...	27 46 0	6.69	...	F; S; R; makes Δ with 2 S st 12 and 14 m.	0
4243	3651			16 41 51.9	4.170	1	130 58 41.7	6.59	1	Cl; eL; eRi (in M. Way) ...	1
4244		IV. 50		16 43 6.4	1.678	1	42 8 38.8	6.56	1	vB; L; R; Disc + F, r, border	1
4245	3652		Δ. 499	16 44 14.5	4.197	1	131 33 37.7	6.39	1	Cl; B; cL; pRi; st 10...13	1
4246	3653	II. 584		16 45 1.0	3.584	1	111 55 47.5	6.35	1	pB; cL; iR; rrr; st 14...16	2
4247		III. 727		16 45 54.8	1.932	1	48 1 25.6	6.32	1	cF; S; E90°	1*
4248		III. 735		16 46 0.4	1.775	1	44 20 25.3	6.31	1	eF; pS	1
4249	3654		Δ. 520	16 46 3.5	4.112	2	129 16 12.2	6.24	2	Cl; B; L; Ri; st 8...11	3
4250			D'Arrest, 103	16 47 1	0.56	...	26 47 5	6.26	...	F; pS; irr	0
4251	3655			16 47 33.9	4.327	2	134 33 17.6	6.12	2	Cl; pRi; vL; iF; st L & S...	2
4252	3656			16 47 40.8	+4.378	2	135 42 21.0	6.10	2	Cl; L; iRi; iC; st 8...12	2
4253		III. 974		16 47 55.2	-6.988	1	7 9 0.3	6.41	1	cF; S; bM; p of 2	1
4254		III. 975		16 47 58.3	-7.058	1	7 5 59.7	6.39	1	vF; vS; f of 2	1
4255	3657		Δ. 374?	16 48 3.0	+4.734	1	142 28 51.8	6.06	1	Cl; S; Δ ar; st 13	1
4256	{ 1972 = 3659}		M. 10	16 49 47.6	3.159	3	93 52 6.8	5.96	3	!; ⊕; B; vL; R; gvmbM; rrr; st 10...15.	7
4257	1973	III. 689		16 49 47.6	2.121	1	53 16 46.7	5.99	1	eF; cL; E90°	2
4258	3658			16 50 8.6	4.032	2	126 53 32.3	5.91	2	⊕; vF; vL; iR; vgbM; rrr; st 20.	2
4259	1974			16 50 11.5	2.008	1	50 9 37.8	5.96	1	vF (?)	1*
4260	3660		Δ. 456	16 50 37.1	4.326	3	134 26 53.8	5.86	3	!; Cl; B; vL; vRi; st 11...	2
4261	3661		{ M. 62 = Δ. 627}	16 52 18.7	3.810	5	119 53 42.9	5.73	5	!; ⊕; vB; L; gmbM; rrr; st 14...16.	5+
4262		III. 123		16 52 22.2	2.523	1	66 46 53.8	5.76	1	vF; pL; R; lbM	1
4263	3662		Δ. 521	16 52 26.0	4.130	2	129 30 46.3	5.71	2	Cl; B; pL; cRi; st 10	2
4264	{ 1975 = 3663}		M. 19	16 53 59.2	3.701	3	116 3 12.0	5.60	2	⊕; vB; L; R; vCM; rrr; st 16... red.	7
4265	3664		Δ. 556	16 55 16.3	4.067	1	127 40 56.4	5.48	1	Cl; L; pRi; iC; st 9...11	1
4266		III. 124		16 55 22.1	2.521	1	66 46 39.3	5.51	1	vF; stellar	1
4267		III. 728		16 55 52.8	1.534	1	39 51 36.0	5.50	1	vF; cS; iR	1
4268	{ 1976 = 3665}	VI. 11		16 55 55.3	3.661	2	114 33 37.9	5.43	2	⊕; B; L; R; gCM; rrr; st 16.	3
4269	3666	II. 195		16 56 43.5	3.606	1	112 30 9.8	5.36	1	⊕; cB; L; R; gpmCM; rrr; st 16.	3
4270	{ 1977 = 3667}	VI. 12		17 1 28.2	3.716	2	116 23 13.1	4.97	2	⊕; vB; L; R; psbM; rrr; st 16; F neb f.	4
4271	1978			17 1 34.2	3.715	1	116 22 33.5	4.95	1	F; S; vgbM; ⊕ p	1
4272			D'Arrest, 104	17 3 40	0.57	[1]	27 21 54	4.86	[1]	vF; vS; R	0
4273	3668			17 3 51.8	5.588	2	152 39 3.3	4.71	2	F; vL; vL; am st; 2 st inv...	2
4274		IV. 57		17 4 10.8	1.883	2	47 29 24.4	4.78	2	F; stellar	2
4275	3670	I. 147		17 5 40.2	3.803	2	119 17 18.0	4.60	2	⊕; B; cL; R; s, vglbM; rrr; st 16...17.	3
4276	3669			17 5 46.4	5.242	1	149 0 21.5	4.55	1	vF; vS; R; glbM	1
4277			D'Arrest, 105	17 6 3	0.73	[1]	29 3 54	4.59	[1]	pF; vS; R; *13 nr	0
4278			D'Arrest, 106	17 6 15	+0.71	[1]	28 49 54	+4.58	[1]	F; pL; lE	0

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
4279	h. 3671	H. I. 45		h m s 17 7 49.0	s +3.765	2	117° 58' 27.9	+4.43	2	⊕; cB; pS; R; gvmbM; rrr; st 16...17.	4
4280	3672		Δ. 522	17 8 4.2	4.140	1::	129 17 11.4	4.38	1::	Cl; pL; Ri; R; gbM; st 12...14.	1
4281	3673			17 8 45.0	+4.279	2	132 43 29.9	4.33	2	Cl; vL; pRi; IC (* nf taken)	2
4282		III. 945		17 9 14.5	-2.000	1	14 23 21.1	4.47	1	vF; S; E; S * s	1
4283	3676			17 9 30.9	+3.643	1	113 35 51.4	4.28	1	pF; L; R; rr	1
4284	3675			17 9 41.0	4.722	2	141 35 30.9	4.23	2	II; O; pB; vS; R	3+
4285	3674			17 9 57.7	+5.861	2	154 51 25.4	4.18	2	vF; vS; vLE; glbM	2
4286		III. 951		17 10 45.4	-3.497	...	11 12 41.4	4.38	1	eF; S	1
4287	{1979 = 3677}		M. 9	17 10 57.6	+3.507	2	108 22 59.8	4.16	2	⊕; B; L; R; cCM; rrr; st 14	6
4288	3678			17 11 31.9	4.021	2	125 54 52.0	4.10	2	cF; vL; icE; vglbf; * 8 inv.	2
4289	3679			17 11 35.8	3.828	1::	120 0 3.0	4.10	1::	Diffused neb in patches	1
4290	3680			17 12 40.1	4.109	3	128 20 0.0	4.00	3	III; ⊙; eF; S; am St.	3+
4291		II. 812		17 12 40.7	+1.011	1	32 24 7.4	4.08	1	F; S; R; vglbM	1
4292	1980	II. 767		17 12 44.1	-1.062	1	17 32 6.2	4.14	1	cF; pL; R; vgmbM	2
4293		I. 149		17 12 47.8	+3.535	1	109 25 12.0	4.00	1	eB; pS; IE; er	1
4294			M. 92	17 12 56.9	1.840	...	46 43 31.2	4.04	...	⊕; vB; vL; eCM; rrr; st S.	8*
4295	3681	I. 46		17 15 16.2	3.719	1	116 12 43.4	3.78	1	cF; L; R; gbM; rrr	2
4296	3683	I. 48		17 15 23.2	3.491	1	107 40 23.4	3.78	1	⊕; vB; cL; vgvmbM; rrr; st 20.	3
4297	3682			17 15 28.0	3.960	1	124 3 6.8	3.76	1	F; L; E; vglbM; * inv	1
4298			D'Arrest, 107	17 16 23	0.61	[1]	28 4 54	3.77	[1]	pB; S; R; bMN=*12	0
4299	3685			17 16 28.0	3.827	1	119 51 37.1	3.67	1	Neb in patches (M. Way)	1
4300	3684		Δ. 225	17 17 23.3	6.165	2	156 55 47.9	3.53	2	⊕; cB; L; R; vgmbM; rrr; st 14...17.	2
4301			Auw. N. 36	17 20 19.2	3.184	...	94 57 2.0	3.37	...	F; L; vlbM (Winnecke, April 12, 1860).	0
4302	{1981 = 3686}	IV. 11		17 20 50.4	3.650	2	113 37 47.3	3.21	3	II; ⊙; pB; S; R	5*+
4303		III. 137		17 22 2.6	2.412	1	63 25 22.2	3.24	1	vF; pL; iF	1
4304	3687			17 22 58.1	3.913	1	122 28 48.3	3.11	1	Cl; S; P; B * inv	1
4305	3688			17 24 47.4	4.142	1	128 58 1.5	2.95	1	eF; pS; IE; * 9 att	2+
4306	3689			17 25 34.1	3.915	3	122 28 38.7	2.89	3	Cl; st 6.7, 13	2
4307	3690		Δ. 457	17 26 3.2	4.380	2	134 38 32.2	2.84	2	⊕; vB; L; R; pg, psvmbM; rrr; st 17...	2
4308		II. 901		17 26 3.6	2.680	1	73 29 2.4	2.88	1	F; S; iF; er	1
4309	3691			17 27 57.9	6.657	2	159 41 19.0	2.60	2	cF; S; R; glbM; * 13 sp	2
4310	3693			17 28 51.0	3.998	1	124 54 49.0	2.60	1	Cl; pL; iRi; IC	1
4311	3692		Δ. 366	17 29 17.3	4.871	2	143 35 5.2	2.54	2	⊕; B; vL; Ri; st 13	3
4312	3694			17 29 48.6	5.527	1	151 36 5.1	2.47	1	eF; S; R; p of 2	1
4313	3696		Δ. 568	17 30 1.9	4.067	1	126 51 8.0	2.50	1	Cl; pL; pRi; iR; st 9...10...	1
4314	{1982 = 3697 1983}	I. 44		17 30 6.9	3.658	4	113 48 55.0	2.50	4	pB; pL; R; * 12 finv	6
4315	{3698 = 3698}		M. 14	17 30 16.0	3.146	2	93 9 25.0	2.50	2	⊕; B; vL; R; eRi; vgmbM; rrr; st 15...16.	7
4316	3695			17 30 18.8	5.528	1::	151 36 3.9	2.43	1	eeF; f of 2	1
4317	4020	h. o. n.		17 30 26.3	3.939	1	123 9 16.8	2.46	1	Cl; F; L; pRi; IC; st 13...15	1
4318	3699		M. 6	17 30 55.8	3.905	1	122 7 4.9	2.43	1	Cl; L; iR; IC; st 7, 10...	4
4319	3700			17 32 15.0	5.436	1	150 39 39.1	2.27	1	eF; S; R; 3 st nr	2
4320			D'Arrest, 108	17 33 56	+0.69	[2]	29 6 12	2.26	[2]	vS; gbM	0
4321		VI. 41		17 34 7.6	-2.177	1	14 11 11.9	2.33	1	⊕; cL; R; vglbM; rr	1
4322	3701'			17 35 0.1	+4.002	1::	124 56 32.1	2.07	1	Nebulous portion of M. Way	1
4323	3702		Δ. 612	17 35 9.3	3.912	2	122 17 0.5	2.05	2	Cl; vL; Ri; IC	2
4324	3702'			17 36 30.1	3.957	1	123 37 33.2	1.94	1	Cl; vL; pRi; st 8...12	1
4325		II. 587		17 37 53.6	+2.997	1	86 44 25.5	+1.85	1	F; cL; iF	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
4326	3703			17 37 56.4	+3.887	1	121° 27' 58".6	+1.82	1	Cl; pS; lRi; lC; st 10...12 ..	1
4327	1984			17 38 46.0	+3.689	1	114 49 51.5	1.75	1	Cl; st vS	1
4328	1987	III. 741		17 39 11.8	-1.069	1	17 49 22.7	1.69	1	vF; vS; R; stellar; *8 s.....	2
4329	3704			17 39 39.5	+4.019	1::	125 22 49.8	1.66	1::	Cl; F; eL; vS st + neb	1
4330	3701			17 39 43.4	19.744	1	175 24 24.0	1.20	1	pB; R; vgbM	1
4331	1985	I. 150		17 40 32.5	3.565	1	110 18 21.0	1.60	1	pB; pL; R; bM	3
4332	3705		Δ. 557	17 40 41.6	4.077	2	127 0 0.1	1.57	2	⊕; vB; pL; R; vgmbM; rrr; st 18...20.	2
4333	1986	II. 586		17 40 54.7	3.557	2	109 58 3.8	1.56	2	pB; pS; R; gbM; r; *15 np	3
4334	3706		Δ. 597?	17 40 55.0	3.999	1::	124 48 46.5	1.55	1::	Cl; vL; vRi; st 12...13	1
4335	3707	VI. 13		17 41 40.5	3.847	2	120 10 12.7	1.49	2	Cl; pL; pRi; bifid; st 12... ..	3+
4336	3708			17 41 59.7	3.991	1::	124 34 43.8	1.46	1::	eL; iR; pmbM; r	1
4337	3709			17 42 37.5	4.018	1::	125 20 42.0	1.40	1::	Cl; rr; st eS + neb	1
4338	1988			17 44 16.8	3.705	1	115 21 32.1	1.27	1	eF; S; (?)	1
4339	3711			17 44 29.7	3.620	1	112 18 23.5	1.25	1	Cl; pRi (in M. Way)	1
4340	3710		M.7 (Lacaille)	17 44 39.6	3.999	1	124 46 37.9	1.23	1	Cl; vB; pRi; lC; st 7...12 ...	3
4341	3712			17 44 54.5	3.815	1::	119 5 36.3	1.21	1::	Neb or nebulous patch of M. Way.	1
4342	3713'			17 45 36.6	3.855	1	120 23 56.5	1.15	1	Neb or nebulous patch of M. Way.	1+
4343	1989			17 45 57.6	2.502	1	66 53 15.5	1.15	1	!; vF; S; R; vsymbMvSRN.	1+
4344	3713			17 46 1.3	5.767	1	153 38 2.5	1.05	1	F; S; E; bM; bet 2 st 10 ...	1
4345	3714			17 48 31.2	6.132	1	156 24 42.9	0.83	1	pF; S; pmE 90°; *12 f, att...	1
4346	1990		M. 23	17 48 40.9	3.532	3	108 59 43.7	0.89	3	Cl; B; vL; pRi; lC; st 9...10, 11...13.	6
4347	3715		Δ. 460?	17 48 51.4	4.372	2	134 13 52.2	0.84	2	Neb + Cl; pL; mE; gvlbM ...	2
4348		III. 957		17 49 25.1	2.631	1	71 42 59.5	0.85	1	vF; vS; p of 2	1
4349		III. 958		17 49 31.1	2.629	1	71 38 58.5	0.85	1	vF; vS; f of 2	1
4350	3716			17 50 20.0	+5.991	2	155 23 58.1	0.67	2	vF; vS; f * of * inv	2
4351			Auw. N. 37	17 50 54.1	-0.638	...	19 49 12.3	0.81	...	pF; L; mE (Auwers, July 22, 1854).	0
4352	3717			17 51 10.6	+3.685	1	114 38 24.1	0.67	1	Cl; Ri; eL; vLc	1
4353		VIII. 53		17 51 30.0	3.491	1	107 23 18.2	0.64	1	Cl; pS; lRi; lC	1
4354			D'Arrest, 109	17 53 43	0.49	[2]	27 20 24	0.53	[2]	vF; R; 1st of 3	0
4355	{ 1991 = 3718 }	{ IV. 41 V. 10, 11, 12 }	M. 20	17 53 51.8	3.640	3	113 1 39.9	0.43	3	!!!; vB; vL; trifid; * inv ...	8 Mon.†
4356			D'Arrest, 110	17 54 9	0.48	[2]	27 18 30	0.50	[2]	vF; vS; 2nd of 3	0
4357	3719	II. 199		17 54 13.9	3.282	2	98 56 37.3	0.41	2	pB; pL; R; rr	3
4358	3721	VII. 7		17 54 36.4	3.779	1	117 53 32.8	0.36	1	Cl; pS; Ri; lC; st 9...13 ...	3
4359	3720	I. 49		17 54 37.2	3.845	2	120 1 29.8	0.36	2	⊕; B; pL; R; gvmbM; rrr; st 16...17.	3
4360			D'Arrest, 111	17 54 38	0.49	[2]	27 22 0	0.46	2	F; pL; 3rd of 3	0
4361	3722		M. 8	17 55 17.9	3.677	3	114 21 15.3	0.31	3	!!!; vB; eL; eiF; with LCl...	8 Mon.†
4362	1992			17 55 27.9	2.811	1	78 57 6.6	0.32	1		1
4363		V. 9		17 55 39.8	3.652	1	113 27 28.4	0.28	1	F; L; eE	1
4364	3723	II. 200		17 55 51.7	3.846	2	120 3 24.5	0.25	2	⊕; pF; cS; R; gbM; rrr; st 16...17.	3*
4365	3724		Δ. 569	17 56 1.6	4.054	1::	126 18 6.9	0.23	1::	Cl (in M. Way)	1
4366	3725			17 56 6.0	3.677	2	114 19 58.9	0.23	2	Cl; B; L; pRi; vL neb p ...	2
4367	1993		M. 21	17 56 13.8	3.626	1	112 30 8.6	0.22	1	Cl; pRi; lC; st 9...12	2
4368		V. 13		17 56 32.3	3.692	1	114 53 23.7	0.19	1	eL; eiF; st f	1*
4369			Auw. N. 38	17 56 38.5	3.076	...	90 17 44.9	0.20	...	pF; vS; vS neb * p (Hind, Ap. 1852).	0
4370			Auw. N. 39	17 57 15.0	3.247	...	97 34 58.4	0.15	...	(Brorsen, 1856.) No description.	0
4371		II. 198		17 57 27.3	3.778	1	117 49 20.6	0.12	1	pF; S; iE; er or Cl	1
4372	3726		Δ. 473	17 57 50.9	+4.350	2	133 43 25.8	0.06	2	⊕; B; R; eC; gbM; rrr; st 15...16.	2*
4373		IV. 37		17 58 20.0	-0.023	1	23 22 9.5	+0.15	1	○; vB; pS; sbMvSN	1
4374	1994	II. 197		17 58 42.4	+3.696	1	115 0 52.0	0.00	1	eF; pL; iR; r	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	N of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s						
4375	3727			17 58 45.8	+5.788	1	153 47 28.2	-0.06	1	eeF; ceS; R	1†
4376	3729			17 58 47.5	3.646	1::	113 14 0.0	0.00	1::	Cl; vL; vRi	1
4377		III. 555		17 59 39.1	2.624	1	71 27 7.8	-0.04	1	ceF; S; IE; r	1
4378	3730	IV. 12		18 0 44.2	3.722	1	115 56 17.6	0.18	1	⊕; F; L; IE; vglbM; rr; st 20.	3
4379	1995			18 0 50.8	3.518	1	108 26 46.6	0.18	1	Cl; pRi; vIC; st L & S	1
4380	3732			18 1 9.8	3.769	1	117 32 31.7	0.21	1	F; vL; cE; lbM; rr	1
4381	3728			18 1 11.4	8.690	1	166 36 39.2	0.36	1	vF; vS; R; glbM	1
4382	3731			18 1 12.2	3.900	3	121 46 50.4	0.22	3	⊕; pB; pL; R; glbM; rrr; st 16	3
4383		II. 902		18 1 18.7	2.649	1	72 25 54.3	0.19	1	F; L; R; vglbM	1
4384	{ 1996 = 3733 }			18 1 23.9	3.671	2	114 7 30.1	0.23	2	vF; vL; IE; * inv	2
4385	1997	VIII. 54		18 2 24.7	3.476	1	106 48 57.7	0.31	1	Cl; L; IC; st cL	3
4386	3734			18 2 47.5	3.970	1	123 53 29.2	0.36	2	○; F; L; cE; hazy border...	2
4387			D'Arrest, 112	18 3 51	1.34	[1]	37 44 0	0.38	[1]	ceF; vS; R; * 16 nr.....	0
4388	{ 1998 = 3735 }	VII. 30		18 4 20.6	3.602	2	111 37 33.6	0.48	2	Cl; vL; IC	3
4389	3736	II. 201	Δ. 619	18 4 33.3	3.902	3	121 51 0.7	0.51	3	⊕; cB; L; R; rrr; st 15.....	5
4390	2000		Σ. 6	18 5 17.8	2.912	1	83 10 53.5	0.55	2	○; vB; vS; R; l hazy	3*
4391	1999			18 5 23.5	3.617	1:	112 10 47.6	0.58	1	Cl; st vS	1
4392	{ 2001 = 3739 }	VII. 31		18 7 24.1	3.616	2	112 10 21.5	0.75	2	Cl; pRi; pC; cE; st 13	3
4393	3737		Δ. 376	18 7 24.6	4.797	2	142 15 13.3	0.79	2	⊕; cB; cL; R; gmbM; rrr; st 15	2
4394	3738			18 7 56.9	5.793	1	153 51 18.2	0.86	1	ceF; S; *6, sp	1
4395	2002			18 8 47.2	3.529	1	109 55 8.6	0.88	3	F; pL; cE; * inv	3†
4396	2003	VIII. 55		18 9 28.2	3.473	1	106 41 22.1	0.93	1	Cl; IC	2
4397	2004		M. 24	18 10 13.7	3.518	2	108 28 7.3	0.99	2	l; Cl; vRi; vmC; R; st 15 (M. Way).	4*
4398	3740	VIII. 15		18 10 14.2	3.363	1	102 17 10.3	0.99	1	Cl; lRi; IC	2
4399	2005			18 10 20.9	3.429	1	104 59 37.7	1.01	1	Cl; lRi; IC; st 10...12	1
4400	2006		M. 16	18 10 57.0	3.401	1::	103 50 2.2	1.06	1	Cl; at least 100 st L & S ...	3
4401	2007		M. 18	18 11 44.6	3.485	1::	107 11 8.1	1.13	1	Cl; P; vIC	4
4402	3741			18 11 45.3	5.724	2	153 17 47.0	1.20	2	vF; S; R; gvlbM; *9 p.....	2
4403	2008		M. 17	18 12 33.1	3.460	2	106 13 36.0	1.20	2	III; B; eL; eiF; 2-hooked...	9†
4404	3742	I. 50		18 14 41.9	3.855	3	120 25 26.0	1.40	3	⊕; vB; pL; R; rrr; st 16 ...	4
4405	2009			18 15 20.7	3.358	1::	102 5 58.8	1.44	1	IC; lRi; IC; st 11...12	1
4406	{ 2010 = 3743 }		M. 28	18 15 55.4	3.692	2	114 56 30.0	1.50	2	l; ⊕; vB; L; R; geCM; rrr; st 14...16.	8
4407	3744	II. 204		18 17 13.4	3.645	1	113 16 30.4	1.62	1	○ or ⊕; pB; eeS; R	2
4408	3745			18 19 12.0	5.726	1	153 22 22.5	1.85	1	pF; S; R; gbM	1
4409	3746			18 19 23.4	3.358	1	102 6 42.3	1.79	1	Cl; pL; pRi; st 12...15	1
4410		VIII. 72	C. H.	18 20 43.1	2.921	1	83 31 15.3	1.89	1	Cl; IC; st L	3
5076				18 20 44.6		...	123 30 27.3		...	See No. 5076	0
4411			M. 69=Δ. 613	18 22 13.2	3.917	3	122 26 33.5	2.05	3	⊕; B; L; R; rrr; st 14...16	4*
4412	3748	I. 51		18 22 16.6	3.708	1	115 34 55.5	2.05	1	⊕; B; S; R; rr	2
4413	2011			18 23 4.6	3.385	1	103 15 12.7	2.11	1	Cl (in M. Way)	1
4414	{ 2012 = 3749 }	II. 205		18 23 24.2	+3.651	3	113 33 57.5	2.15	3	⊕; pB; pL; iR; gpmbM; rrr; st 16.	4
4415			Auw. N. 40	18 23 35.4	-1.719	...	15 29 47.7	2.01	...	II; pB; pL; E 50°; 2 st p Var. (Tuttle.)	0*
4416	2013	VI. 23		18 24 31.5	+3.477	1	106 59 3.8	2.24	1	Cl; pL; vRi; pC; st 11...15	2
4417		II. 907		18 24 55.8	1.967	1	50 14 52.8	2.24	1	F; S; iF	1
4418	2014	VIII. 14		18 25 4.6	3.488	1::	107 26 2.3	2.29	1	Cl; L; Ri; IC; st vS	2
4419			Auw. N. 41	18 25 4.9	0.230	...	25 5 44.2	2.20	...	S; pmE; * inv (Σ. neb No. 7)	0
4420	3751			18 25 44.1	3.317	1	100 29 32.8	2.34	1	Cl; P; IC; pS; st 9-10, 12...13	1
4421	3752		Δ. 607	18 26 32.0	+3.936	4	123 5 5.1	-2.43	4	B; S; IE; rrr; st 15	5

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
4422	3750			18 27 8.1	+7.511	1	163° 22' 49.3	-2.59	1	vF; S; lE; glbM.....	1
4423			Auw. N. 42	18 27 19.3	3.210	...	96 4 46.7	2.45	...	pF; vS; E (Winnecke, June 1855).	0
4424	{ 2015 = 3753 }		M. 22	18 27 52.1	3.662	4	114 0 25.8	2.54	4	ll; ⊕; vB; vL; R; vRi; vmC; st 11...15.	10
4425	2016			18 28 5.5	2.496	1	66 32 22.1	2.53	1	Cl; P; IC.....	1
4426	3754	VIII. 12		18 29 5.1	3.266	1	98 20 0.1	2.63	1	Cl; L; pRi; vIC.....	2
4427	3755			18 32 2.0	5.603	2	152 25 36.5	2.95	2	pF; S; R; pSBM; r.....	2
4428	3756		M. 70 = Δ. 614	18 34 4.9	3.910	3	122 25 12.6	3.08	3	⊕; B; pL; R; gbM; st 14...17	4*
4429	2017			18 34 9.9	3.185	1	94 53 29.9	3.07	1	Cl; L; Ri; st 10...18	1
4430	2018			18 34 41.1	3.219	1	96 21 12.4	3.12	1	Cl; vRi; vIC (in M. Way)...	1
4431	3757			18 35 9.5	5.941	2	155 19 18.8	3.24	2	vB; pL; R; vg, psmbM; *7p	2
4432	3758		M. 26	18 37 32.6	3.293	1	99 31 57.9	3.37	1	Cl; cL; pRi; pC; st 12...15.	6
4433		VI. 15		18 39 37.8	3.715	1	116 3 40.2	3.56	1	Cl (?) ; vF; cL	1
4434	3759			18 40 6.8	5.131	1	147 27 46.8	3.64	1	pF; pS; lE 90°; psbM	1
4435			Auw. N. 43	18 43 18.8	3.192	...	95 22 6.4	3.86	...	Cl; B; 60 st 13 (Winnecke, 1854).	0
4436	3760			18 43 36.6	5.679	1	153 19 39.2	3.96	1	Neb. No description.....	1
4437	2019		M. 11	18 43 37.2	3.219	2	96 26 7.6	3.88	2	l; Cl; vB; L; iR; Ri; st 1 L, 11..	8+
4438	3761			18 44 2.1	4.873	2	143 58 46.9	3.97	2	F; S; vIE; gbM	3
4439	4021	h. o. n.		18 44 18.1	4.866	2	143 53 19.3	3.99	2	pF; S; R; glMbM; last of gr	2
4440	2020			18 44 52.5	2.836	1	79 49 0.6	3.98	1	Cl; pRi; IC; iF	1
4441	3762	I. 47		18 45 29.2	3.276	1	98 52 8.5	4.05	1	⊕; pB; vL; ir; vglbM; rrr	2
4442	3763		M. 54 = Δ. 624	18 46 6.2	3.846	3	120 38 29.7	4.11	3	⊕; vB; L; R; g, smbM; rrr; st 15.	6
4443	2021			18 46 17.5	3.549	1	110 4 8.1	4.13	1	Cl; pRi; st 9...13	1
4444	{ 2022 = 3766 }	III. 143		18 46 39.8	3.623	2	112 52 17.2	4.16	2	F; S; rr Cl + neb	3
4445	3764			18 47 25.9	6.042	2	156 17 14.0	4.30	2	vF; S; R; glbM; *9 sp	2
4446	3765			18 48 2.6	6.430	1	158 47 1.2	4.36	2	vF; pL; R; vglbM	2
4447	2023		{ M. 57 D'Arquier }	18 48 20.1	2.228	2	57 8 57.2	4.26	3	{ III; ⊙; B; pL; cE (in } Lyra).	14+
4448	3767			18 48 51.4	5.153	1	147 56 49.3	4.39	1	pF; cS; R; vmbM	1
4449	3768			18 49 59.0	5.874	1	155 5 21.7	4.51	1	pF; S; E; glbM; 2 st 8 p	1
4450	3770		Δ. 573	18 50 5.5	4.047	1	126 48 45.9	4.47	1	⊕; vL; vIE; vgbM; rrr; st 14...16.	1
4451	2024			18 50 25.1	2.838	1:	79 49 16.2	4.46	1	Cluster.....	1
4452	3769			18 50 26.1	4.871	1:	144 7 32.7	4.51	1	eF; pL; R	1
4453		VIII. 13		18 52 20.9	3.281	1	99 7 33.8	4.64	1	Cl; vL; P	1
4454	3771			18 52 32.0	6.472	1	159 6 45.8	4.74	1	vF; S; R; pmbM; *7, 8 nf	1
4455	3772			18 53 15.0	5.552	1	152 23 37.6	4.78	1	eeF; vglbM; v difficult	1
4456	3773			18 53 29.2	5.938	1	155 39 40.7	4.81	1	vF; S; R; glbM; p of 2	1
4457	2025			18 53 33.1	3.086	1	90 39 5.1	4.73	1	Cl; vL; P; st 12...	1
4458	3774			18 53 44.6	5.934	1	155 38 10.4	4.82	1	eF; S; R; glbM; f of 2	1
4459	2026			18 53 59.2	3.512	1	108 44 8.6	4.78	1	Cl; pL; pRi; R; st 12...15...	1
4460	2027			18 54 50.0	2.810	1	78 35 18.1	4.83	1	Cl; P; IC	1
4461	3775			18 55 7.2	5.463	2	151 34 23.8	4.94	2	eF; vS; cE; psbM; 3 st p	2
4462		III. 742	D'Arrest, 113	18 55 32.0	1.618	1	41 45 14.2	4.86	1	vF; stellar	1*
4463	2028			18 55 58.2	2.351	1	60 55 35.4	4.92	1	Cl; pL; P; st 11...12	1
4464	3776		Δ. 262	18 56 26.8	5.730	2	154 3 58.5	5.05	2	cB; cL; R; vg, psmbM; r...	2
4465	3777			18 57 31.7	5.518	1	152 10 36.1	5.13	1	eF; cS; R; glbM	1
4466	2029			18 57 56.6	3.035	1	88 25 11.0	5.10	1	Cl; L; IC; st L & S	1
4467	3778		Δ. 295	18 58 28.0	5.324	5	150 11 49.7	5.21	5	⊕; B; vL; iR; rrr; st 11...16	5
4468	3779			18 59 37.5	5.085	1	147 15 30.0	5.30	1	pB; pL; R; gbM	1
4469	3780			19 0 33.6	4.653	2	140 51 32.9	5.37	2	pF; pL; mE 63°; vglbM	2
4470	2030	VII. 19		19 0 50.5	2.981	3	85 59 17.5	5.35	3	Cl; vL; vRi; pC; st 12...14.	5
4471	2031	VII. 62		19 1 46.5	2.971	2	85 32 43.1	5.43	1	Cl; S; Ri; IC; st 11...12	5
4472	3781			19 2 15.9	5.014	1	146 32 24.1	5.53	1	pB; S; R	1
4473			Auw. N. 44	19 4 4.8	+3.051	Au	89 11 51	-5.62	Au	ll; pB; pL; gbM; Var.? (Hind)	0*

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
4474	3782			19 4 14.3	+4.647	2	140° 53' 11".6	-5.68	2	vF; pS; iR	2
4475	3786			19 6 50.4	4.160	1	130 25 44.6	5.88	1	vF; S; R; pslbM	1
4476	3783			19 6 53.8	5.349	1	150 44 16.4	5.92	1	vF; S; R; lbM; 1st of 3.....	1
4477	3784			19 7 6.3	5.349	1	150 44 46.1	5.93	1	eF; vS; 2nd of 3.....	1
4478	3785			19 7 7.3	5.351	1	150 46 25.8	5.94	1	eF; S; 3rd of 3	1
4479	2032	IV. 14		19 7 14.8	3.137	2	92 57 5.3	5.89	2	vF; L; R; vvlbM; r	3
4480	2033			19 8 3.7	2.968	1	85 22 9.5	5.95	1	Cl; P; IC.....	1
4481	2034			19 8 38.7	3.449	1	106 30 34.4	6.02	1	Cl; vL; IC	2
4482	2035			19 9 37.9	3.097	1	91 9 48.3	6.09	1	Cl; P; IC; st 10...11	1
4483	3787			19 10 22.9	5.690	1	154 8 23.1	6.23	1	pB; S; R; pgbM	1
4484			Lac. I. 13	19 10 45.5	6.917	...	161 45 35.7	6.21	...	Neb without st (Lac. Auw. 40)	0
4485	2036		M. 56	19 11 7.2	2.339	4	60 3 41.6	6.18	6	⊕; B; L; iR; gvmCM; rrr; st 11...14.	14
4486	3788			19 11 17.6	4.954	1	146 1 45.6	6.28	1	vF; L; R; vglbM	1
4487	2037	III. 743		19 11 37.3	2.931	1	83 42 46.5	6.25	1	○; F; L; R; vsbM disc; S* nf	3*
4488	3789			19 11 39.2	5.283	3	150 10 43.4	6.32	3	cF; cS; R; lbM; *9 s	3
4489	3790			19 12 48.3	5.896	2	155 52 59.1	6.43	2	eeeF; pS; am S st	3
4490	2038			19 13 23.1	3.102	1	91 21 22.0	6.40	1	eS; stellar	1
4491	3791			19 14 30.8	4.890	2	145 13 36.8	6.54	2	pB; S; mE; pslbM	2
4492			Auw. N. 45	19 15 55.9	2.098	...	52 28 32.1	6.53	...	vF (Winnecke, Dec. 1853)	0
4493	2039	VIII. 81		19 17 13.9	2.565	3	68 6 5.0	6.70	3	Cl; P; IC.....	4
4494	3792			19 18 23.8	4.094	1	129 9 44.8	6.84	1	eF; pS; R; vglbM	1
4495	2040			19 19 6.7	3.000	1	86 44 46.9	6.87	1	Cl; Ri; bet 2 st 9	1
4496	3793			19 20 47.4	4.941	2	146 11 39.2	7.06	2	eF; vS; R; lbM; 3 vS st nr	2
4497	2041	VIII. 21		19 21 22.5	2.491	1	65 8 23.8	7.04	1	Cl; vL; pRi; vIC; st 10...	3
4498	2042	VI. 14		19 24 29.7	2.628	3	70 1 4.0	7.30	3	Cl; L; vC; E 0°; st 14...18	5
4499	2043	VI. 38		19 24 53.0	2.875	4	81 3 37.8	7.34	4	eB; S; L; R; rrr	5
4500	3796			19 27 18.0	4.068	1	128 51 32.9	7.57	1	eF; R; vgbM	1
4501	3795			19 27 18.9	4.209	1	132 36 18.9	7.57	1	eF; vS; * 14 att	1
4502	3794			19 28 30.3	6.638	2	160 57 27.8	7.74	2	pB; E; biN; * 8 f	2
4503	3798		M. 55 = Δ. 620	19 31 7.9	3.817	2	121 15 44.2	7.86	2	{ ⊕; pB; L; R; vRi; } vgbM; st 12...15.	5
4504	3797			19 31 43.0	5.109	2	148 58 44.5	7.95	2	pS; R; vgbM	2
4505	2044			19 33 58.6	1.790	2	43 44 48.8	8.04	2	Cl; L; pRi; IC; st 11...14...	2
4506	3799			19 34 7.2	4.858	1	145 40 17.8	8.14	1	pB; pS; pmE; glbM	1
4507	2045	III. 744		19 35 0.0	3.300	...	100 38 29.9	8.17	...	pF; pL; R; bM; r	2
4508	2046			19 35 6.9	2.462	1	63 30 55.5	8.15	1	Cl; vL; pRi; IC; st 10...15	1
4509	3800			19 35 16.0	3.744	1	118 52 41.0	8.20	1	eF; pS; R; vlbM; * np.....	1
4510	2047	IV. 51		19 36 3.0	3.386	3	104 28 52.5	8.25	3	○; B; vS; R	8†
4511	2048		Harding	19 36 24.3	2.053	1	50 7 59.5	8.25	1	Cl; vL; vRi; st 11...15 (Harding, 1827).	1
4512	2049	VII. 18		19 37 13.6	2.555	1	67 1 48.4	8.32	1	Cl; cRi; E; st 11...12	2
4513		II. 878		19 40 47.3	1.298	1	34 18 29.2	8.56	1	pB; iF; bM	1
4514	2050	IV. 73		19 41 7.5	1.622	1	39 49 41.7	8.61	1	○; B; pL; R; * 11 M	2†
4515	2051	VIII. 73		19 43 30.0	2.912	1	82 26 25.1	8.83	2	Cl; P; IC	3
4516	2052	VII. 9		19 45 4.3	2.569	1	67 15 45.5	8.95	1	Cl; L; pRi; pC; st 11...12 ..	2
4517	2053			19 45 43.7	1.072	1	30 55 55.2	8.96	1	Cl; vL; IC; st 7, ...	1
4518	2054	VIII. 16		19 46 33.8	2.408	1	60 57 6.5	9.05	1	Cl; P; IC; st 11...12	2
4519	2055	VIII. 18		19 46 53.0	2.832	1	78 40 27.3	9.09	1	Cl; S; P	2
4520	2056		M. 71	19 47 28.8	2.674	3	71 34 55.1	9.13	3	Cl; vL; vRi; pmC; st 11...16	13
4521	2057	VI. 16?		19 48 13.9	2.696	1	72 28 24.0	9.20	1	Cl; vS; vC	3
4522	2058	VIII. 19		19 48 41.7	2.824	1	78 15 35.1	9.23	1	Cl; P; IC.....	3
4523	3802			19 48 57.9	3.818	1	122 11 30.6	9.28	1	vF; S; R; psbM	1
4524	2059			19 49 31.7	2.823	1	78 12 33.0	9.30	1	Cl; S; P	1
4525	3801			19 49 42.9	5.672	1	155 36 45.6	9.38	1	eF; vS; R; psbM; * 11 up...	1
4526	3803			19 50 49.6	4.355	1	137 27 28.8	9.44	1	vF; S; vIE; glbM	2
4527		II. 202		19 51 22.9	2.417	1	61 1 41.1	9.43	1	Neb; r	1
4528	3804			19 51 33.2	4.845	2	146 28 18.7	9.51	2	eF; cL; R; vglbM; 2 st f ...	2
4529	3805			19 51 47.6	4.079	1	130 35 26.7	9.51	1	pB; S; R; vS * np	1
4530	3806			19 52 27.6	4.760	1	145 13 47.9	9.57	1	vF; S; R; bM	1
4531	3807			19 53 19.4	+4.402	1	138 39 10.1	-9.63	1	pF; S; vIE; psbM	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos	Other Authorities.								
4532	h. 2060	H.	M. 27	19 53 29.3	+2.588	4	67° 39' 43.0	- 9.60	3	!!!; vB; vL; BiN; iE (Dumb-bell N).	13+
4533	3808			19 54 40.9	4.723	2	144 45 44.5	9.75	2	F; S; vLE; glbM.....	2
4534	3809			19 55 31.7	4.849	1	146 47 1.7	9.81	1	pF; S; R.....	1
4535	2063			19 56 8.8	1.354	1	34 15 9.9	9.77	1	Cl; pS; pmC; iR; st 12...16	1
4536	2062	III. 144		19 56 23.1	2.309	3	56 50 48.7	9.81	3	F; am M. Way st	4*
4537	2061			19 56 23.6	2.847	1	79 7 24.4	9.82	1	Cl; cL; E; pRi; st 13... ..	1
4538			Auw. N. 46	19 56 39.0	3.068	...	89 56 49.0	9.85	...	F; *10 p 1 ^s , s 1' 29" (Bond, Nov. 1852).	0
4539	3810			19 56 41.0	5.207	1	151 29 30.7	9.91	1	F; pS; gbM	1
4540	3811		Δ. 425	19 57 5.0	4.395	2	138 46 5.7	9.91	2	B; S; cE; gpmbM	2
4541	3812			19 57 39.9	4.849	1	146 47 18.2	9.96	1	F; S; iE; glbM	1
4542	2065			19 57 46.5	3.151	1	93 57 0.8	9.94	1	Cl; S; vmC; st 19	1
4543	2064		M. 75	19 57 49.1	3.547	3	112 18 47.5	9.95	3	⊕; B; pL; R; vmbMBN; rr	10
4544	2066	VII. 59		19 59 11.2	1.967	1	46 23 49.7	10.01	1	Cl; L; vRi; cC	2
4545	3813			19 59 31.7	4.733	1	145 11 4.4	10.12	1	eeF; L; pmE	1
4546	3814			19 59 40.6	4.387	2	138 46 26.4	10.12	2	vB; S; R; pgvmbM	2
4547	3815			19 59 57.7	4.382	2	138 41 20.8	10.14	2	cF; cS; E 90°; gbM	2
4548	2067		Σ. 2630	20 0 38.3	2.258	1	54 37 13.1	10.13	1	Cl; st L & S; * inv	1
4549	3816			20 2 2.9	6.426	2	161 11 51.8	10.34	2	F; pS; iE; glbM; *9 p 10 ^s .5; 1st of 4.	2
4550	2068		Σ. 2631	20 2 4.8	2.636	1	69 17 54.5	10.25	1	Cl; iC; st 10...13; * inv.....	1
4551	2069	VIII. 86		20 2 44.4	2.179	1	52 9 34.3	10.29	1	Cl; P; iC	2
4552	3819			20 3 15.0	4.278	1	136 34 21.6	10.38	1	F; vS; R; vgmbM; *7 nf...	1
4553	3817			20 3 27.6	6.429	2	161 17 6.5	10.45	2	pB; S; R; eS*sf; 2nd of 4...	2
4554	3818			20 3 44.5	6.428	2	161 17 10.6	10.48	2	vF; vS; R; 3rd of 4	2
4555	3821			20 4 2.8	4.209	1	134 56 31.8	10.44	1	vF; pL; R; glbM	1
4556	3820			20 4 40.6	6.419	2	161 17 3.8	10.54	2	F; S; R; r; vS*att; 4th of 4	2
4557		VIII. 22		20 5 58.2	2.509	1	63 42 34.8	10.54	1	Cl; P; iC	1
4558	2070			20 5 59.3	2.261	1	54 34 19.8	10.54	1	Cl; pRi; * inv.....	1
4559	2071	VIII. 20		20 6 7.1	2.514	1	63 55 43.5	10.55	1	Cl; vB; vL; Ri; iC; st 6...11	2
4560	3822			20 6 37.5	4.584	1	143 12 32.8	10.64	1	pF; cL; pmE; glbM	1
4561		IV. 72		20 7 22.0	2.185	1	52 1 28.8	10.64	1	F; vL; vmE; * att	1
4562	3823			20 8 5.5	4.644	1	144 23 0.5	10.75	1	vF; L; iE	1
4563	3824			20 8 29.7	4.206	2	135 13 51.9	10.77	2	pF; S; R; vglbM	2
4564	3825			20 10 43.4	4.344	2	138 40 53.1	10.93	2	pF; S; R; svbM*12	2
4565	2072	IV. 13		20 10 45.8	2.419	2	59 51 33.3	10.89	2	!!; ⊙; F; S; vLE	7+
4566		VIII. 83		20 12 23.7	1.746	1	40 12 6.0	11.00	1	Cl; pRi; iC	1
4567			D'Arrest, 114	20 12 26	2.42	[2]	59 47 42	11.01	[2]	Cl+neb; S; st vS	0
4568	3826			20 14 4.0	4.433	1	140 52 0.6	11.18	1	F; S; R; glbM; am st.....	1
4569	3827			20 14 52.7	4.143	1	134 5 51.8	11.24	1	F; cS; R; bM.....	1
4570	2073			20 15 40.2	3.468	3	109 45 13.6	11.28	3	cL; E; bM*17; *10 att.....	3*
4571	2074			20 15 46.1	2.547	1	64 41 32.2	11.26	1	Cl; S; vLC; st 10...11	1
4572	2075	IV. 16		20 16 7.9	2.676	3	70 20 19.3	11.29	3	!!; ⊙; B; pS; R; 4St nr ...	6+
4573	2076	III. 141		20 16 44.2	3.590	1	115 14 51.2	11.36	1	cF; cL; vLE; vglbM; r; 3st p	3
4574	3828			20 17 43.6	4.267	1	137 28 57.8	11.44	1	pB; pL; gbM; 2st 10 nr.....	1
4575	2077	VIII. 56		20 18 5.5	2.137	2	49 40 14.4	11.42	2	Cl; pB; pS; P; pC; st 10...12	5
4576	2078		M. 29	20 18 51.9	2.212	1	51 56 3.6	11.48	1	Cl; P; iC; st L & S.....	6
4577	3830			20 20 50.3	4.274	1	137 56 31.9	11.67	1	vF; *12 att sp.....	1
4578	3831			20 22 0.0	4.141	1	134 40 55.8	11.74	1	eF; pS; R; vglbM	1
4579	3829			20 22 36.0	9.441	1	170 28 50.4	11.92	1	pB; cS; R; psmbM	1
4580	3832			20 23 2.0	3.727	4	121 17 38.7	11.81	4	pF; cS; R; gbM; bet 2 st ...	4
4581	3834			20 25 35.6	3.752	3	122 26 56.3	11.99	3	cB; L; mE 6 ^s .0; psibM	3
4582	2079	III. 142		20 25 52.1	3.118	2	92 29 44.3	11.99	2	vF; pL; E 0° or biN; p of 2	5
4583	2080			20 26 8.3	3.118	2	92 30 28.7	12.01	2	vF; vS; sf of 2	2
4584	3833			20 26 29.1	6.798	1	164 6 50.1	12.13	1	F; S; R; gbM; 5 st p.....	1
4585		I. 103		20 26 43.0	2.935	1	82 44 5.5	12.05	1	vB; L; gmbM; er	1
4586	2081			20 27 19.8	2.941	3	83 3 41.3	12.09	3	⊕; B; L; R; rrr; st 16...; *9p	3*
4587	3835			20 27 59.4	4.457	2	142 35 10.9	12.17	2	pB; cL; R; glbM; r	2
4588	3836			20 28 25.0	4.458	2	142 38 11.7	12.21	2	vF; cS; R; sibM; f of 2	2
4589	2082	VIII. 17		20 28 35.0	+2.646	1	68 13 54.9	-12.17	1::	Cl; vL; P; vLC	4

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
4590	2083	VI. 42		20 28 36.4	+1.211	3	29 49 56.8	-12.14	3	Cl; pL; eRi; pCM; st 11...16	5
4591		VII. 8		20 28 42.5	2.510	5	62 9 51.9	12.17	5	Cl; vB; vL; vRi; eC; st pL..	5
4592	3837			20 30 3.7	4.563	1	144 47 44.4	12.32	1	pB; pL; R; psbM	1
4593	3838			20 31 3.8	5.831	1	159 14 17.4	12.42	1	pF; L; mE; vglbMvs*	1
4594	2084	IV. 76		20 31 57.3	1.269	3	30 20 17.9	12.37	3	vF; vL; vg; vsbM; rr.....	4†
4595	3839			20 32 30.6	3.750	1	122 58 55.9	12.47	1	vF; L; R; gbM	1
4596	3840			20 33 4.6	4.497	2	143 51 23.1	12.53	2	vF; pS; eE; lbM	2
4597	2085	VIII. 23		20 34 40.7	2.769	1	73 50 36.0	12.60	1	Cl; P; vIC	2
4598	2086	III. 219		20 37 14.2	2.852	1	77 59 34.9	12.77	1	vF; S; stellar; * att	2
4599	3841			20 39 39.2	3.882	1	128 30 26.2	12.96	1	B; cS; R; pgmbM; 4 st p ...	1
4600	2088	V. 15		20 39 53.0	2.478	3	59 47 14.8	12.94	2	!!; pB; cL; eF; & Cygni inv	6†
4601	2087	II. 426		20 40 7.8	3.074	2	90 11 12.9	12.97	2	eF; S; R; bM	4
4602	2087, a		R. 3 novæ	20 40 ±	3.074	...	90 11 ±	12.97	...	Group of 5 with many st ...	0
4603											
4604											
4605	2089	II. 427		20 40 13.6	3.074	2	90 12 34.6	12.98	2	F; vS; R; bM	4
4606	3842			20 42 15.1	4.243	1	139 17 41.8	13.14	1	pB; S; IE; gbM	1
4607		II. 206		20 45 16.1	2.454	1	58 23 48.0	13.30	1	F; S; iE; r	1
4608	2090		M. 72	20 45 44.9	3.302	3	103 3 35.8	13.34	3	⊕; pB; pL; R; gmCM; rrr..	10
4609	3843			20 47 8.1	4.356	2	142 24 21.2	13.46	2	vF; S; E; p of 2	2
4610	3844			20 47 18.9	4.043	1	134 30 30.2	13.46	1	eF; cS; R	1
4611	3845			20 47 48.0	4.353	3	142 24 12.7	13.51	3	F; pL; vIE; vgbM; f of 2 ...	3
4612	3846			20 48 20.9	4.211	2	139 10 13.8	13.54	2	pF; S; vIE; gpmbM; vB * p	2
4613		VIII. 82		20 49 13.4	2.092	1	45 15 32.5	13.55	1	Cl; cL; st pS	1
4614	3847			20 49 27.4	4.529	2	146 6 24.7	13.61	2	ecF; vS; vmE0°; *13 att, n	2
4615	2091	VIII. 76		20 49 51.7	2.024	1	43 15 22.6	13.58	1	Cl; L; P; vIC	3
4616	2092	V. 14		20 50 35.2	2.478	2	58 50 11.8	13.64	1	!!; eF; cE; eE; eiF; bifurcate	4†
4617			M. 73	20 51 16.0	3.299	...	103 10 31.0	13.70	...	Cl??; eP; vIC; no neb	3
4618	2093			20 51 19.8	2.490	1	59 19 15.3	13.69	1	F; eL; neb & st	1†*
4619	2094			20 51 30.9	2.095	1	45 3 59.3	13.69	1	Cl; P; IC	2
4620		VIII. 58		20 51 33.5	2.123	3	45 53 24.0	13.70	3	Cl; P; IC; st L	3
4621	2096	V. 37?		20 53 48.2	2.142	(1)	46 13 4.8	13.84	1	F; cE; diff nebulosity	3
4622	2095			20 53 54.5	3.083	1	90 44 34.9	13.87	1	eF; S; E 0°	1
4623	3848			20 53 57.1	4.201	2	139 35 0.3	13.89	2	eF; cS; R; bM	2
4624	3849			20 54 13.9	4.203	1	139 40 27.7	13.91	1	eF; R; lbM; *11 f	1
4625	2097	I. 52		20 54 43.8	2.801	(3)	74 21 51.7	13.91	1	B; pL; R; gbM	4
4626	3850			20 55 23.1	4.345	3	143 6 3.6	13.98	3	pB; S; R; psbM; am st	3
4627	2099	I. 192		20 56 17.5	1.749	1	35 59 39.6	13.98	2	eB; L; E 45° ±; r; * att ...	5†
4628	2098	IV. 1	Lal. 40765	20 56 31.2	3.273	4	101 55 4.8	14.04	4	!!!; O; vB; S; elliptic	23*†
4629	2100			20 57 5.8	3.291	1	103 3 8.9	14.07	1	eF; pL; R; r	1
4630	2101			20 57 8.1	2.054	1	43 13 48.5	14.05	1	Cluster; no description	1
4631	3851			20 57 21.1	4.033	2	135 22 12.0	14.10	2	F; pL; E; vglbM; * p	2
4632	2102	II. 203		20 57 38.9	2.535	2	60 39 30.3	14.09	2	pB; cS; R; psbM; pB* np...	4
4633	3852			20 58 18.4	4.111	1	137 44 44.2	14.16	1	pF; S; R; bM; 2 st 12 n ...	1
4634		IV. 74		20 59 22.9	0.771	1	22 26 51.2	14.16	1	eF; *7 m in neb (?)	1
4635	3853			20 59 38.5	5.136	1	154 35 45.2	14.26	1	pB; cS; IE; pgbM	1
4636	3854			20 59 39.2	5.060	1	154 5 34.2	14.26	1	pF; cS; R; psbM; *7.8 p ...	1
4637	3855			20 59 45.4	4.185	1	139 52 10.5	14.25	1	eeF; S; R; B * * sf	1
4638	2103	VIII. 57		21 0 46.2	2.255	2	49 3 58.9	14.27	2	Cl; P; IC; st 10...	3
4639	3856			21 2 7.3	4.173	1	139 51 26.0	14.40	1	B; cS; R; pgmbM	1
4640	2105	VIII. 74		21 2 45.7	1.948	(1)	39 43 13.3	14.39	1	Cl of triple st; IC	2
4641	3857			21 2 47.1	5.457	1	158 51 57.9	14.47	1	vF; cS; R; glbM	1
4642	2104			21 3 36.5	2.825	1	75 7 5.2	14.46	1	Cl; IC	1
4643	2106			21 5 3.4	2.471	1	56 50 48.0	14.60	1	Cl; pRi; iF; st 11...15	1
4644	3858			21 5 39.3	4.079	2	137 47 45.7	14.61	2	pB; pL; IE; gbM	2
4645	2107			21 6 14.3	2.150	1	44 53 32.7	14.61	1	Cl; vL; pRi; E; st 10... ..	1
4646		III. 209		21 6 52.3	2.860	1	76 58 16.2	14.66	1	vF; S; R	1
4647	3859			21 6 55.9	4.114	2	138 56 41.3	14.69	2	B; cS; cE; psmbM; *10 f...	2
4648	2110	VI. 24		21 7 45.4	2.253	2	48 4 44.0	14.70	2	Cl; vF; pL; vRi; vC; st 15...18	4
4649	2108			21 7 48.6	3.008	1	86 3 33.4	14.72	1	eF;	1
4650	2109	III. 858		21 7 52.9	+3.035	2	87 44 17.4	-14.72	2	eF; pL; R; lbM	3

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
4651	3860		Δ. 406	21 9 26.5	+4.109	2	139 8 52.8	-14.84	2	vB; pS; E; mbM	2
4652	2111			21 9 33.0	2.424	1	54 23 16.7	14.81	1	Cl; no description	1
4653	2112	III. 145		21 12 36.6	2.640	1	64 9 17.3	14.99	1	F; S; vIE; r	3
4654	2113			21 14 20.1	3.216	1	99 22 29.7	15.11	1	vF; R; gbM; * nr	1*
4655	2114			21 15 26.3	1.717	1	32 59 50.8	15.14	1	Cl; F; pS; P	1
4656	3861			21 15 58.1	3.884	1	133 3 41.7	15.21	1	eF; vS; R; p of 2	1
4657	2115			21 16 16.6	2.021	1	39 47 4.3	15.19	1	Cl; P; IC	1
4658	3862			21 16 41.6	4.607	1	150 37 16.9	15.27	1	B; pL; IC; gpmbM	1
4659	3863			21 16 53.5	3.879	1	133 0 26.2	15.26	1	vF; pS; R; f of 2	1
4660	3864			21 17 58.9	4.085	1	139 40 10.1	15.33	1	eeF; vS; R	1
4661	2116	VII. 51		21 18 12.1	2.181	3	44 13 3.7	15.31	3	Cl; pS; pRi; pC; st 13... ..	4
4662	2117			21 18 44.9	2.447	1	54 5 44.1	15.33	1	Cl; P; st 10... ..	1
4663	3865			21 19 9.1	4.222	1	143 23 17.3	15.39	1	eF; pL; vme 90° 8; * s	1
4664	2118	VII. 50		21 19 37.1	2.135	1	42 34 45.6	15.38	1	Cl; P; ? neb	2
4665	3866			21 21 26.5	3.879	2	133 41 50.4	15.52	2	F; cL; IE; gvlbM; p of 2 ...	2
4666	2119			21 21 31.0	2.148	1	42 40 21.6	15.48	1	Cl; S; C; cE	1
4667	3867			21 21 37.5	3.880	2	133 45 57.1	15.53	2	F; S; R; vglbM; f of 2	2
4668	3868			21 22 49.4	3.757	1	129 14 1.3	15.59	2	eF; eS; R; pgbM	2
4669		III. 936		21 22 51.7	1.459	1	27 43 29.5	15.55	1	vB; er	1
4670	2120	 {	M. 15= Lal. 40815 }	21 23 9.9	2.895	1	78 27 22.3	15.59	2	{ !; ⊕; vB; vL; iR; } vsmbM; rrr; st vS. }	16
4671	3869			21 23 31.9	3.898	1	134 40 55.1	15.63	1	B; R; eS; psbM	1
4672	2121	III. 859		21 24 18.3	3.043	1	88 7 23.2	15.66	1	F; S; R; mbM; *14s	2
4673	2122	VII. 52		21 24 22.5	2.188	1	43 31 7.5	15.65	1	Cl; L; eRi; IC; st 10...13 ...	2
4674	3870		Δ. 263 ?	21 24 38.7	4.809	2	154 31 11.7	15.71	2	pF; cL; vIE; vgpmbM; r ...	2
4675	2123			21 25 11.3	2.819	1	73 11 28.0	15.70	1	Cl; IC	1
4676	2124	VI. 32		21 25 42.8	2.044	2	39 1 55.7	15.71	2	Cl; cL; vRi; pC; st 11...16	3
4677	3871			21 25 44.2	3.800	2	131 26 32.5	15.75	2	eF; S; R; gbM	2
4678	2125	 {	M. 2= Lal. 41928 }	21 26 12.5	3.091	2	91 26 37.2	15.76	3	{ !; ⊕; B; vL; g, pmbM; } rrr; st eS. }	19+
4679	3872			21 26 36.3	4.251	1	145 10 46.0	15.80	1	pB; pL; vme 127° 1; g, psbM	1
4680	3873			21 26 51.0	3.699	1::	127 24 5.7	15.81	1::	eF; pL; vgbM; *6 f 40s	1
4681	2126		M. 39	21 27 12.6	2.160	1	42 10 58.0	15.80	1	Cl; vL; vP; vIC; st 7...10 ...	3
4682	2127			21 29 5.1	2.244	1	44 37 6.0	15.90	1	Cl; P; IC	1
4683	3875			21 30 3.9	3.827	2	133 10 21.6	15.98	1	F; pL; R; vglbM; *13 inv...	2
4684	3874			21 30 20.1	4.755	1	154 32 2.4	16.02	1	vF; S; R; vS * nf	1
4685	3877			21 31 24.1	3.821	2	133 10 19.5	16.05	2	B; S; vIE; mbM	2
4686	3876			21 31 59.5	6.179	1	165 44 25.1	16.13	1	pF; R; g, psmbM; am st ...	1
4687	{ 2128 = 3878 }		M. 30	21 32 26.0	3.422	2	113 47 59.0	16.10	2	!; ⊕; B; L; IE; gpmbM; st 12...16.	3+
4688	3879			21 32 59.7	4.134	1	143 20 29.8	16.14	1	eF; cS; IE; vglbM	1
4689	3880			21 33 22.2	3.873	1	135 25 37.2	16.16	1	vF; cL; R; vglbM	1
4690	3881			21 33 31.6	3.626	1	125 4 56.2	16.16	1	eF; vS; am st	1
4691	3882			21 33 48.4	3.620	1	124 48 14.9	16.17	2	F; S; R; bM	2
4692	3883			21 36 32.5	3.965	2	139 3 59.4	16.32	2	F; S; R; glbM; p of 2	2
4693	3884			21 36 57.1	3.961	2	138 59 48.8	16.34	2	F; S; R; glbM; f of 2	2
4694	3885			21 37 10.8	3.904	1	137 9 38.5	16.35	1	F; S; R; gbM	1
4695			Anw. N. 47	21 38 22.8	3.296	...	99 28 19.2	16.36	...	Nebulous *10-11 or vSC (Cooper).	0
4696	3886			21 38 39.8	5.278	1	160 58 43.5	16.45	1	pB; S; R; vgbM; *9 f	1
4697	3888			21 38 45.1	4.022	3	141 12 49.8	16.44	3	pB; L; pmE; vgbM	3
4698	3887			21 38 56.2	4.466	1	151 21 18.5	16.45	1	eF; pL; R; p of 2	1
4699	3889			21 38 58.8	4.460	1	151 15 13.5	16.45	1	pB; pS; IE; gbM; f of 2 ...	1
4700	2129			21 39 8.8	2.013	1	36 1 22.4	16.42	1	Cl; S; P; IC	1
4701	2130	VII. 40		21 39 21.8	2.049	1	36 56 49.1	16.43	1	Cl; S; pRi; has a ruby * ...	2
4702	2131	IV. 75		21 39 45.0	1.387	2	24 32 15.8	16.44	2	!; eF; pL; gbM **	4
4703	3890			21 39 51.6	3.605	1	125 5 43.6	16.48	1	pB; S; R; glbM	1
4704	3891			21 41 22.0	3.609	2	125 31 43.2	16.56	2	pB; pL; R; vgbM; *14 att p	2
4705	2132	II. 261		21 41 44.2	2.773	2	68 29 7.2	16.56	2	F; pS; R; vglbM; r	3
4706		III. 696		21 42 15.3	+1.576	4	26 49 29.9	-16.57	4	vF; eS; R; r	4

No. of Cata- logue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s						
4707	3892			21 42 28.6	+4.229	1	147° 12' 6.1	-16.63	1	pF; cS; R; bM	1
4708	3893			21 42 30.4	4.187	1	146 13 34.1	16.63	1	F; L; R; g, psmbM	1
4709	2134	VII. 66		21 42 34.2	1.441	1	24 50 39.6	16.58	1	Cl; cL; cRi; pC; st 11...14	3
4710	2133			21 42 41.2	2.650	1	60 41 28.0	16.60	1	vF; ? a * inv in neb	1*
4711	3894			21 43 34.1	3.921	2	138 54 28.9	16.67	2	vB; pS; R; mbMN	2
4712	3895			21 44 13.2	3.907	1	138 32 26.0	16.70	1	B; S; R; in Δ of st 13.....	1
5077				21 45 12.5		...	40 53 46.4		...	See No. 5077.	
4713	3896			21 45 46.0	3.984	3	141 19 11.6	16.78	3	vF; pL; lE; vgbM; r.....	3
4714	3897			21 45 50.0	3.493	1	119 56 36.9	16.77	1	ceF; vS	1*
4715	3898			21 46 30.0	3.486	1	119 41 30.0	16.80	1	eF; S; E towards eF*	1
4716	3900			21 46 58.7	3.588	2	125 28 21.1	16.83	2	B; pL; iR; glbM; r	2
4717	3899			21 46 59.2	3.940	2	140 10 49.8	16.84	2	pB; S; lE; mbM.....	2
4718	2135	III. 452		21 47 29.0	3.042	3	87 42 49.8	16.84	3	F; pL; R; bM; r	4
4719	2136	VIII. 67		21 49 48.4	1.722	1	28 3 3.1	16.93	1	Cl; P; vLc	2
4720			D'Arrest, 115	21 49 53	3.04	[2]	87 42 42	16.95	[2]	Cl; vS; st 19 m; bet 2 st 16...	0
4721	3901			21 50 58.1	3.752	1	133 58 28.4	17.02	1	cF; cL; cE; glbM	2
4722	3902			21 51 10.5	3.521	3	122 33 8.4	17.02	3	F; pL; vLE; vglbM	3
4723	2137	III. 930		21 51 49.3	3.289	1	107 10 39.5	17.05	1	eF	2*
4724	3903			21 51 53.6	3.750	1	134 3 32.2	17.06	1	cB; S; vLE; svbMN	2
4725	3905			21 52 37.2	3.401	1	115 18 18.3	17.09	1	F; pS; R; vglbM; *10 f ...	1
4726	3904			21 52 54.3	3.975	3	142 25 3.4	17.12	3	pB; S; R; psbM.....	3
4727	3906			21 53 6.8	3.852	1	138 21 46.4	17.12	1	eF; S; R; *8 np	1
4728	2138	III. 692		21 53 29.4	3.243	1	103 57 3.1	17.13	3	vF; cL; E 135° ±; vgbM ...	4
4729	3908			21 53 52.3	3.513	4	122 32 38.5	17.15	4	pB; pL; lE; gbM; 1st of 4...	4†
4730	3909			21 53 53.1	3.514	4	122 38 8.5	17.15	4	cB; cS; R; sbM*; 2nd of 4	4†
4731	3910			21 53 55.3	3.515	2	122 39 58.5	17.15	2	cF; S; R; p of D neb; 3rd of 4	2†
4732	2141			21 53 55.9	2.106	1	35 50 51.1	17.13	1	Cl; vL; pRi; lC	1
4733	3911			21 53 59.0	3.514	6	122 39 31.5	17.15	6	B; pL; R; f of Dneb; 4th of 4	6†
4734	2139	II. 247		21 54 0.1	2.858	1	72 56 7.5	17.15	3	pB; pS; R; bMN; r; *sp...	4†
4735	2912			21 54 6.3	3.582	1	126 28 45.2	17.16	1	eF; S; R; *8 s 2'	1
4736	3907			21 54 21.4	4.525	2	154 43 13.6	17.18	2	cF; pS; vgbM.....	2
4737	2140	III. 693		21 54 26.8	3.339	1	111 13 26.9	17.17	1	vF; S; R; lbM; p of 2	2
4738	2142	II. 595		21 54 38.9	3.316	1	109 35 22.6	17.18	2	vF; pS; vLE 90°; lbM.....	4
4739	2143	II. 1		21 54 51.2	3.341	1	111 28 50.3	17.19	1	pB; pL; mE64° 3 bet 3 st; er	3
4740	2144			21 55 8.7	3.337	1	111 8 59.7	17.21	1	vF; pL; iR; vglbM; f of 2...	1
4741		III. 165		21 55 13.6	2.599	1	55 33 55.0	17.20	1	vF; am 5 or 6 st	1
4742	3913			21 56 20.4	4.543	1	155 19 8.9	17.27	1	vF; S; lE; vgbM	1
4743	3914			21 56 24.7	4.521	2	154 59 23.6	17.28	2	pB; S; R; pmbM	2
4744	2145			21 56 42.8	2.949	1	79 51 21.9	17.27	1	Cl; lRi; lC; st 9...10	1
4745	3915			21 56 50.6	3.900	3	140 47 57.3	17.29	4	cB; S; R; am st	4
4746	2146	II. 599		21 57 7.6	2.493	1	49 37 29.6	17.28	1	F; cS; cE; vglbM; er	2
4747	3916			21 58 4.3	4.529	2	155 22 59.5	17.35	2	vF; S; R; psbM; *11 p 3'...	2
4748	3917			21 58 7.9	3.839	1	140 40 49.5	17.35	1	pF; S; R; smbM	1
4749	3918			21 58 27.0	3.488	3	121 55 30.5	17.35	3	F; R; gbM; 1st of 4	3
4750	3920			21 58 39.2	3.486	1	121 51 49.2	17.36	1	eF; S; stellar; 2nd of 4	1
4751	3921			21 58 40.2	3.488	3	121 49 42.2	17.36	3	cF; R; stellar; 3rd of 4	3
4752	3922			21 58 47.8	3.484	1	121 43 31.9	17.37	1	pB; L; lE; gbM; 4th of 4 ...	1
4753	3919			21 58 53.7	4.140	1	148 6 29.6	17.38	1	pB; L; cE; gpslbM	2
4754	3923			21 59 24.6	3.451	1	119 43 57.3	17.39	1	vF; vS; R; almost a O	1
4755	2147	VII. 53		21 59 42.2	2.384	2	44 11 36.3	17.39	2	Cl; L; cRi; pC; st 9...12 ...	4
4756	2148			21 59 57.7	2.741	1	63 34 17.7	17.41	1	eF; R; bM; vF * np	1*
4757	3924			22 0 29.0	3.798	1	137 50 36.5	17.45	1	vB; pS; R; gbM	1
4758	3926			22 1 8.3	3.427	1	118 29 42.9	17.47	1	⊕; pL; iR; rr.....	1
4759	3925			22 1 30.8	4.794	1	159 20 46.7	17.51	1	pF; S; R; gbM	1
4760	2149	II. 207		22 1 37.4	2.682	1	59 20 35.6	17.48	1	B; pL; gbM; er	2
4761	2150	II. 897		22 2 34.9	3.273	1	107 19 55.1	17.53	1	pB; lE; r	2
4762	3927			22 2 47.3	4.487	2	155 32 6.2	17.56	2	pB; S; R; 2 st nr.....	2
4763	3928			22 3 13.8	3.462	4	121 14 22.2	17.56	4	F; S; R; gbM; r; 2vSstur...	4
4764	2151	III. 862		22 4 16.2	2.525	1	49 40 48.3	17.59	1	eF; pS; lE; r; am 3 st.....	2
4765	3929			22 5 15.4	3.392	1	116 50 37.8	17.64	1	pF; S; lE; bM	1
4766	3930			22 6 4.9	+3.436	4	120 4 3.9	-17.67	4	F; pL; R; vglbM	4

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s						
4767	2152	III. 931		22 6 37.7	+3.272	1	107° 45' 40.3	-17.69	1	vF; S; R; bM.....	2
4768	2153	II. 606		22 6 43.4	2.447	1	45 20 52.3	17.69	1	eF; S; cr.....	4
4769	3931			22 6 57.4	3.731	1	136 32 31.7	17.71	1	pB; S; pmE; psbM; p of 2...	1
4770	3932			22 7 7.9	3.730	1	136 32 16.4	17.72	1	F; vS; R; *8f; f of 2.....	1
4771		VIII. 63		22 7 8.2	2.127	1	33 42 56.0	17.70	1	Cl; S; P; IC.....	1
4772	2154			22 7 34.6	2.119	2	33 25 12.4	17.72	2	Cl; pC; has a ruby *10.....	2
4773	2155	VIII. 75		22 9 41.1	2.358	1	40 48 55.7	17.81	1	Cl; L; P; IC; st vL.....	3
4774	2157	VI. 29		22 10 2.5	2.236	1	36 22 2.4	17.82	1	Cl; C; st eS.....	2
4775	2156	III. 932		22 10 8.7	3.249	1	106 16 47.8	17.84	1	vF; S; vLE; vgbM; *13n.....	2*
4776		III. 863		22 11 0.4	2.563	1	50 10 13.2	17.86	1	vF; S; mbM.....	1
4777	3933			22 11 26.0	3.957	1	145 48 53.0	17.90	1	eeF; R; (?).....	1
4778		III. 864		22 12 28.8	2.569	1	50 8 9.4	17.92	1	vF; S; mE 165°+.....	1
4779	2158	III. 933		22 12 53.8	3.247	2	106 28 21.5	17.95	2	F; pS; R; gpmbM.....	4
4780	3934	III. 458		22 12 57.6	3.353	1	115 22 44.5	17.95	1	F; S; R; cr.....	2
4781	3935			22 15 6.5	3.395	1	119 3 25.8	18.04	1	vF; S; E; glbM; ?biN.....	1
4782	3936			22 15 10.0	3.403	1	119 39 16.8	18.04	1	eF; pL; R; vlbM.....	1
4783	2159			22 15 23.6	2.153	1	32 36 51.1	18.03	1	Cl; L; pRi; IC.....	1
4784	3937			22 15 26.0	3.450	1	123 3 31.5	18.05	1	eF; S; R; lbM.....	1
4785	3938			22 16 21.1	3.466	2	124 24 1.6	18.08	1	cB; pS; vLE; glbM; B ** sp.....	2
4786	3939			22 16 44.2	3.429	3	121 53 45.0	18.10	3	F; cS; vLE; p of 2.....	4
4787	3940			22 18 15.5	3.424	4	121 51 23.5	18.15	3	F; cS; vLE; f of 2.....	4
4788	3941			22 18 57.4	4.079	1	150 52 54.3	18.19	1	eeF; lE; vgvlbM; 3 st sf.....	1
4789	3942			22 19 7.4	3.478	2	125 51 19.3	18.19	2	vF; pS; R; vgvlbM.....	2
4790	2160	II. 248		22 19 38.1	2.916	2	74 33 55.0	18.20	2	F; cS; R; gbMS*; 3 st nr.....	4
4791	2161			22 19 39.4	2.198	1	32 52 26.3	18.19	1	Cl; L; pRi; IC; st 10...16...	1
4792	3943	II. 469		22 20 50.7	3.336	2	115 34 20.5	18.25	2	cF; cS; lE; r; * inv.....	3
5078				22 20 50.4		...	115 34 +		...	(See No. 5078).....	0
4793	2162			22 20 52.2	2.770	1	61 36 48.8	18.24	1	vF; S; R; am st.....	1
4794	3944			22 21 12.0	3.474	1	126 10 21.2	18.26	1	vF; S; R; gbM.....	1
4795			Auw. N. 48	22 22 5.6	3.285	...	111 32 59.8	18.29	...	!; pF; vL; E or biN (Harding)	0
4796	2163			22 22 21.2	2.365	1	37 53 26.3	18.29	1	Cl; P; lC; st 12...13.....	1
4797		VII. 41		22 22 37.4	2.380	2	38 24 45.0	18.30	2	Cl; iR; lC; st vS.....	2
4798	3945			22 22 59.4	3.503	1	128 33 0.1	18.33	1	eF; S; R; p of 2.....	2
4799	2165, a		R. nova	22 23 13.6	3.213	::	104 40 30.3	18.34	::	vF; E np to sf (nisi=h.2164)	0
4800	3946			22 23 23.8	3.501	1	128 32 9.8	18.34	1	eF; S; R; f of 2.....	2
4801	2164			22 23 28.3	3.213	2	104 44 11.8	18.34	2	vF; cS; vglbM.....	2
4802	2165	IV. 31		22 24 53.0	3.213	4	104 50 30.3	18.39	4	F; pS; R; vsbMFSRN.....	5
4803			D'Arrest, 116	22 24 57	2.76	[1]	59 45 18	18.38	[1]	vF; pL; vlbM; h.2166 dist 2'	0
4804	2166			22 24 58.6	2.760	2	59 45 25.3	18.39	2	vF; S; R; vglbM.....	2
4805	3948			22 25 28.5	3.352	1	117 58 4.4	18.42	1	vF; S; lE; *11 p.....	1
4806	3947			22 25 32.6	3.541	1	131 39 53.4	18.42	1	F; pL; pmE.....	1
4807	2167	II. 476		22 26 55.7	3.173	5	101 4 53.2	18.46	5	vF; pL; R; glbM; r.....	6
4808	2168	II. 428		22 27 3.9	3.027	2	85 9 17.9	18.47	2	pF; S; R; psbM; r.....	4
4809	2169	III. 180		22 27 40.9	2.886	1::	70 23 45.6	18.48	1::	eF; vS; R; *9 s.....	2
4810	3949			22 28 2.3	3.331	2	116 45 55.0	18.50	3	cB; L; mE 0°; vlbM.....	3
4811	2170	III. 237		22 29 46.2	2.877	1	69 5 50.5	18.55	1	F; S; iR; vglbM.....	2
4812	3950			22 29 46.2	3.466	2	127 57 13.5	18.55	2	vF; S; vLE; gbM.....	2
4813	2171			22 30 32.5	2.984	3	80 11 4.6	18.58	4	eF; pS; lE 90°; vglbM.....	4
4814	3951			22 30 37.2	4.257	1	157 12 16.0	18.60	1	pB; pS; mE 90°.....	1
4815	2172	I. 53		22 30 39.5	2.736	1	56 20 5.6	18.58	3	B; pL; pmE 160°; smbM.....	4†
4816											
4817											
4818	2172, a		R. 5 novæ	22 30 ±	2.736	...	56 20 ±	18.58	...	5 near; positions measured; no distances.	0*
4819											
4820											
4821	2173	II. 233		22 30 44.0	2.857	2	66 55 34.6	18.58	2	cB; S; mE 163° 0; vsmbM *11; p of 2.	5
4822	3950	(No. 2?)		22 30 50.5	3.466	1	127 55 48.4	18.62	1	eF; ??.....	1?
4823	2174	III. 166		22 30 54.0	2.736	1	56 17 51.3	18.59	1	cF; vS; E.....	2
4824	2175	II. 234		22 31 5.7	+2.858	2	66 55 53.3	-18.59	2	F; pS; mE 90°±; vglbM; f of 2.	5

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and II.
	Sir J. II.'s Catalogues of Nebulæ.	Sir W. II.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s		° ' "	"			
4825	2176			22 32 57.9	+2.982	1	79 42 40.2	-18.66	1	eF; pL; E	2
4826	2177			22 34 15.4	2.333	1	33 20 2.3	18.69	1	Cl; vL; pRi; vLC	1
4827	2178	II. 705		22 35 6.6	2.218	1	29 27 5.4	18.72	1	B; S; R; pgvbm; er	2
4828	3952			22 35 26.3	3.451	1	128 36 27.8	18.74	1	eeF; S; R; * f 90°, 40°	1
4829	3953		Δ. 255?	22 36 9.6	4.117	1	155 51 19.9	18.77	1	F; S; R; bM	1
4830	3954			22 36 30.8	3.351	1	120 47 3.9	18.77	1	F; pL; vmE0°; vglbm	1
4831	2179	II. 442		22 37 14.0	3.079	2	90 53 42.0	18.80	2	F; S; R; psbM	5
4832	3955			22 37 30.5	3.475	1	130 4 35.7	18.81	1	F; cS; lE; glbm	1
4833	2180	II. 477		22 38 40.4	3.166	4	101 44 29.8	18.84	5	vF; pL; R; lbM	6
4834	2181	II. 598		22 40 12.8	3.262	3	113 2 10.3	18.89	3	pB; S; vLE; vgmbM; S * nr	4
4835	2181, a		R. 7 novæ	22 40 ±	3.262	...	113 2 ±	18.89	...	No descriptions	0
4836											
4837											
4838											
4839											
4842	2182	VIII. 77	C. H.	22 41 23.2	2.374	2	32 39 8.7	18.91	2	Cl; pL; pRi; lC; st 9...13 ..	3
4843	3956			22 42 23.0	3.408	1	127 34 56.5	18.95	1	eF; vS; R; * 12 att np	1
4844			{ D'Arrest, 117 = R. nova }	22 42 30	2.99	[2]::	79 10 30	18.94	[2]	vF; vS; R; III. 216 nf	0
4845	2183	III. 216		22 42 56.0	2.988	1	79 7 46.2	18.96	2	eF; S; R; glbm; * 11 np ...	5
4846	2184	III. 217		22 43 2.5	2.987	1	79 2 36.2	18.96	2	eF; S; R; pgbm; f of 2	5
4847			D'Arrest, 118	22 43 20	2.99	[2]	79 6 0	18.97	[2]	eF; vS; R; 2 st 11, s	0
4848	2184, a		R. nova	22 43 22.5	2.99	...	79 8 2.2	18.96	...	One of 5. See h. 2183, 2184, D'Arrest, 117, 118.	0*
4849	2185	II. 443		22 43 25.0	3.088	2	92 16 54.6	18.98	2	eF; cS; R; sbM * 13; * np ..	4
4850	2186	II. 702		22 44 15.4	3.238	3	111 21 1.0	19.00	3	pB; pS; lE 120°+; mbM ...	4
4851	2187	II. 453		22 44 21.1	3.118	2	96 18 6.0	19.00	2	vF; pL; lE; vgbM; r	3
4852	2188			22 44 36.2	2.534	1	38 33 54.0	19.00	1	Cl; vP	1
4853	2189			22 45 12.8	3.069	2	89 38 56.4	19.02	2	pF; pS; R; gbM	2
4854	2189, a		R. 3 novæ	22 45 ±	3.069	...	89 39. ±	19.02	...	"A group of 4," incl h. 2189	0
4855											
4856											
4857	3957			22 45 42.2	3.506	1	136 5 36.8	19.04	1	pF; lE; glbm; vS * inv	1
4858	3958			22 46 23.9	3.420	1	130 3 30.2	19.06	1	vF; S; R	1
4859	3959			22 46 49.0	3.930	1	154 26 29.6	19.08	2	pB; pS; R; vglbm	1
4860	3960		Δ. 518	22 47 2.8	3.421	2	130 24 35.6	19.08	2	cB; L; vmE 43°3; mbM ...	2
4861	3961			22 47 45.7	3.456	1	133 23 49.0	19.10	1	eF; vL; * 7 nf	1
4862	3962			22 48 42.2	3.940	1	155 46 26.1	19.13	1	pB; cS; R; gpmbM	2
4863	3963			22 48 42.4	3.383	3	127 46 23.4	19.12	3	cB; vL; vLE; vglbm	3
4864	2190	VII. 43		22 48 46.6	2.356	2	29 55 6.3	19.09	2	Cl; pRi; cC	3
4865	3964			22 49 1.9	3.385	3	128 5 31.1	19.13	4	cB; L; vLE; gpmbM; rr	4
4866	2191	III. 745		22 49 18.9	2.466	1	33 38 14.1	19.13	1	vF; pL; iF; er	2
4867	3965			22 49 21.4	3.428	2	131 49 4.8	19.14	2	F; cL; vLE; vgmbM	2
4868	2192	III. 576		22 49 28.5	2.784	1	54 22 38.8	19.14	1	vF; cS; R; stellar; * p	3
4869	2193			22 50 18.4	2.404	1	30 45 23.2	19.16	1	Cl; P; pC; st 9...11	1
4870	2194	III. 465		22 51 3.9	2.986	1	77 37 14.6	19.18	1	eF; S; R	4
4871	2195	III. 243		22 51 12.3	2.886	1	64 37 4.6	19.18	2	F; pS; E 90°; gbM; er	3
4872	2195, a		R. 2 novæ	22 51 ±	2.886	...	64 37 ±	19.18	...	{ 2 of a group of 3; inv in F neb. }	0
4873											
4874	2196			22 51 36.5	2.543	1	36 23 55.3	19.19	1	Cl; vL; E	1
4875			D'Arrest, 119	22 52 32	2.97	[2]	75 12 12	19.21	[2]	pF; R; bet 2 st 16; * 13 nr...	0
4876	2197	II. 450		22 52 46.6	3.163	2	103 33 10.4	19.22	2	F; vS; vLE; smbM; er; p of 2	4+
4877	2198	II. 451		22 52 46.9	3.163	2	103 34 46.4	19.22	3	F; vS; vLE; smbM; er; f of 2	5+
4878			Auw. N. 49	22 53 6.3	3.144	...	101 16 41.1	19.23	...	* 11-12 in neb (Markree Obs. Oct. 8, 1855).	0
4879	2199	II. 251		22 53 8.0	2.968	2	74 46 10.1	19.23	2	pB; L; E 175°; vgbM	3
4880		II. 249		22 54 3.1	2.967	2	74 21 17.2	19.26	2	F; cS; lE; lbM; pB * p	2
4881	3966			22 54 13.6	+3.387	1	130 19 19.2	-19.26	1	F; L; mE 33°8; vglbm	1

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	"			
4882	3967			22 54 16.6	+3.387	1	130 20 14.9	-19.27	1	eF; vL; vlbM.....	1
4883	2201	II. 212		22 54 17.6	2.861	1	60 36 32.2	19.26	1	cB; cL; lE; gmbM; r; 2S st n	2
4884	2200	II. 590		22 54 20.5	3.065	2	88 59 35.2	19.26	2	cF; cS; psbM.....	3
4885	3968			22 54 55.7	3.397	2	131 34 57.6	19.28	2	cF; pS; vmE 5° ±; * 11 np...	2
4886	2202	III. 210		22 54 56.3	2.971	1	74 46 22.6	19.28	1	vF; S; lE; p of 2.....	2
4887	2203	III. 211		22 55 5.3	2.971	1	74 46 59.6	19.28	1	vF; vS; f of 2.....	2
4888	2204	III. 230		22 56 12.2	3.020	1	81 52 19.7	19.31	1	vF; vS; vsmbM * 12.....	2
4889	3969			22 56 13.3	3.514	1	140 52 10.7	19.31	1	eF; pL; R; glbM; * 11 np...	1
4890		III. 202		22 56 22.3	2.969	1	74 8 43.7	19.31	1	eF; vS.....	1
4891	3970			22 57 19.6	3.411	2	133 51 40.8	19.34	2	F; S; R; Δ with 2 st 7.....	2
4892	2205	I. 55		22 57 56.4	2.999	4	78 25 53.5	19.35	4	pB; cL; mE 11° 9; bet 2 st...	7*+
4893	2206			22 58 40.4	3.055	1	87 12 42.2	19.36	1	vF; S; E; psbM.....	1
4894	{ 3971 = 3972 }			22 59 21.2	3.330	4	127 1 37.6	19.38	4	pB; S; R; lbM; * 8.9 att s...	4*
4895	2207			22 59 23.7	2.845	1	56 39 4.6	19.38	1	vF; S; R; bM; * 10 p.....	1
4896	2208	III. 558		23 1 2.4	3.169	1	106 22 33.4	19.42	1	eF; L; bet 2 D st.....	2
4897	3973			23 1 56.3	3.379	2	134 10 34.8	19.44	2	pB; cL; lE; vgbM * 13.....	2
4898	2209	III. 203		23 2 5.5	2.971	2	72 35 4.8	19.44	2	vF; L; pmE 45°; lbM.....	3
4899	2210	III. 184		23 4 29.6	3.087	2	92 54 44.3	19.48	2	cF; vS; R; sbM * 15.....	4
4900	{ 2211 = 3974 }	II. 2		23 4 33.6	3.247	2	119 17 53.3	19.49	2	pB; cS; R; psymbM; * 10 np	5
4901	2212			23 4 45.0	3.004	2	77 49 34.0	19.50	2	eF; bM*; (?).....	2
4902	2213	VII. 44		23 5 26.7	2.536	1	30 11 23.7	19.51	2	Cl; pRi; pC; fan-sh; st pB...	4
4903	2214	III. 220		23 5 46.8	3.006	2	78 4 51.4	19.52	3	F; cS; R; vglbM; r.....	7
4904		III. 470		23 6 36.7	3.125	1	99 57 29.1	19.53	1	cF; vS.....	1
4905	3975			23 7 0.8	3.364	1	134 21 49.8	19.54	1	pB; S; lE; pgbM.....	1
4906	2215	II. 429		23 7 28.8	3.052	2	86 15 44.5	19.55	2	vF; cS; R; bM; sp of 2.....	3
4907		II. 706		23 7 34.3	2.537	2	29 14 59.5	19.55	2	vF; L; 2 pB st inv.....	2
4908	2217			23 7 36.2	2.945	1	67 4 19.5	19.55	1	F; S; R; psbM.....	1
4909	2216	II. 430		23 7 37.8	3.052	2	86 13 35.5	19.55	2	B; L; mE 97° 5 (2 est); mbM; nf of 2.	3
4910	3976			23 7 49.1	3.313	1	129 17 54.2	19.56	2	F; S; vLE; vglbM; * 10 att.	2
4911	2218			23 8 5.0	2.975	1	71 46 56.2	19.56	1	np of 2 neb.....	1
4912	2218, α		R. 2 novæ	23 8 ±	2.975	...	71 47 ±	19.56	...	2 of 4 incl h. 2218, 2219...	0
4913			D'Arrest, 120	23 8 17	2.94	[2]	65 29 18	19.57	[2]	vF; vS; (d'Arrest) * 16 p 11 ^s	0
4914				23 8 17.5	2.975	1	71 47 55.9	19.57	1	eF; S; R; sf of 2.....	2
4915	2219	III. 181		23 8 25.2	3.346	1	133 21 29.9	19.57	1	B; S; mE 90° ±; vsbM * 13...	1
4916	3977		Δ. 475?	23 8 27.0	3.041	::	84 3 21.6	19.57	::	No description.....	0
4917	2224, α		R. nova	23 8 30.3	3.010	1::	78 11 12.9	19.57	1::	F; R; bM.....	1
4918	2221			23 8 32.3	3.087	1	93 9 4.9	19.57	1	cF; pL; R; B * f.....	3
4919	2220	II. 235		23 8 44.8	3.007	3	77 27 35.6	19.58	3	F; cS; R; lbM * 6; p of 2.....	6
4920	2222	III. 221		23 8 51.2	3.041	1	84 4 21.6	19.58	1	cB; pS; iR; psbM.....	2
4921	2224	II. 467		23 8 53.6	3.008	2	77 34 16.6	19.58	4	cF; cS; R; sbM * 16.....	6*
4922	2223	III. 222		23 9 8.7	3.087	2	93 6 55.6	19.58	2	vF; pS; E; er; 3S st inv.....	2
4923		III. 185		23 9 43.0	3.007	1	77 17 25.3	19.59	1	eF; eS.....	1
4924		III. 238		23 10 14.0	3.099	1	95 30 25.0	19.60	1	F; S; smbM.....	1
4925		II. 454		23 10 14.2	2.980	1	72 4 20.0	19.60	1	vF; am vS st.....	3
4926	2225	III. 182		23 10 39.7	3.330	1	132 52 58.7	19.61	1	pB; L; pmE; gbM.....	1
4927	3978		Δ. 476?	23 10 47.0	3.098	2	95 24 42.7	19.61	2	pB; pS; iR; gbM.....	6
4928	{ 2226 = 3979 }	II. 236		23 11 9.4	3.329	1	132 59 53.4	19.62	1	pB; pL; pmE; gbM; p of 2..	1
4929	3980		Δ. 477, 1	23 11 21.9	3.097	1	95 12 23.4	19.62	1	eF; vS.....	1
4930		III. 186		23 11 36.8	3.326	1	133 1 8.1	19.63	1	F; pL; pmE; gbM; f of 2...	1
4931	3981		Δ. 477, 2	23 11 38.9	3.112	1	98 19 42.1	19.63	2	cF; S; R; pspmbM.....	4
4932	2227	II. 431		23 11 49.7	+3.116	2	99 15 31.1	-19.63	2	pF; cL; pmE 0° ±.....	3*
4933	{ 2228 = 3982 }	I. 104									

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. II.'s Catalogues of Nebulæ.	Sir W. II.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s		° ' "	"			
4934			D'Arrest, 121	23 12 30	+3.04	[2]	82 42 24	-19.64	[2]	F; S; R; Δ with 2 st 19, n ...	0
4935	2229			23 12 51.0	3.036	1	82 21 19.5	19.65	1	eF; eS	1
4936	2230	II. 439		23 13 9.5	3.037	2	82 33 59.5	19.65	2	eB; pS; R; psbM	3
4937	3983			23 13 21.1	3.584	1	152 53 32.2	19.66	1	eF; eS; am 5 st; (?)	1
4938	2231	III. 435		23 13 25.2	3.036	1	82 21 49.2	19.66	1	F; vS; R; psbM	2
4939	2232	II. 250		23 13 27.2	2.994	2	73 32 34.2	19.66	2	pB; cS; R; snbM	4
4940	2233	II. 440		23 13 38.0	3.037	2	82 32 58.2	19.66	1	eB; pS; R; psbM	3
4941			D'Arrest	23 14 5.8	2.998	[4]	74 12 23.9	19.67	[4]	eB; pS; bM* (D'Arrest, Resultate).	0*
4942	2231, a		R. nova	23 14 13.7	3.036	::	82 21 49	19.67	::	Ep and f	0
4943			D'Arrest, 122	23 14 22	3.04	[2]	82 33 6	19.67	[2]	vF; vS	0
4944	3985			23 14 21.5	3.314	2	133 14 52.6	19.68	2	F; S; R; lbM	2
4945	3986			23 14 30.5	3.717	2	158 25 49.6	19.68	2	F; vS; E90°; psbM	2
4946	2234	II. 441		23 14 36.4	3.035	1	81 52 38.6	19.68	1	F; S; F * att	2
4947	2235	IV. 52		23 14 38.8	2.616	1	29 35 10.6	19.68	1	vF; *9 inv a l excentric	3
4948	3987			23 15 3.6	3.219	1	120 2 41.3	19.69	1	eF; S; R; sbM	1
4949	3984			23 15 6.0	4.986	1	172 40 19.0	19.70	1	vF; pL; R; vlbM; *nr	1
4950	2236	II. 600		23 15 23.6	2.858	2	49 54 46.3	19.69	2	eF; L; mE0°+; vlbM; r ...	4+
4951	3988			23 16 17.5	3.215	1	120 8 51.7	19.71	1	vF; S; R; glbM	1
4952		III. 473		23 16 47.9	3.002	1	74 0 18.4	19.72	1	eF; cL; se st f; (?)	1
4953		III. 218		23 16 48.7	3.032	1	80 50 18.4	19.72	1	eF; pS; lE	1
4954	3989			23 17 17.8	3.464	2	148 33 48.1	19.73	2	pF; pS; R; glbM; p of 2 ...	2
4955	3990			23 17 37.9	3.462	1	148 39 59.1	19.73	1	eF; S; R; f of 2	1
4956	2237			23 17 51.8	3.010	1	75 29 12.1	19.73	1	vF; pS; R; gbM	2
4957	2238		M. 52	23 18 3.2	2.643	1	29 10 20.1	19.73	1	Cl; L; Ri; mCM; R; st9...13	8
4958	3991			23 18 14.6	3.674	1	158 47 39.8	19.74	1	eF; vS; R; psbM; *10 p 22	2
4959	3992			23 18 40.6	3.452	1	148 34 57.5	19.75	1	eF; R	1
4960	3994			23 18 48.0	3.266	1	129 59 55.5	19.75	2	{ eF; S; R } D neb; 4 st p..	2
4961	2239	III. 212		23 18 52.9	3.016	1	76 33 33.5	19.75	1	vF; vS; R; psbM	2
4962	2240			23 18 52.9	2.956	1	63 44 7.5	19.75	1	F; vS; psmbM; *10 p	1
4963	3993			23 18 56.4	3.590	1	156 2 49.5	19.75	1	eF; cL; R; vglbM	1
4964	2241	IV. 18		23 19 9.9	2.864	5	48 13 57.5	19.75	5	III; O; vB; pS; R; blue.....	16*+
4965		III. 438		23 20 0.9	3.112	1	100 11 16.2	19.76	1	eF; S; stellar	1
4966	2242	III. 226		23 20 15.0	3.025	4	78 18 33.9	19.77	4	{ (H.) vF } S; R; vsmbM; { (h.) pB } *9 p (? var).	6*
4967	2242, a		R. nova	23 20 15.0	3.025	::	78 24 33.9	19.77	::	vF; S; 6' s of h. 2242	0
4968	2243			23 20 49.8	3.040	1	81 59 10.6	19.78	1	F; eS; gbM; p of 2	1
4969	2244			23 20 59.8	3.041	1	81 59 55.6	19.78	1	vF; S; R; gbM; f of 2	1
4970	3995			23 21 2.6	3.457	1	150 29 2.6	19.78	1	B; S; lE; vsmbM*11	1
4971	2245	II. 226		23 21 28.0	2.985	2	68 20 38.3	17.79	2	vF; pL; vLE; lbM; am 4 st...	3+
4972	2246	III. 860		23 21 36.7	2.938	(1)	58 25 6.3	19.79	1:	vF; S; R; lbM; r	2
4973	2247	II. 242		23 21 52.1	3.008	2	73 28 4.3	19.79	2	vF; S; iR; r; * f	4
4974			D'Arrest, 123	23 21 55	3.06	[1]	87 19 4	19.79	[2]	eF; *14 p 13.7, l n	0
4975	2248	III. 426		23 23 21.9	3.061	1	86 52 7.7	19.81	1	eF; cL; R; gbM; *nr	3
4976	2249	VIII. 69		23 23 29.0	2.839	2	41 38 50.7	19.81	2	Cl; P; lC; st 7...11	4
4977			D'Arrest, 124	23 23 47	3.06	[2]	87 12 42	19.81	[2]	vF; vS; *11 f 1.8, n 85''	0
4978	3996		Δ. 347?	23 24 48.4	3.352	1	144 52 22.1	19.83	1	pF; L; R; vgbM	1
4979	3997			23 25 20.0	3.325	2	142 28 11.8	19.84	2	eB; S; lE; psbM; *8 f	2
4980	2250	III. 213		23 25 20.6	3.019	1	74 55 35.8	19.84	1	eF; pL; Δ with 2 st 10	2*
5079				23 25 33.1		...	96 22 12.0		...	See No. 5079.	
4981		III. 187		23 25 56.9	3.084	2	93 31 10.5	19.85	2	eF; pL; stellar	1
4982			D'Arrest, 125	23 26 4	3.09	[1]	93 39 12	19.85	[1]	eF; pL; 3 st 11 & 12 f.....	0
4983	3998			23 26 48.5	3.494	1	156 19 38.2	19.86	1	eeF; pL (certain)	1
4984		III. 188		23 27 3.0	3.084	1	93 41 10.2	19.86	1	eF; stellar	1
4985	3999			23 27 43.9	3.347	2	146 47 2.9	19.87	2	B; eS; E; g. sbM; *8.9 p ...	2
4986	2251			23 27 44.2	3.022	1	74 42 26.9	19.87	1	vF; vS; gbM; *nf 1'	1
4987	2252			23 27 53.0	3.059	2	85 52 26.9	19.87	2	eF; *12 p; sp of 2	2
4988	2253			23 28 2.0	3.058	1	85 49 1.9	19.87	1	vF; nf of 2	1
4989	2254	III. 579		23 28 2.6	2.900	2	46 27 42.9	19.87	2	eF; S; R; *9.10 p, v nr.....	3
4990	2255	VIII. 62		23 28 15.5	+2.510	2	17 51 23.9	-19.87	2	Cl; L; P; lC; st 8, 10...15...	5

No. of Cata- logue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulae.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	H.		h m s	s		° ' "	"			
4991	2256	II. 244		23 28 35.2	+3.026	1	75 27 48.6	-19.88	2	F; S; R; psbM; stellar	4
4992	4000			23 28 56.7	3.209	1	128 13 2.6	19.88	1	pB; L; E; vgbM.....	1
4993	2257			23 29 2.6	3.067	1	88 36 51.3	19.89	1	pB; S; R; psbM; *12 sp ...	1
4994	2257, a		R. nova	23 29 10.6	3.067	::	88 36 51.3	19.89	::	No description	0
4995	2258			23 29 20.0	3.073	1	90 28 43.3	19.89	1	F; pL; IE; gbm; *10 s	1
4996	2259	III. 146		23 31 28.2	2.992	1	63 45 34.7	19.91	1	F; S; IE; bM; am st	2
4997	2260	II. 432		23 31 35.5	3.092	3	97 17 32.7	19.91	3	pF; cL; E 12° ±; vgbM.....	7
4998	2261	I. 110		23 31 41.7	3.111	2	103 43 56.7	19.91	2	{ (H.) cB } ; cL; E; gmbM; { (h.) cF } r (? var.)	4*
4999		III. 189		23 32 21.2	3.087	1	95 24 5.4	19.92	1	eeF	1
5000	2262	I. 111		23 32 38.8	3.107	2	103 4 12.4	19.92	2	pB; pL; iR; mbM	4
5001	4001			23 34 41.4	3.397	1	156 45 1.5	19.95	1	eF; S; R; p of 2	1
5002	4002			23 34 49.4	3.396	1	156 44 21.5	19.95	1	eF; cS; R; f of 2	1
5003	2263	II. 208?		23 35 17.3	3.006	1	64 32 13.5	19.95	1	vF; *14 att 255°	1*
5004		II. 208		23 36 45.2	3.010	1	64 38 3.2	19.96	1	cL; R; *10-11 np	1*
5005	2264	II. 255		23 37 8.5	3.049	3	80 0 26.2	19.96	5	cB; cS; gmbM; r; B * f.....	7
5006	2265	II. 256		23 37 13.4	3.051	1	80 50 20.2	19.96	3	pF; S; R; *15 sf	4
5007	4003			23 37 35.5	3.190	3	133 41 27.9	19.97	3	cB; S; vLE; svbmM*14	3
5008	2266			23 38 26.1	2.757	1	21 1 30.9	19.97	1	vL; surrounds *8	1
5009	4004			23 38 29.5	3.141	2	120 17 4.6	19.98	2	vF; S; R; gmbM; *12 f ...	2
5010	2267	III. 427		23 39 27.3	3.066	2	86 58 39.6	19.98	2	cF; pL; vLE 0°; lbM; 2 Bstnr	4
5011	2268	II. 213		23 40 2.3	3.011	1	61 17 42.3	19.99	1	cF; cL; vLE; vglbM; r	2
5012	4005			23 40 35.8	3.137	2	121 17 45.3	19.99	2	B; cL; R; psbmM	2
5013	2269	III. 437		23 40 50.1	3.064	1	83 54 46.3	19.99	2	F; S; R; gbm; er	3
5014	2270			23 41 35.4	3.066	1	86 36 8.0	20.00	2	vF; cL; R; vglbM; *13 n ...	2
5015	2271	III. 854		23 42 7.7	3.015	5	59 47 45.0	20.00	5	cB; vS; R; psbM; rr	7*
5016	2272	VII. 53		23 43 4.8	2.849	1	22 45 49.7	20.01	1	Cl; pRi; pC; st 11...15	4
5017	4006			23 43 34.1	3.151	1	131 31 2.7	20.01	1	B; pL; R; gbm	1
5018	2273, a		R. nova	23 43 ±	3.028	...	63 37 ±	20.01	...	vvF; a little np h. 2273	0
5019	2273			23 43 52.2	3.028	1	63 37 55.7	20.01	1	vF; S; iF; vF * inv	1
5020	2274	II. 230		23 43 55.7	3.041	2	70 37 26.7	20.01	2	pF; S; R; mbM; sp of 2 ...	4*
5021	2274, a		R. nova	23 44 13.9	3.04	...	70 40 43.7	20.01	...	Seen and meas. with h. 2274, 2275.	0
5022	2275	II. 231		23 44 17.1	3.042	2	70 39 16.7	20.01	2	pB; pL; E 90°; bM; nf of 2	4
5023	2276			23 44 38.0	3.049	1	74 31 22.4	20.02	1	Cl of sc st 10 m	1
5024	2277	II. 851		23 45 4.8	3.024	2	59 31 6.4	20.02	3	vF; cS; R; * nf	5
5025	2278	III. 231		23 46 11.5	3.063	2	82 54 5.4	20.02	2	cF; S; R; psbM; stellar; 1st of 4.	3
5026	2279	III. 232		23 46 19.0	3.063	2	82 53 50.4	20.02	2	pF; S; R; psbM; stellar; 2nd of 4.	3
5027	2280			23 46 38.2	3.063	1	82 54 18.1	20.03	1	F; S; R; 3rd of 4	1
5028	2281	III. 233		23 46 45.2	3.063	2	82 48 10.1	20.03	2	pF; pL; IE; gbm; 4th of 4	4
5029	2282	II. 468		23 48 10.1	3.066	4	84 51 58.1	20.03	4	pB; pS; iR; psbM; r; *7 p30°	5
5030	2283			23 49 42.8	2.975	2	29 23 16.8	20.04	2	Cl; S; pRi; vC; st 10, 13...	2
5031	2284	VI. 30	C. H.	23 49 58.5	2.994	1	34 3 46.8	20.04	1	Cl; vL; vRi; vmC; st 11...18	3
5032	2285	VII. 56		23 50 0.7	2.979	1	29 33 50.8	20.04	1	Cl; pRi; pC	2
5033	2286			23 50 47.4	3.063	1	80 0 18.8	20.04	1	vF; vS; (?)	1
5034	2287			23 51 34.3	2.999	1	30 45 41.8	20.04	1	Cl; vL; P; iC; st 7, 10...	1
5035	4009			23 51 45.2	3.136	1	146 14 16.8	20.04	1	pB; cS; R; gmbM	1
5036	2288	III. 466		23 51 47.0	3.065	1:	80 2 53.5	20.05	1	vF; pS; iR	2
5037	2289	III. 867		23 51 50.3	3.070	1	87 8 41.5	20.05	1	eF; pS; iR; lbM	2
5038	2290	II. 232		23 52 17.4	3.058	1	70 0 13.5	20.05	1	pF; S; R; sbM; * 10 sp.....	3
5039	2291	II. 10		33 52 28.1	3.063	1	75 58 24.5	20.05	1	F; pS; iE 15° ±	3
5040	2292			23 53 17.8	3.033	1	40 3 38.5	20.05	1	Cl; pRi; pC; st 9...	1
5041	2293			23 53 49.6	3.069	1	84 32 7.5	20.05	1	vF; S; R; psbM	1
5042	2294	III. 855		23 54 14.4	3.056	2	59 20 47.5	20.05	3	eF; S; R; sbM; stellar sp of 2	4
5043	2295	III. 856		23 54 17.0	3.056	3	59 20 0.5	20.05	3	eF; S; R; stellar; nf of 2 ...	4
5044	2296	III. 984		23 55 10.4	3.067	2	77 48 0.5	20.05	3	cF; stellar; Δ with 2 st	4*
5045	4010			23 55 43.1	3.085	1	125 1 38.5	20.05	1	vF; S; R; am st	1
5046	2297	II. 240		23 56 4.5	3.067	1	74 37 31.5	20.05	1	cB; cL; ir; vgbM	2†
5047	2298	III. 436		23 56 32.1	+3.070	1::	83 25 16.5	-20.05	1	vF; pL; R; lbM	2

No. of Catalogue.	References to			Right Ascension for 1860, Jan. 0.	Annual Precession in Right Ascension for 1880.	No. of Obs. used.	North Polar Distance for 1860, Jan. 0.	Annual Precession in N.P.D. for 1880.	No. of Obs. used.	Summary Description from a Comparison of all the Observations, Remarks, &c.	Total No. of times of Obs. by h. and H.
	Sir J. H.'s Catalogues of Nebulæ.	Sir W. H.'s Classes and Nos.	Other Authorities.								
	h.	II.		h m s	s						
5048	2299			23 56 39.9	+3.070	1	83 17 43.5	-20.05	1	vF; pL; R; gbM.....	1
5049	2300	II. 227		23 56 48.7	3.067	1	70 0 3.5	20.05	1	pF; cL; mE 45°+; lbM.....	4
5050	2301			23 57 19.7	3.072	1	85 34 38.5	20.05	1	pB; vS; mE; vsmbM.....	2
5051	2302			23 57 32.6	3.051	1	22 6 18.2	20.06	1	l; eeF; eeL.....	1*
5052	4011			23 57 35.2	3.087	2	152 50 42.2	20.06	2	F; S; R; gbM.....	2
5053	2303			23 57 55.7	3.072	1	83 51 42.2	20.06	1	pF; S; R; * 9 np.....	1
5054	2304			23 57 56.2	3.072	1	85 34 12.2	20.06	1	vF; S; gbM.....	1
5055	2305	VIII. 29		23 58 1.9	3.070	1	111 29 37.2	20.06	1	Cl; vP; vC.....	2
5056	2306			23 58 17.7	3.072	1	85 33 47.2	20.06	1	vF; S; R; * nf.....	1
5057	4013	III. 190		23 59 17.4	+3.072	1	94 29 48.2	-20.06	1	vF; vS; R; vg, psmbM; 2 st 9 sf.	2

SUPPLEMENTARY LIST OF NEBULÆ AND CLUSTERS.


5058			G. P. Bond	0 35 1.0	+3.073	...	89 50 6.6	-19.80	...	F; S; R; * 11 sp 1'; disc Sept. 16, 1862.	0
5059	79, b		R. nova	0 50 7.9	3.241	...	60 23 27	19.56	...	No description; γ in Lord R.'s diagram.	0
5060			S. Coolidge	3 6 55.2	3.089	...	89 4 27.5	13.67	...	F; disc Jan 25, 1860.....	0
5061			S. Coolidge	3 16 29.0	+3.086	...	89 19 3.0	13.07	...	F; disc Dec. 16, 1859.....	0
5062	(123)			4 57 18.4	-0.385	1	159 36 32.1	-5.43	1	No description.....	1
5063	(374)			5 20 52.9	-0.469	1::	159 36 35.4	-3.42	1::	No description.....	1
5064			J. H. Safford	6 5 7 :	+3.093	...	88 50 39 :	+0.43	...	2 Cls; near 2 st 9.10 & 10.11; disc Mar. 19, 1863.	0
5065			J. H. Safford	6 6 40.9	3.093	...	88 58 11.2	0.44	...	Cl; bet 2 st 9.10 & 10.11; disc Mar. 19, 1863.	0
5066			G. P. Bond	7 55 12.5	3.509	...	65 25 19.4	9.73	...	vF; cometic; disc Sept. 1, 1852.	0
5067			S. Coolidge	9 59 51.1	3.079	...	89 15 7.3	17.38	...	Neb; no description; disc Mar. 31, 1859.	0
5068			S. Coolidge	10 16 14.1	3.079	...	89 13 46.0	18.05	...	F; disc Mar. 31, 1859.....	0
5069	939, c		R. nova	11 33 +	3.122	...	71 29 +	19.93	...	No description.....	0
5070	2849, a		D'Arrest	12 12 41	3.06	...	83 46 42	20.02	...	A nebula; no description.....	0*
5071			S. Coolidge	12 31 0.5	3.069	...	89 2 46.5	19.87	...	* 12, in F neb; disc May 3, 1859.	0
5072			S. Coolidge	13 24 32.0	3.067	...	89 18 29.7	18.67	...	* 12, in F neb; disc Apr. 30, 1859.	0
5073			S. Coolidge	13 42 38.3	3.065	...	89 14 2.4	18.03	...	* 12, in Fneb; disc Apr. 30, 1859.	0
5074			G. P. Bond	13 49 18.6	3.068	...	89 31 9.0	17.77	...	S; R; * 92'; disc June 8, 1855.	0
5075			S. Coolidge	13 53 58.0	3.065	...	89 13 53.3	+17.60	...	* 12, in neb; disc Apr. 29, 1859.	0
5076			Lac. I. 11	18 20 44.6	3.952	...	123 30 27.3	-1.89	...	Neb. without stars.....	0
5077			G. P. Bond	21 45 12.5	2.217	...	40 53 46.4	16.67	...	Neb; no description; disc Feb. 10, 1848.	0
5078	3943, a		Lassell	22 20 50 :	3.336	...	115 34 +	18.25	...	Neb* * 1' dist from h. 3943...	0
5079			G. P. Bond	23 25 33.1	+3.093	...	96 22 12.0	-19.83	...	Neb; * 9.10 sf; disc Oct. 23, 1848.	0

Of this supplementary list, the objects Nos. 5058, 5060, 5061, 5064, 5065, 5066, 5067, 5068, 5071, 5072, 5073, 5074, 5075, 5077, and 5079 were communicated to me by Professor BOND, Director of the Observatory of Harvard College, U.S., too late for insertion in the body of the Catalogue.

ERRATA.

In page 7, lines 13, 14, for 5063 read 5079, and for six read 22.





U. C. BERKELEY LIBRARIES



C054582036

